



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*



## Evaluation of quality and variety of Indonesian cassava (*Manihot esculenta* Crantz)

Amarullah

Received 28 April 2020, Revised 19 May 2020, Accepted 20 June 2020, Published online 30 June 2020

### ABSTRACT

The evaluating cassava varieties for productivity necessary to assess the cassava stem characteristics for their suitability as planting material and to improve the capability of cassava farmers to select good quality cassava varieties. Cassava varieties are generally distinguished from each other by their morphological characteristics, which include leaf, stem and tuber colour. The cassava planting material used in this study and some of characteristics by five cassava varieties, consisted of mature stem cuttings of about 20 cm in length, containing between 10 and 12 nodes and planted in a vertical position along the top of the ridges. Each plot consisted of 30 plants, with data being taken from ten plants within each plot. Cassava varieties Malang-6 presented the highest yield average is 13.81 tones ha<sup>-1</sup> followed by Singgah and Adira-4 i.e. 11.98 t ha<sup>-1</sup> and 11.11 t ha<sup>-1</sup> of different varieties 1 yang Ketan, which only weighs 6.63 t ha<sup>-1</sup>. Harvest index varieties of Adira-4, Malang-6, UJ 5 and Singgah were found 0.78, 0.77, 0.77 and 0.76, respectively higher and significantly different IP varieties of glutinous Ketan is 0.58. Malang varieties and Malang-6 that produces the highest tuber weight, has a low dry matter content is 5.65% and 5.62%. The Ketan varieties and UJ-5 has a tuber weight was significantly lower with dry matter content higher than that dried 8.69% and 8.68%. The UJ-5 varieties has a tuber weight starch HCN 230,17 higher and significantly other variety.

**Keywords:** Evaluation, Quality, Cassava variety.

Department of Agronomy, Faculty of Agriculture, University of Borneo Tarakan, Jl. Amal Lama 1, Tarakan. North Borneo, Indonesia.

\*Corresponding author's email: [amarullah70@gmail.com](mailto:amarullah70@gmail.com) (Amarullah)

Cite this article as: Amarullah. 2020. Evaluation of quality and variety of Indonesian cassava (*Manihot esculenta* Crantz). *Int. J. Agril. Res. Innov. Tech.* 10(1): 108-116. <https://doi.org/10.3329/ijarit.v10i1.48102>

### Introduction

Cassava (*Manihot esculenta* Crantz) is one of the most important crops in the tropics (Nnodu *et al.*, 2006; Ikeh *et al.*, 2012) and subtropics around the world (Raji *et al.*, 2009; Perera *et al.*, 2013), globally, cassava plant is the sixth most important after wheat, rice, potatoes, corn and barley, and became a staple food for more than 800 million people in the world, especially in tropical countries (Lebot, 2009; Ladeira *et al.*, 2013) while in Indonesia, cassava is the third staple food after rice and maize (Ristono and Amarullah, 2011).

Prospects for the development of cassava commodities continued to increase in line with the increasing diversification of needs for raw materials are sourced from cassava. In some countries, cassava is the staple food and some are used for processed into food alternatives. The economic value of cassava can be enhanced when used as biofuels, especially with the depletion of world oil reserves that make cassava demand is

increasing in some other countries (Amarullah *et al.*, 2016).

According to Alexandratos (1995), cassava plays an important role in alleviating food problems, because it thrives and produces stable yields under conditions in which other crops fail. Cassava is a versatile crop and can be processed into a wide range of products such as starch, flour, tapioca, beverages and cassava chips for animal feed. Cassava is also gaining prominence as an important crop for the emerging biofuel industry and, as corroborated by Ziska *et al.* (2009), is a potential carbohydrate source for ethanol production. A well planned strategy for the development and utilization of cassava and cassava products can provide incentives for farmers, crop vendors and food processors to increase their incomes. It can also provide food security for households producing, consuming cassava products (Plucknett *et al.*, 1998).

Traditionally, cassava has been grown by farmers throughout the Indonesian and has been of particular importance to small farmers, where it is still cultivated. It is a crop that is generally grown on marginal lands with a minimum of agricultural 2 inputs (Hillocks *et al.*, 2002). Once established, the cassava crop is given little attention, but still is able to tolerate weed competition, as well as insect pests and diseases. The potential exists for improving the productivity of cassava through better agronomic practices, superior varieties and pest and disease management.

Cassava varieties are generally distinguished from each other by their morphological characteristics, which include leaf, stem and tuber colour, leaf shape and number of storage roots per plant. The plant produces all year round and can be harvested over an extended period. Although it is easily propagated by stem cuttings, the lack of quality planting material is a major constraint to the development of a viable cassava production system. Eze and Ugwuoke (2010) report that tuber yield of cassava is influenced by both the quality of planting material used and the agronomic practices employed. Access to high yielding cassava varieties and an improvement of the production system will result in increased economic benefits to local farmers.

Among the objectives of the seed and tuber crops programme by Balitkabi at the Malang Indonesian are to identify high yielding cassava varieties, to evaluate and preserve cassava germplasm and to provide good quality planting material for local farmers. Documented information on the performance of cassava on the inceptisol soils of Java, under improved agronomic practices, is lacking. Until this present study, very little effort had been made to evaluate existing varieties and new introductions for their yield potential under local conditions. Consequently, there is a need for conducting

research on cassava in order to identify varieties suitable for growing on Java soils. The productivity of this crop can be improved through the selection of cassava varieties are most suitable for local conditions and the application of the appropriate agronomic practices

### Objectives

The objectives of this study were to evaluate five cassava varieties for their productivity on Java soils, to assess the cassava stem characteristics for their suitability as planting material and to improve the capability of local farmers to select good quality cassava varieties.

### Materials and Methods

#### Experiment site

The study was carried out at the experiment, Gajah Mada University, Yogyakarta, from February 2013 to September 2013. Research station is situated at altitude 113 m above mean sea level.

#### Experiment design

The cassava planting materials used in this study and some of characteristics by variety are listed in Table 1. These varieties are already a part of the production systems of local farmers. Though no nutritional studies have been done to determine the levels of hydrogen cyanide (HCN) within their edible portions, they have been designated by farmers as sweet cassava varieties, with apparently low HCN contents. To date, there have been no reports of HCN toxicity with any of the three varieties evaluated. Their origins could not be independently verified at the time of this study; the information found in Table 1 was obtained from the farmers who supplied the plant material, or identified it as such from stock plants growing at the Balitkabi, Malang Indonesia.

Table 1. Cassava varieties used as planting material for cassava tests.

No	Variety	Origin	Status variety	Description
1	Adira-4	Balitkabi, Malang Indonesian	VUB National	Early branching, erect
2	Malang-6	Balitkabi, Malang Indonesian	VUB National	Early branching, spreading
3	UJ-5	Balitkabi, Malang Indonesian	VUB National	Early branching, spreading , menyebar
4	Singgah	Park P3SG UBT Tarakan North Borneo	Local Variety North Borneo (Promotion)	Late branching, erect
5	Ketan	Balitkabi, Malang Indonesian	Local Variety Ponorogo East Java, Indonesian (Balance sample)	Early branching, spreading

*Balitkabi: Research Institute for Nuts and Tubers; P3SG UBT: Research and Development Center for Elephant Cassava, University of Borneo Tarakan; VUB: new high yielding varieties; Stop: Cassava Elephant; East Java: East Java.*

The five cassava varieties evaluated in this experiment were established in an open field in a block randomized design with four replications, on ridges 1.5 m apart, with a 1.0 m spacing between plants within rows. The planting material consisted of mature stem cuttings of about 20 cm in length, containing between 10 and 12 nodes and planted in a vertical position along the top of the ridges. Each plot consisted of 30 plants, with data being taken from ten plants within each plot. Tuber yield was determined from the actual area of each plot, which, according to [Romani \*et al.\* \(1993\)](#), provides a good estimate of true yield. This is also supported by [Neppel \*et al.\* \(2003\)](#) whose study indicated that

interactions of centre row with border row were insignificant.

The usual cultural practices were observed to ensure an even stand of plants in the experimental plots. The cassava trial plots were grown under rain-fed conditions. Fertilizer was applied at a rate of 250 kg per hectare (223 lb per acre) one month after planting, then again at three months after planting. Before each application of fertilizer, the plots were weeded and cultivated. Plant characteristics (Table 2) were described according to the descriptor list for cassava genetic resources ([CIAT, 2009](#)).

Table 2. Observed cassava plant characteristics and rating system ([CIAT, 2009](#))

No	Characteristics	Determination Method	Key
1	Colour of first fully expanded leaf	Observing and estimating	1. Light green, 2. Dark green, 3. Green-purple), 4. Purple
2	Shape of central leaf lobe	Observing and estimating	1. Oblanceolate, 2. Linear, 3. Elliptic, 4. Pandurate, 5. Lanceolate, 6. Other
3	Colour of unexpanded apical leaves	Observing and estimating	1. Light green, 2. Dark green, 3. Green-purple, 4. Purple, 5. Other
4	Pubescence of young Leaves	Observing and estimating	1. Sparse, 2. Intermediate, 3. Dense
5	Petiole colour	Observing and estimating	1. Light green, 2. Dark green, 3. Green-purple, 4. Purple, 5. Other
6	Stem colour	Observing and estimating	1. Silver green, 2. Light brown or orange, 3. Dark brown, 4. Other
7	Storage roots per plant	Counting	
8	Storage root pulp colour	Observing and estimating	1. White or cream, 2. Yellow, 3. Pink, 4. Other
9	Storage root surface colour	Observing and estimating	1. White, 2. Cream, 3. Light brown, 4. Dark brown, 5. Other
10	Colour of outer surface of storage root cortex	Observing and estimating	1. White, 2. Yellow, 3. Pink, 4. Purple, 5. Other

Farming practices commonly observed to ensure even stand of plants in the experimental plots. Cassava planted in experimental plots during the dry season. Urea fertilizer, SP36 and KCl was applied at a rate of 200, 100, and 100 kg ha<sup>-1</sup>, given half a dose at age 1 month after planting (bst), then the rest is given again at age 3 bst. Before the application of fertilizers, which weeded plots of weeds, while manure is given at age 1 bst with a dose of 20 t ha<sup>-1</sup>. Plant characteristics are presented in Table 2, according to a list of descriptions of cassava genetic resources ([CIAT, 2009](#)).

The minimum and maximum monthly temperature average for the trial period each 20°C and 28°C, total rainfall of 1,124 mm and the duration of sunlight monthly average for the period was 7.7 hours. Ten months after planting,

in a plot harvested cassava and tubers counted, weighed and measured. Data taken from ten plants randomly selected from each experimental plot. The parameters observed and measured include; the number of tubers per plant (grain), root length (cm), root diameter (cm), weight of each tuber (g), total tuber yield per plant (kg), and the harvest index (IP). In addition, the following trunk characteristics were evaluated: plant height (cm), number of segments, stem diameter (cm), segment length (cm) and number of primary branches. Plant height is measured from the base of the plant to the highest tops.

Samples of the tubers harvested for each of the five varieties of cassava to be analyzed HCN and starch content of the Food Science Laboratory FTP UGM. Varieties of cassava tuber samples are weighed in water, 1.5 or 3.0 kg to determine the

value of dry matter. Analysis of N, P and K of total land is analyzed in the laboratory of soil science FP UGM to determine the level of absorption and utilization of nutrients N, P and K by each variety of cassava.

### Statistical analysis

Data were analyzed by analysis of variance (ANOVA). The values with significant differences were tested further with; Duncan's multiple range test (DMRT) at 5% level of significance. The data is processed and analyzed using Excel Program continued into portable 9.1.3 SAS program (SAS, 2004).

## Results and Discussion

Five cassava varieties were evaluated for their morphological characteristics according to morphological characteristics of cassava plants (Table 3). No signs of harmful pests or diseases on the leaves, stems and roots during this trial

period were observed. During the growing season, there are many leaves are falling, and there is probably not too have adverse effects on tuber yields. Decreased level of production, leaf and root dry weight can occur at temperatures too low even reaching 17°C (Alves, 2002), but (Irikura *et al.* (1979) have found that the same cassava varieties can produce higher production at various temperatures, this shows that the effect of natural selection is very significant on the adaptation of varieties. These varieties are distinguished from one another based on their leaves, stems and tubers characteristics. Variations encountered in the characteristics of the five varieties of cassava in this study. Cassava Adira-4, Malang-6 and UJ-5 has a dark tuber surface with white flesh and a light yellow cortex, while the Transit and sticky rice varieties tend to be the same as the tuber flesh is white with white storage root surface and cortex crimson

Table 3. Morphological characteristics of cassava plant.

Parameter	Characteristics					Information
	Adira-4	Malang-6	UJ-5	Singgah	Ketan	
Observing age 3 MAP						
1. Color of Leaf Buds	1	1	2	1	1	1. Light green 2. Dark green (Hijau Gelap) 3. Purplish green 4. Purple
2. Pubertas of leaf buds	0	0	0	0	0	0. Absent 1. Present
Observing age 6 MAP						
3. Leaf retention	2	1	3	2	3	1. Very poor retention 2. Less than average retention 3. Average retention 4. Better than average retention 5. Outstanding leaf retention
4. Central leaflet	1	2	2	2	1	1. Ovoid 2. Elliptic-lanceolate
5. Petiole color	1	4	3	6	6	0. Yellowish-green, 3. Green 4. Reddish-green, 5. Greenish-red, 6. Red, 7. Purple
6. Leaf color	2	1	2	1	2	1. Light green, Dark green 2. Purple green, Purple
7. Number of leaf lobes	6,7,8	4,5,6,7,8	4,5,6	3,4,5,6,7	4,5,6,7	1. 3 leaf; 2. 4 leaf; 3. 5 leaf; 4,6 leaf; 5. 7 leaf; 6. 8 leaf; 7. 9 leaf. 8. 10 leaf; 9. 11 leaf
8. Length of leaf lobe	12-27	17-36	19-34	19-34	12-29	leaf located in the middle
9. Width of leaf lobe	1.2-4.7	2.4-8.6	2.0-7.9	2.1-7.8	1.2-6.9	leaf located in the middle
10. Ratio of lobe length to lobe width of central leaf lobe	5/1	4/1	4/1	4/1	3/1	-
11. Lobe margins	1	2	1	2	1	1. Smooth, 2. Winding
12. Petiole length	18-30	30-35	20-32	29-39	23-31	cm
13. Color of leaf vein	1	1	1	1	1	1. Green, 2. Reddish-green in less than half of the lobe 3. Reddish-green in more than half of the lobe, 4. All red
14. Orientation of petiole	1	3	4	2	2	1. Inclined upwards, 2. Horizontal, 3. Inclined downwards, 4. Irregular
15. Flowering	2	2	2	2	2	1. Absent, 2. Present
16. Pollen	2	2	2	2	2	1. Absent, 2. Present
Observing age 9 MAP						
17. Prominent of foliar scarsee	2	2	1	2	2	1. Semi-prominent 1. Prominent
18. Color of stem cortex	2	2	2	2	2	1. Orange, 2. Light green, 3. Dark green

19. Color of stem epidermis	2	2	2	2	2	1. Cream, 2. Light brown 3. Dark brown, Orange
20. Color of stem exteriors	2	7	2	3	3	1. Orange, 2. Greenish-yellowish, 3. Golden
21. Distance between leaf scars	3	3	3	3	3	1. Short (<8 cm), 2. Medium (8-15 cm), 3. Long (>15 cm)
22. Growth habit of stem	1	1	1	1	1	1. Straight, 2. Zig-zag
23. Color of end branches	1	1	2	2	2	1. Green, 2. Green-purple, 3. Purple
24. Length of stipules	1	1	1	1	1	1. Short, 2. Long
25. Stipule margin	1	1	1	1	1	1. Entire, 2. Split or forked
Observing Harvest						
26. Plant height	2.7-3.1	2.3-2.9	2.5-3.0	2.8-3.4	25.3.2	cm
27. Height to first branching	1 (180-245)	2 (50-220)	2 (10-180)	1 (70-240)	1 (40-120)	1. Primary Branch (cm) 2. Second Branch (cm)
28. Level of branching	3	2	3	3	5	
29. Branching habit	2/3	1/5	2/3	2/3	2/3/4	1. Erect, 2. Dichotomous 1. Trichotomous 2. Tetrachotomous, 5. Irregular
30. Angle of branching	70-145	60-80	50-90	45-90	70-120	O
31. Shape of plant	3	4	2	3	3	1. Compact, 2. Open 3. Umbrella, 4. Cylindrical all tubers are formed
32. Number of storage roots/plant	4-11	4-11	5-13	5-14	5-12	
33. Number of commercial roots/plant	4-10	4-11	5-9	5-14	5-12	3 tubers with a length of more than 20 cm
34. Extent of root peduncle	1	1	1	1	1	1. Sessile, 2. Pedunculate 1. Mixed
35. Root constrictions	1	1	3	1/2	1/2	1. Few to none, 2. Some, 3. Many
36. Root shape	1/2/3	3/4	1/3	2/4	4	1. Conical, 2. Conical-cylindrical 2. Cylindrical, 4. Irregular
37. External color of storage root	3	1	2	3/4	3/4	1. White or cream, 2. Yellow 3. Light brown, 4. Dark brown
38. Color of root pulp	2	2	2	1	1	1. White, 2. Cream
39. Color of root cortex	1	2	2	4	4	1. White or cream, 2. Yellow, 3. Pink, 4. Purple
40. Cortex	2	2	2	2	2	1. Easy, 2. Difficult
41. Texture of root epidermis	3	1	1	3	3	1. Smooth, 2. Intermediate, 3. Rough
42. Root taste	3	1	1	3	3	1. Sweet, 2. Intermediate, 3. Bitter
43. Cortex thickness	3	1	1	3	3	1. Thin, 2. Intermediate, 3. Thick
44. Dry matter content	1.20	0.91	1.05	1.02	0.85	3 plant
45. Starch content	23.06	27.29	29.57	25.64	28.26	3 plant
46. Harvest index	0.49	0.42	0.46	0.48	0.39	4-6 plant/varieties
47. Cyanogen potential/CNP	108.19	109.33	236.84	133.64	100.44	(mg/100g)

Analysis of the performance characteristics of stem growth are presented in Table 4, showed a significant response to the parameters of the results of the five varieties of cassava. Plant height, number of segments, stem diameter and number of primary branches are significant at 1% level of confidence, while the segment length is significant at 5% level.

Table 4. Plant height conditions, number of segments and branch conditions on five cassava varieties in one plant per m<sup>2</sup> age 9 MAP.

Varieties	Plant height (cm)	Number of segments	Stem Diameter (cm)	Number of branch	Branch level	shape and angle of the branch
Adira 4 (V <sub>1</sub> )	293.04ab	142.44b	3.20b	2.60b	1.98b	Umbrella, (35-60°)
Malang 6 (V <sub>2</sub> )	274.23b	163.90a	3.56	4.33a	1.20d	Cylindrical, lateral branches and Irregular, (35-50°)
UJ 5 (V <sub>3</sub> )	241.26c	110.43b	2.72d	1.76d	1.55c	Open, decumbent or horizontal, (65-90°)
Singgah (V <sub>4</sub> )	307.80a	120.82b	2.66cd	2.33c	1.40c	Cylindrical, (25-35°)
Ketan (V <sub>5</sub> )	295.33ab	138.27b	3.02bc	2.13c	3.27a	Compact, (50-70°)

Note: Numbers in columns followed by the same letter show no difference according to the Duncan's test at 5% level.

Analysis of variance test performed showed a significant difference between the responses for all five varieties of tuber yield. Data analysis (Table 4) shows the significance of the results of the tuber. Tuber yield is determined by the

number, length, diameter and weight as well as weight of the tuber and tuber per plant. Bulbs separated into valuable and worthless bulbs, depending on the size and weight.



Fig. 1. Results of the stems of five cassava varieties aged 4 MAP; V: Cassava varieties; V1 Adira-4; V2: Malang-6; V3: UJ-5; V4: Singgah; V5: Ketan.

Based on the cassava harvest at the age of 9 bst obtained significant differences between the varieties of cassava to the parameters of tuber yield and harvest index. Weight of five cassava tuber crops increase until at harvest age 9 bst. The harvest index (IP) for five cassava varieties has increased since the age of 3-9 years but shows different results among the varieties, where the varieties of Adira-4, Malang-6, UJ-5 and Singgah IP show higher than Ketan varieties (Table 4).

Number and length of tuber of Singgah cassava variety was found more, longer and significantly different from the four other varieties. Cassava varieties Adira-4 and Malang-6 produces very large diameter tubers, while the glutinous

varieties produce small tubers that are uniform in size. This condition may make the plant more efficient in collecting the storage of carbohydrates in the roots. Higher number of tubers in the varieties Transit UJ-5 and Ketan not in line with the results of cassava tuber weight. These results supported the research results of Hayford (2009), which indicates that there is a negative correlation between the number of tubers per plant and mean weight of cassava tubers. Weight of tubers per plant was lower in glutinous varieties allegedly because of competition between the roots of the bulbs during the charging process.

Table 4. Tuber yields and harvest index (HI) of five cassava varieties.

Varieties	Number of Tuber	Length of tuber (cm)	Diameter of tuber (cm)	Weight of tuber (g)	Tuber weight per plant (kg)	Index harvest (IP)
Adira-4 (V <sub>1</sub> )	9.11d	40.77c	9.59a	992.98b	11.11c	0.78a
Malang-6 (V <sub>2</sub> )	9.33d	64.40b	8.74b	1033.95a	13.81a	0.77a
UJ-5 (V <sub>3</sub> )	10.67b	34.87d	7.12d	777.33c	9.63d	0.77a
Singgah (V <sub>4</sub> )	11.22a	75.98a	7.75c	1053.02a	11.98b	0.76a
Ketan (V <sub>5</sub> )	10.22c	52.77b	6.09d	605.77d	6.63e	0.58b

Note: Numbers in columns followed by the same letter show no difference according to the Duncan's test at the 5% level.

The vegetative growth of the five cassava tested did not show consistency and direct relationship between growth parameters, but from the data in Table 4. It was found that cassava varieties with medium plant height produced a higher number of segments and a number of branches such as those in Adira-4 and Malang-6. This condition is similar to the results of the study of [Ntui et al. \(2006\)](#), which states that cassava varieties with stems that continue to produce longer segments, fewer internodes and fewer tubers per plant.

Cassava varieties Singgah produces a higher number of tubers, tuber length and weight per tuber and is different from other varieties. The cassava Malang-6 variety has a larger tuber diameter and produces a higher weight per tuber and plantations and is different from other

varieties. Whereas, the sticky cassava variety which has the smallest number of tubers and diameter produces the lowest weight per tuber or plant and is different from other varieties. Cassava with a high tuber yield in terms of number, diameter and high tuber weight also results in a high Harvest Index. Proven on the sticky cassava varieties produce the smallest IP of 0.58. The difference in cassava tuber yield is determined by several factors, such as the number of tubers, tuber length and tuber weight. [Ntawuruhunga and Dixon \(2010\)](#) concluded that the number and size of tuber diameters were components of the main yield that contributed to the increase of cassava



Fig. 2. Tuber yields of five cassava varieties aged 4 bst; V: Cassava varieties; V1 Adira-4; V2: Malang-6; V3: UJ-5; V4: Singgah; V5: Ketan.

Dry matter content of cassava is affected by a number of factors such as the age of the plant, season, and various plant locations. Dry matter content of cassava varieties in the range of high value, as reported by [Chávez et al. \(2005\)](#), ranging from 10.7% to 57.2%, with an average of 34.7%. Various varieties that produce the highest tuber weight, has the lowest dry matter content. There are also some varieties that produce significantly lower weighty tuber with a high dry matter. [Ntawuruhunga et al. \(2001\)](#), and recently [Adeniji et al. \(2011\)](#), reported the results of a

similar study on the relationship between the dry matter content and weight of tubers. Dry matter content was significantly negatively correlated with storage root weight. This shows that when the weight of high storage root studied, the dry matter content tends to be low.

Cassava is consumed by taking into account the levels of cyanide and starch in the tuber. There are significant differences between the levels of cyanide and starch of cassava varieties, as presented in Table 5.

Table 5. Dry matter, HCN and starch content of cassava variety.

No	Variety	Dry matter (%)	Content of HCN (mg kg <sup>-1</sup> )	Content of starch (%)
1	Adira-4 (V <sub>1</sub> )	8.59d	111.52c	24.32d
2	Malang-6 (V <sub>2</sub> )	8.65b	112.66c	28.23b
3	UJ-5 (V <sub>3</sub> )	8.68a	230.17a	30.26a
4	Singgah (V <sub>4</sub> )	8.62c	136.97b	26.15c
5	Ketan (V <sub>5</sub> )	8.69a	103.77c	28.85b

Note: HCN: Cianida acid, Numbers in columns followed by the same letter show no difference according to the Duncan's test at the 5% level.

UJ-5 cassava varieties contain the highest HCN, starch content and significantly different from other varieties. The five cassava varieties containing HCN above 100 mg kg<sup>-1</sup> means that including cassava is rather bitter. [Sundaresan et al. \(1987\)](#) classified cassava into three categories based on HCN content, namely non-bitter cassava (HCN <100 mg kg<sup>-1</sup> of fresh cassava), bitter cassava (HCN between 100-450 mg kg<sup>-1</sup> of

fresh cassava), and very bitter cassava (HCN > 450 mg kg<sup>-1</sup> of fresh sweet potatoes).

Cassava is a plant that is ideal for integrating into local farmers' crop production systems. This study shows that the three varieties can effectively compete under the conditions of the soil and the environment. Different varieties of cassava will give farmers the opportunity to choose the most suitable cassava for its growing conditions. This study shows that there is great

variability in the morphological characteristics of the roots and stems of cassava varieties. Morphological characteristics presented in this study will assist farmers in identifying these varieties. Tuber yield can be improved through selection based on components of the results, in particular, plant height, number of bulbs and bulb size (Aina *et al.*, 2007; Akinwale *et al.*, 2010). Selection of cassava varieties with good plant height and good branching habit, will facilitate the implementation of planting and harvesting, and provide quality planting material at the end of the harvest.

Characteristics of cassava tubers play an important role in the recommendation and acceptance of cassava varieties to farmers and consumers. Cassava development programs should be taken into account not only for the performance results of the bulbs, but also must take into account the characteristics of quality. Quality characteristics may be more difficult to assess objectively, however, because they relate to features such as appearance, color, taste and texture.

The results showed differences in the response of cassava varieties with the measured parameter, an indication that these properties depend on the variety. It is therefore important that cassava varieties were evaluated to assess its potential. Additional research is also needed, with respect to the feasibility of cassava production. Introduction and evaluation of new cassava varieties could increase the productivity of local farmers through increased yields for consumption bulbs and stems for planting material, with a minimum of manpower and financial input.

## Conclusion

Cassava varieties Malang-6 has a rod with highest average number of segments (163.90), followed by 142.44 Adira-4 as much, so it will be very much beneficial if used as source of planting materials. Cassava varieties Malang-6 presented the highest yield average is 13.81 tones ha<sup>-1</sup> followed by Singgah and Adira-4 i.e. 11.98 and 11.11 tons ha<sup>-1</sup> but the variety 1 yang Ketan which only weighs 6.63 tons ha<sup>-1</sup>. Harvest index of varieties Adira-4, Malang-6, UJ 5 and Singgah were 0.78, 0.77, 0.77 and 0.76, respectively and the values were higher and significantly form different IP varieties of Ketan is 0.58. Malang-6 varieties and Singgah that produces the highest tuber weight, has a low dry matter content is 8.65% and 8.62%. The Ketan varieties and UJ-5 has a tuber weight was significantly lower with dry matter content higher than that dried 8.69% and 8.68%. Dry matter content was significantly negatively correlated with tuber weight, showed

that when high tuber weight, then dry matter content tends to be low.

## References

- Adeniji, O.T., Odo, P.E. and Ibrahim, B. 2011. Genetic relationships and selection indices for cassava root yield in Adamawa State, Nigeria. *African J. Agril. Res.* 6(13): 2931-2934.
- Aina, O.O., Dixon, A.G.O. and Akinrinde, E.A. 2007. Genetic variability in cassava as it influences storage root yield in Nigeria. *J. Biol. Sci.* 7: 765-770. <https://doi.org/10.3923/jbs.2007.765.770>
- Akinwale, M.G., Akinyele, B.O., Dixon, A.G.O. and Odiyi, A.C. 2010. Genetic variability among forty-three cassava genotypes in three agro-ecological zones of Nigeria. *J. Plant Breed. Crop Sci.* 2(5): 104-109.
- Alexandratos, N. ed. 1995. World Agriculture: Towards 2010: A FAO Study, Chichester, England. Food and Agriculture Organization of the United Nations; John Willey and Sons. pp. xxvi + 488.
- Alves, A.A.C. 2002. Cassava botany and physiology. In: Hillocks, R.J., Thresh, J.M. and Bellotti, A.C. (Eds.). Cassava: biology, production and utilization. UK: CABI Publishing. pp. 67-89. <https://doi.org/10.1079/9780851995243.0067>
- Amarullah, Indradewa, D., Yudono, P. and Sunarminto, B.H. 2016. Photosynthetic activity of superior varieties super and local Cassava (*Manihot esculenta* Crantz) in Indonesia. *J. Agril. Sci.* 8(8): 194-200. <https://doi.org/10.5539/jas.v8n8p194>
- Chávez, A.L., Sánchez, T., Jaramillo, G., Bedoya, J.M., Echeverry, J., Bolaños, E.A., Ceballos, H. and Iglesias, C.A. 2005. Variation of quality traits in cassava roots evaluated in landraces and improved clones. *Euphytica*. 143: 125-133. <https://doi.org/10.1007/s10681-005-3057-2>
- CIAT. 2009. Key access and utilization descriptors for cassava genetic resources. Bioversity International, Rome, Italy, International Center for Tropical Agriculture (CIAT), Cali, Colombia. 5p.
- Eze, S.C. and Ugwuoke, K.I. 2010. Evaluation of different stem portions of cassava (*Manihot esculenta*) in the management of its establishment and yield. *Res. J. Agric. Biol. Sci.* 6(2): 181-185.
- Hayford, M.A. 2009. Growth, Yield and Quality of Cassava as Influenced by Terramend 21, Poultry Manure and Inorganic Fertilizer. Master of Agronomy Thesis, School of Graduate Studies, Kwame Nkrumah University of Science and Technology (KNUST), Kumasi, Ghana. 66p.

- Hillocks, R.J., Thresh, J.M. and Bellotti, A.C. ed. 2002. Cassava: biology, production and utilization. New York: CABI Publishing. 332p. <https://doi.org/10.1079/9780851995243.0000>
- Ikeh. A.O., Udaeyo. N.U., Udoh. E.I., Iboko.K.O. and Udounang. P.I. 2012. Growth and yield of Cassava (*Manihot esculenta* Crants) as influenced by the number of shoots retained per stand on as Ultisol. *Nat. Sci.*10(8): 17-20. <http://www.sciencepub.net/nature>
- Irikura, Y., Cock, J.H. and Kawano, K. 1979. The physiological basis of genotype-temperature interactions in cassava. *Field Crops Res.* 2: 227-239. [https://doi.org/10.1016/0378-4290\(79\)90025-X](https://doi.org/10.1016/0378-4290(79)90025-X)
- Ladeira, T., Souza, H. and Pena, R. 2013. Characterization of the roots and starches of three cassava cultivars. *Int. J. Agril. Sci. Res.* 2(1): 12-20.
- Lebot, V. 2009. Tropical roots and tuber crops: cassava, sweet potato, yams and aroids. Crop Production science in horticulture series. 17, CABI Publishing, UK. pp. 50-103. <https://doi.org/10.1079/9781845934248.0000>
- Neppel, G.P., Wehner, T.C. and Schultheis, J.R. 2003. Interaction of border and center rows of multiple row plants in watermelon yield trials. *Euphytica.* 131: 225-234. <https://doi.org/10.1023/A:1023958321626>
- Nnodu, E.C., Ezulike, T.O. and Asumugha, G.N. 2006. Cassava. In: Idem NUA, Showemimo FA (eds.). Tuber and Fibre Crops of Nigeria: Principles of production and Utilization. 2006: (XXII), 239p.
- Ntawuruhunga, P. and Dixon, A. 2010. Quantitative variation and interrelationship between factors influencing cassava yield. *J. Appl. Biosci.* 26: 1594-1602.
- Ntawuruhunga, P., Rubaihaho, P.R., Whyte, J.B. A., Dixon, A.G.O. and Osiru, D.S.O. 2001. Inter-relationships among traits and path analysis for yield components of cassava: A search for storage root yield indicators. *African Crop Sci. J.* 9: 599-606. <https://doi.org/10.4314/acsj.v9i4.27582>
- Ntui, V.O., Uyoh, E.A., Affangideh, U., Udensi, U. and Egbonyi, J.P. 2006. Correlation and genetic variability in cassava (*Manihot esculenta* Crantz). *J. Food Agric. Environ.* 4(3/4): 147-150.
- Perera, P.I., Quintero, M., Dedicova, B., Kularatne, J.D. and Ceballos, H. 2013. Comparative morphology, biology and histology of reproductive development in three lines of *Manihot esculenta* Crantz (Euphorbiaceae: Crotonoideae). *AoB Plants.* 5: plso46. <https://doi.org/10.1093/aobpla/plso46>
- Plucknett, D.L., Phillips, T.P. and Kagbo, R.B. 1998. A Global Development Strategy for Cassava: Transforming a Traditional Tropical Root Crop: Sparing Rural Industrial Development and Raising Incomes for the Rural Poor. Available online: <http://www.fao.org/docrep/006/y0169e/y0169e04.htm> (accessed 9 August 2011).
- Raji, A.A.J., Anderson, J.V., Kolade, O.A., Ugwu, C.D., Dixon, A.G.O. and Ingelbrecht, I.L. 2009. Gene-based microsatellites for cassava (*Manihot esculenta* Crantz): prevalence, polymorphisms, and cross-taxa utility. *BMC Plant Biol.* 9: 118. <https://doi.org/10.1186/1471-2229-9-118>
- Ristono and Amarullah. 2011. Singkong Gajah Berjuang (*Indonesian*). Petrogas Press. Cetakan II. Balikpapan. 202p.
- Romani, M., Borghi, B., Alberici, R., Delogu, G., Hesselbach, J. and Salamini, F. 1993. Intergenotypic competition and border effect in bread wheat and barley. *Euphytica.* 69: 19-31. <https://doi.org/10.1007/BF00021722>
- SAS. 2004. *Statistical Analysis System for Windows* 9.0. SAS Institute Inc. SAS Campus Drive. Cary. North Carolina. USA. 5121 p.
- Sundaresan, S., Nambisan, B. and Eswari, A. 1987. Bitterness in cassava in relation to cyanoglucoside content. *Indian J. Agric. Sci.* 57: 37-40.
- Ziska, L., Runion, G.B., Tomecek, M., Prior, S.A., Torbet, H.A. and Sicher, R. 2009. An evaluation of cassava, sweet potato and field corn as potential carbohydrate sources for bioethanol production in Alabama and Maryland. *Biomass and Bioenergy.* 33: 1503-1508. <https://doi.org/10.1016/j.biombioe.2009.07.014>