AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

**Health Effects of Sustained Exposure to Fine Particulate Matter:
Evidence from India ^a**

Yashaswini Saraswat ^{1,2} & Sangeeta Bansal ^{1,3}

*Selected Paper prepared for presentation at the 2020 Agricultural & Applied Economics
Association Annual Meeting, Kansas City, MO
July 26-28, 2020*

^a **JEL Classification:** I12, I15, Q51, Q53, Q56

¹ Centre for International Trade and Development, Jawaharlal Nehru University, Delhi-67

² E-mail: yashaswini.saraswat@gmail.com

³ E-mail: sangeeta.bansal7@gmail.com

Copyright 2020 by [authors]. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

ABSTRACT

India suffers from severe air pollution. Nine out of the ten most polluted cities in the world lie here. Evidence is lacking on the effect of air pollution on health outcomes in India. The objective of this study is to estimate the causal effect of long-term exposure to PM_{2.5} on health outcomes in India, viz., early life mortality (neo-natal, infant, under-five) and life expectancy at birth. We utilise the “valley effect” owing to the peculiar topography and location of the Indo-Gangetic Plains as a basis for a natural experiment. The lower boundary of the Indo-Gangetic Plains drawn from the geographical literature acts as the exogenous threshold for regression discontinuity design (RDD). Applying RDD and two-stage least square method, the study provides the first causal estimates of the health impact of long-term exposure to PM_{2.5} using health and pollution data from India. We find PM_{2.5} exposure is 49% higher and life expectancy is 2.6 years lower in the Plains relative to other districts in the sample. Early life mortality is positively and significantly affected by sustained exposure to PM_{2.5}. Life expectancy at birth reduces by 1.2 years due to additional 10 µg/m³ of PM_{2.5} exposure, *ceteris paribus*. Around 5.2 and 8.8 years of life can be saved in the Indo-Gangetic Plains if the national and the WHO standards for PM_{2.5} are met in the region, respectively. For India as a whole, these figures are 1.7 and 5.3 years. India needs immediate policy attention to curb air pollution.

Keywords: Early life mortality, Fine particulate matter, India, Life expectancy, Regression discontinuity design, Sustained exposure.

INTRODUCTION

Air pollution has reached dangerously high levels in India. More than 99% of the Indian population is exposed to PM_{2.5} concentration levels exceeding the World Health Organization standard of 10 µg/m³ (Apte and Pant, 2019). Higher incidence of cardiorespiratory diseases like heart stroke, lung cancer, asthma has been observed in heavily polluted regions in the world (WHO, 2016). Studies have also found early life mortality to rise and life expectancy to fall as a result of exposure to PM_{2.5} (Pope et al., 2002; Tanaka, 2015; Heft-Neal et al., 2018). Ambient air pollution hinders the natural growth process in the early life stages (Zhang et al., 2018; Lavigne et al., 2018) and may even lead to a modification in the DNA of the population (Carre et al., 2017).

Several studies find a robust relationship between particulate matter and health in different countries. Chay and Greenstone (2003) have utilised a regression discontinuity design (RDD) to study the impact of total suspended particulate (TSP) matter on infant mortality rate (IMR) for US counties during 1971-72. They find a fall in infant mortality rate by 0.5% due to a 1% decline in TSP concentration. Arceo et al. (2016) find a rise in IMR by 8.8% due to a rise in PM₁₀ by 10 µg/m³ in Mexico city. Heft-Neal et al. (2018) find an IMR effect of 9.2% due to a rise in PM_{2.5} by 10 µg/m³ in Africa. Greenstone and Hanna (2014) conducted an analysis of the impact of environmental regulations targeting air and water pollution on infant mortality rate in India. They find that though the regulations improved air quality, it did not have a statistically significant impact on infant mortality rate. Tanaka (2015) finds a significant decline of 20% in infant mortality rate in China due to the first large-scale environmental regulations in the country.

Many studies find that a reduction in long-term exposure to particulate matter improves life expectancy. Pope et al. (2002) find a gain of 0.7 years due to $10 \mu\text{g}/\text{m}^3$ fall in the long term exposure to $\text{PM}_{2.5}$ in the United States. Chen et al. (2013) and Ebenstein et al. (2017) find gains in life expectancy to be 0.3 years and 0.64 years in China due to $10 \mu\text{g}/\text{m}^3$ decline in the long term exposure to TSP and PM_{10} , respectively. Lelieveld et al. (2020) utilise the Global Exposure Mortality Model (GEMM) derived from several studies to estimate loss of life expectancy (LLE) in different regions of the world. They find LLE of 3.3 years attributed to exposure to $\text{PM}_{2.5}$ and ozone in South Asia in 2015.

Most of these studies are conducted either in the developed countries that are exposed to modest levels of particulate matter or countries like China that are behaviourally and socioeconomically different from India; not many examine effects of $\text{PM}_{2.5}$ on health.

Moreover, no such long-term exposure study on health effects of $\text{PM}_{2.5}$ that utilises data from Indian population exists. Such a study is immensely important for environmental policy making in India where the emission standards are still too high and a systematic tackling of the air pollution problem is urgently required.

More recently, Balakrishnan et al. (2019) estimate the loss of life expectancy due to exposure to ambient particulate matter $\text{PM}_{2.5}$ in India. The study conducts a state-level analysis and find that if exposure to $\text{PM}_{2.5}$ was at the theoretical minimum risk level, then average life expectancy in India would have risen by 1.7 years in 2017. This improvement would have exceeded 2 years in North Indian states. They utilise estimates from integrated exposure-response functions based on several observational studies and randomized control trials conducted in different populations.

The objective of this study is to estimate the causal effect of long-term exposure to $\text{PM}_{2.5}$ on health outcomes in India. More specifically, we examine the impact of sustained exposure to $\text{PM}_{2.5}$ on early life mortality (neo-natal mortality rate, infant mortality rate and under-five

mortality rate) and life expectancy at birth in India. Evidence on such effects is important because recent scientific studies have found that the small size of $PM_{2.5}$ enables it to enter the blood stream and placenta of pregnant women, thereby affecting the health status of their offspring (Zhang et al., 2018; Lavigne et al., 2018). The evidence will facilitate better understanding of the benefits of reducing the concentration levels of $PM_{2.5}$ in India and other countries that face similar levels of air pollution.

We identify a physical boundary from the geographical literature and use it as an exogenous threshold to facilitate a regression discontinuity design (RDD). According to the Plate Tectonic Theory, collision of the Indian subcontinent with the Eurasian continent led to the origin of the Himalayan Ranges and a deep depression on their south (Burrard, 1915; Aitchison and Davis, 2007). This depression, known as the Indo-Gangetic Plains (IGP), is one of the most populated as well as polluted regions of the world. The peculiar topography of the region being sunken and landlocked by the Himalayas on the North and the Central Highlands on its South restricts the wind passage thereby making displacement of the particulate matter generated in the region difficult (Guttikunda and Gurjar, 2012). Districts lying south of the Plains do not suffer from this unfavorable “valley effect” and hence are exposed to much lower pollution levels.

For the study, we consider districts in the Indo-Gangetic Plains and districts that lie below the plains, but within a distance of approximately five degree latitude. The lower boundary of the Indo-Gangetic Plains (Fig. 1) exogenously divides this sample into treatment and control group such that the districts in the Indo-Gangetic Plains form the treatment group while the other districts form the control group. The districts to the north and the south of the boundary are similar in several ways and some of them even lie in the same state. Differences in

ecology, degree of urbanization, and socioeconomic variables may influence health indicators such as life expectancy and early life mortality. We check for the validity of a regression discontinuity design by using predicted life expectancy as a proxy variable for the health impact of these factors (Chen et al., 2013; Ebenstein et al. 2017). We find that the difference in predicted life expectancy is insignificant at the boundary. This helps in examining a causal relationship between human health and air pollution. Moreover, we control for variables like income, literacy rate, share of rural households, share of minority population, share of households with access to clean drinking water and clean cooking fuel in our regression equations.

Our findings are as follows. The unfavorable location and topography of the Indo-Gangetic Plains has severely affected the pollution concentrations in the region and the health of its inhabitants. The $PM_{2.5}$ exposure is about $23 \mu g/m^3$ or 49% higher, and life expectancy is 2.6 years lower in the region as compared to the control group. The early life mortality rates are also significantly higher in the region suggesting that when individuals are exposed to high levels of pollution for a long period of time, the probability of survival of their future offspring may reduce considerably. Further, life expectancy at birth reduces by 1.2 years due to additional $10 \mu g/m^3$ of $PM_{2.5}$ exposure, *ceteris paribus*. Hence, India can raise life expectancy on average by 1.7 and 5.3 years if the national and WHO standards for $PM_{2.5}$ are met, respectively.

This study addresses several gaps in the literature assessing the health impact of air pollution. First, the study estimates the impact of sustained exposure to $PM_{2.5}$ on life expectancy and early life mortality in India by utilizing differences in long run exposure to $PM_{2.5}$ in the Indo-Gangetic Plains (IGP) and the districts below it. The studies from India have estimated health

effects of short-term exposure to particulate matter (Guttikunda and Goel, 2013; Chowdhury and Dey, 2016). These studies often underestimate the loss in life expectancy since they only capture deaths of the vulnerable population such as old and sick that are accelerated due to a sudden rise in air pollution (Lvovsky, 1998).

Second, this is the first study that attempts to estimate the causal effect of long-term exposure to PM_{2.5} on life expectancy and early life mortality by utilizing data from India. The evidence for India comes from studies that have utilized coefficients estimated in other countries, and/or on other pollutants to extrapolate health effects of PM_{2.5} exposure in India. Greenstone et al. (2015) have utilized estimates from a study conducted in China by Chen et al. (2013) to extrapolate life expectancy gains in India of 3.2 years due to a reduction in PM_{2.5} levels to the annual Indian National Ambient Air Quality Standard of 40 µg/m³. Similarly, Greenstone and Fan (2018) have used estimates from the study by Ebenstein et al. (2017) to find life expectancy gains of 1.8 years on achieving the Indian standards for PM_{2.5}. Further, the studies by Chen et al. (2013) and Ebenstein et al. (2017) utilize data on TSP and PM₁₀ exposure, respectively, to estimate their impact on mortality and life expectancy in China. Greenstone et al. (2015) and Greenstone and Fan (2018) use these estimates to extrapolate the health impact of PM_{2.5} exposure in India by using conversion ratios. Arguably, these extrapolated estimates may not correctly reflect the health effects of improvements in air quality in India. Thus it is important to estimate the impact of improvement in air quality on the health status of the Indian population using data from India.

A novelty of the study is to use the lower boundary of IGP as an exogenous threshold for the regression discontinuity design thereby advancing the literature from observational studies. It further establishes the causal relationship by using two-stage least square method. Fourth,

most studies have dealt with the mega-cities in India (Cropper et al., 1997; Shah and Nagpal, 1997a; Kandlikar and Ramachandran, 2000; Nema and Goyal, 2010; WHO, 2018). Although nine of the ten most polluted cities in the world lie in the IGP (Guttikunda and Jawahar, 2012), there has been limited literature covering the entire region. The Indo-Gangetic Plains are the main focus of this study. Fifth, the study fills the deficit in the literature on the health impact of PM_{2.5} exposure in India. PM_{2.5} is much more harmful to health than other pollutants due to its easy penetration in lungs (since the diameter of PM_{2.5} is below 2.5 µm).

METHODS

We adopt three main approaches - the Ordinary Least Squares method (OLS), the Regression Discontinuity Design (RDD) and the Two-Stage Least Squares method (2SLS). Under the OLS, the four health outcomes are regressed on PM_{2.5} exposure as shown in the equation below:

$$Y_j = \alpha_0 + \alpha_1 PM_j + X_j\phi + e_j \quad (1)$$

where j represents a district in the sample. Y_j indicates health outcome in district j , which could be life expectancy at birth or an early-life mortality rate. PM_j is the level of fine particulate matter exposure in district j over a period of 17 years, from 1998 to 2014. X_j is a vector of demographic covariates in a district which are likely to affect the mortality rate or expected life years. e_j is the random error which is assumed to be distributed normally with mean zero.

We use the regression discontinuity design (RDD) for our study (Imbens and Wooldridge,

2007). This methodology is based on the hypothesis that treated and untreated observations are similar near the threshold level of the treatment, thus, data can be analysed as if it were a (conditionally) randomized experiment. An important assumption is that the assignment mechanism is exogenous, and individuals are unable to precisely manipulate it (Lee and Lemieux, 2010). We identify the physical boundary of the Indo-Gangetic Plains from the geographical literature and use it as an exogenous threshold. For the validity of RDD a necessary identifying assumption is that there should not be a discontinuous jump in the unobserved determinants of the outcome variable. We check continuity around threshold for all the covariates as well as for predicted life expectancy.

The following equations are estimated to test whether the valley effect of the IGP causes a discontinuous change in the particulate exposure and health outcomes:

$$PM_j = \beta_0 + \beta_1 N_j + \beta_2 f(L_j) + X_j k + v_j \quad (2)$$

$$Y_j = \gamma_0 + \gamma_1 N_j + \gamma_2 f(L_j) + X_j \phi + u_j \quad (3)$$

where PM_j is the average annual exposure to $PM_{2.5}$ in district j over the period 1998-2014. Y_j is the health outcome in district j . N_j is a dummy variable which takes value one if a district lies in the IGP, and zero, otherwise. L_j denotes degrees north of the IGP boundary. Separate regressions using $f(L_j)$ as a linear function and as a quadratic function are conducted. X_j is a vector of other covariates that may affect health outcomes such as income, literacy rate, access to clean drinking water, access to clean cooking fuel, share of rural households and minority share in the population in a district. β_1 and γ_1 are the coefficients of interest,

measuring the local average treatment effect of being in the Indo-Gangetic Plains on exposure to fine particulate matter, and the health outcomes, respectively.

To find the average treatment effect of PM_{2.5} exposure on early life mortality and life expectancy at birth we use two-stage least squares method where PM_{2.5} exposure is instrumented by dummy variable indicating whether a district lies in IGP or not.

Equation (2) serves as the first stage equation, and the predicted values of PM_j are then used in the second stage, as shown in equation (4) below.

$$Y_j = \delta_0 + \delta_1 \widehat{PM}_j + \delta_2 f(L_j) + X_j \phi + e_j \quad (4)$$

where $\widehat{PM}_j$ denotes the fitted values obtained by estimating equation (2). The coefficient δ_1 estimates the average treatment effect of PM_{2.5} exposure on health outcomes. Instrumental variable estimation of equation (4) will result in a consistent estimator of δ_1 , if (i) there is strong partial correlation between the instrumental variable and the endogenous regressor in equation (2) ($\beta_1 \neq 0$), and (ii) there is no correlation between the instrumental variable and the error term in equation (4).

DATA SOURCES

Data on early life mortality rates viz., neonatal mortality rate (NNMR), infant mortality rate (IMR) and under-five mortality rate (U5MR) are computed from the National Family Health Survey (NFHS), Round 4, 2015-16. In the absence of data on life expectancy and age-specific death rates at district level, we use the United Nations method to compute life expectancy at birth (LEB). This method has been used by the Population Branch of the

United Nations, Department of Social Affairs to compute life expectancy using only infant mortality rate in countries lacking data on vital statistics (Kesarwani, 2015). The underlying assumption here is that each district follows the same fertility and mortality patterns as the state it belongs to. In this method, first, LEB is estimated as a function of IMR separately for each state using the following linear regression equation.

$$\ln(LEB_s) = a + b * IMR_s \quad (5)$$

where subscript s denotes variable at the state level. The values of a and b thus estimated are used to compute LEB at district level by using the formula below.

$$LEB_d = \exp(\hat{a} + \hat{b} * IMR_d) \quad (6)$$

where subscript d denotes variable at the district level, $\hat{a}$ and $\hat{b}$ denote the estimated values of a and b from equation (5). The methodology uses two data sources. The state-level data on life expectancy and infant mortality rate for a period from 1995-99 to 2012-16 are taken from the Sample Registration System (SRS) Abridged Life Table and the SRS Bulletin Reports, respectively. The data on infant mortality rate have been computed for 234 districts from NFHS (2015-16) using DHS guidelines.

District-level data on exposure to $PM_{2.5}$ was obtained from Van Donkelaar et al. (2016) for 234 districts in the sample from 1998 to 2014. They estimate $PM_{2.5}$ exposure by combining Aerosol Optical Depth (AOD) retrievals from the NASA MODIS, MISR, and SeaWiFS instruments with the GEOS-Chem chemical transport model, and then adjusting to the global ground-based observations of $PM_{2.5}$ using Geographically Weighted Regression (GWR). We

define “PM_{2.5} exposure” for a given year as the average of the PM_{2.5} readings in all the previous years. For instance, the PM_{2.5} exposure relevant to the year 2000 is calculated as the average of the PM_{2.5} concentrations in the years 1998, 1999 and 2000. Subsequently, PM_{2.5} exposure is averaged across all years for each district to arrive at the average PM_{2.5} exposure. This is used as a measure for the sustained level of PM_{2.5} exposure in each district. The data includes districts from the states of Bihar, Haryana, Punjab, Rajasthan, Madhya Pradesh, Uttar Pradesh and West Bengal and the union territory of Delhi. Jharkhand and Chhattisgarh have been excluded since they are newly formed states and the districts have been reorganized over time, which makes district level pollution data not comparable. The coordinates for the lower boundary of the IGP have been taken from the *Geomorphologic Atlas of Indo-Gangetic Plains* (Hecht and Sinha, 2003) created by the University of Technology, Dresden, Germany and the Indian Institute of Technology, Kanpur, India, with the help of ArcGIS software. The latitudinal difference between each district and the boundary is then estimated. Data on the demographic determinants of health are taken from Census 2011, NSSO-Round 68 and NFHS-4.

RESULTS

In this section, we discuss the validity of regression discontinuity design in our setup, the regression results based on RDD and 2SLS method, potential improvements in life expectancy on achieving the air quality standards (viz., national and WHO recommended), and robust checks done.

Validity of Regression Discontinuity Design

Table 1 presents the summary statistics of the key variables in our analysis. Columns (1) and (2) report the mean values of the variables in the districts lying north and south of the IGP boundary, respectively. Column (3) reports the difference in the mean values of the variables found in columns (1) and (2). Column (4) shows the difference in means of the variables once they are adjusted for a quadratic function of the difference in the degrees of latitude. This serves as a test for discontinuity at the IGP boundary. Rejection of the null hypothesis for a variable implies that there is a significant difference in the mean values of the variable on the two sides of the boundary.

PM_{2.5} exposure is observed to be significantly higher in the districts lying above the IGP boundary. This implies that the northern districts lying in the IGP are much more polluted (about 1.6 times) than the districts lying below the IGP boundary. On an average, the actual life expectancy at birth is lower by 1.77 years in the northern districts lying in the IGP relative to the southern districts in the sample. Checking if the observed determinants of health differ significantly across the boundary, we find that while the difference in income, literacy rate, and share of rural households is insignificant, the share of minority population is significantly lower, access to clean drinking water and access to clean cooking fuel is significantly higher in the treatment group. The direction of these latter differences is likely to lead to improvements in the health outcomes in the treatment group, thus, unlikely to weaken the basis of our analysis. We further test the validity of the RDD in the IGP setting by examining the predicted life expectancy in the two groups (Chen et al., 2013; Ebenstein et al. 2017). Predicted life expectancy is estimated as the fitted value from an OLS regression of life expectancy at birth on demographic and socioeconomic determinants of health except

PM_{2.5}. The difference in the predicted life expectancy is insignificant at the boundary as shown in column (4).

Graphically, we observe a striking discontinuous jump in the PM_{2.5} exposure of 22.5 µg/m³ of PM_{2.5} (around 47%) at the IGP boundary implying that the geography of the IGP has led to extremely high pollution levels in districts lying in the Indo-Gangetic Plain relative to those districts which are close to the boundary but lie on the other side (Figure 2). Similarly, a discontinuous fall in life expectancy of approximately 1.8 years is seen at the boundary (Figure 3).

Figure 4 presents the graphical analysis of the test for internal validity of the RDD. The fitted line is obtained by regressing predicted life expectancy at birth on a quadratic function in latitude. There is an insignificant difference in the predicted life expectancy at the IGP boundary. This implies that predicted life expectancy (excluding the impact of PM_{2.5}) moves smoothly across the boundary. The insignificant difference in predicted life expectancy provides support to the validity of our RDD, and shows that the demographic and socioeconomic variables are unable to explain the abrupt fall in life expectancy in districts lying just above the IGP boundary as shown in Figure 3.

Regression Results

The presentation of Tables 2 to 4 is explained as follows. Each cell in each of the tables reports results obtained from a separate regression. Each row presents results for a specific health outcome (namely, neonatal mortality rate, infant mortality rate, under-five mortality rate and life expectancy at birth). The columns report results from different model

specifications for that particular health outcome. Table 3 has an additional row presenting results pertaining to PM_{2.5}.

Table 2 presents the results obtained by using the conventional OLS approach. While Columns (1) and (2) report the regression results by estimating equation (1) excluding and including demographic and socioeconomic variables, respectively. Exposure to PM_{2.5} is found to adversely affect the health status of the population. From column (2), Additional 10 µg/m³ of PM_{2.5} exposure raises the number of deaths of new-borns within first 28 days of life by 1.7 per 1000 live births. Similarly, the number of deaths of infants and children under the age of 5 years rises by 3.1 and 3.7 per 1000 live births, respectively, due to additional 10 µg/m³ of PM_{2.5} exposure. Life expectancy at birth is found to be negatively affected by exposure to fine particulate matter. A rise in PM_{2.5} exposure by 10 µg/m³ reduces life expectancy by 0.54 years. The results are found to be significant at 1%.

Table 3 presents the local average treatment effect of residing in the IGP on PM_{2.5} levels and the above four health outcomes using RDD analysis . Each row presents results for a specific outcome, and different columns report results from different model specifications. Column (1) reports the results when the difference in latitude from the IGP boundary is used linearly. Columns (2) and (3) show the impact of “North” when the polynomial in latitude is quadratic. While column (2) excludes demographic and socio-economic controls, column (3) includes them.

PM_{2.5} exposure in districts in the IGP is significantly higher than the districts south of the boundary, ranging between ~15-23 µg/m³. Neonatal mortality rate is found to be higher on average by approximately 5.6 neonatal deaths per 1000 live births in the districts lying in the

IGP. The infant mortality rate in the IGP exceeds that in the districts lying below it in the sample by almost 6.5 deaths per 1000 live births. Similarly, the under-five mortality rate is also higher by around 7.4 deaths per 1000 live births in the IGP. This implies that neonatal mortality rate in districts lying above and close to the boundary is around 16.7% higher than the average NNMR in the districts south of the boundary. Similarly, IMR in districts north of the boundary is found to be around 13.6% higher than the districts south of the boundary. A 12.4% higher under-five mortality rate is found in northern districts than the average under-five mortality rate in the relevant southern districts. Life expectancy at birth is found to fall by 2.6 years when individuals reside in the IGP.

Table 4 presents results of the 2SLS regression analysis and the organization of the table is similar to Table 3. A rise in $PM_{2.5}$ exposure by $10 \mu g/m^3$ raises NNMR and IMR by 2.6 and 3, respectively. The neonatal period of death constituted around 87% of the infant deaths due to the rise in $PM_{2.5}$. The under-five mortality rate also rose by 3.6 with a $10 \mu g/m^3$ rise in $PM_{2.5}$ exposure. Life expectancy is found to fall by 1.2 years due to a $10 \mu g/m^3$ rise in $PM_{2.5}$, ceteris paribus. The 2SLS estimates are much higher than the OLS estimates.

Life years saved on attaining air quality standards

In our data, the range of $PM_{2.5}$ exposure is $33-108 \mu g/m^3$ and, the exposure within one standard deviation ranges between 45 and $82 \mu g/m^3$. This is similar to the range of $PM_{2.5}$ exposure in the entire country. Thus it is reasonable to generalize these results to the rest of India.

We now compute life years saved when a region is able to attain the national or WHO standards for $PM_{2.5}$ using the formula below:

$$\text{Life years saved} = 0.12 * (\text{Average } PM_{2.5} \text{ exposure} - PM_{2.5} \text{ standard})$$

where 0.12 is the estimated coefficient of $\widehat{PM}_j$ as reported in Table 4, column (3). In estimating the gain in life expectancy, a linear association between $PM_{2.5}$ exposure and life expectancy is assumed. The assumption bears support from the literature (GBD, 2016; Greenstone and Fan, 2018). Figure A1 and Table A1 in Appendix present gains in life expectancy at birth for different states and regions, respectively. The average $PM_{2.5}$ exposure in a region is estimated as its weighted average by using population as the weight. Assuming that the 2014 pollution concentration levels sustain, we find that India can raise life expectancy on average by 1.7 and 5.3 years on meeting the national and WHO standards for $PM_{2.5}$, respectively (*Appendix*, Table A1).

Robustness Checks

The findings are subjected to several robustness checks. First, a variable on interaction between the dummy variable, N_j and the quadratic function, $f(L_j)$ variable is added to the RDD and the 2SLS (*Appendix*, Table A2). Second, elevation above sea level and altitude of a district are included separately as proxies for topography. The estimates for pollution, early life mortality and life expectancy remain significant at 1% in each case (*Appendix*, Tables A3 and A4). Third, the sample is restricted to different bandwidths of latitudinal difference. The estimates for pollution, early life mortality and life expectancy remain significant even when the bandwidth reduces to 5°, 4° and 3° (*Appendix*, Tables A5 and

A6).

CONCLUSION

The study attempts to understand the impact of sustained exposure to PM_{2.5} exposure on four health outcomes (NNMR, IMR, U5MR and LEB) in India. The estimated impact of exposure to PM_{2.5} on life expectancy in this study is 71% higher than the estimated value by Pope et al. (2002) for the US (= -0.07). This strengthens the argument that utilizing the estimates from developed countries may not be desirable for developing countries due to several physiological, geographical, socioeconomic, weather-related and other differences between them. Comparing our results with Greenstone and Fan (2018), the estimates from this study are 22% higher as life expectancy reduces by 0.98 years for every additional 10 µg/m³ of PM_{2.5} above the PM_{2.5} standard in their study, whereas it reduces by 1.2 years in our study. The IMR effect of change in PM_{2.5} by 10 µg/m³ is 6.1% in our study, while it is 8.8% and 9.2% in the studies conducted in Mexico and Africa, respectively (Arceo et al., 2016; Heft-Neal et al., 2018).

Analysing individual cities of India, Delhi, Patna and Allahabad are likely to gain immensely in terms of life years (life expectancy gains exceeding 7 years) if the WHO standard for PM_{2.5} is attained in the districts (*Appendix*, Table A7). The PM_{2.5} exposure has been the highest in Delhi. Delhi has suffered a loss of 1.9 years of life due to increased PM_{2.5} exposure levels as compared to 1998 levels. Gains in life expectancy in Delhi are around 10.9 years when the WHO standards are reached and around 7.3 years when the national standards are met in the city. A significantly higher improvement in life expectancy can be seen in the states lying in the Indo-Gangetic Plains owing to higher levels of PM_{2.5} in

the region (*Appendix*, Fig. A1).

On 25th June 2015, India launched *100 smart cities mission* with a vision to promote cities, which will provide core infrastructure, and sustainable environment. Our results suggest that air pollution risk assessment should be integrated in the goals of smart cities, particularly in the vulnerable cities such as those in IGP that suffer from hazardous levels of air pollution. Measures to monitor and combat air pollution should be embedded in the city plan.

The future work can investigate the impact of exposure to air pollution (primarily PM_{2.5}, and PM₁) on morbidity and the overall mortality rate. An analysis of cause-specific mortality at the district level would be useful. With the presence of more comprehensive datasets, more elaborate studies can be done for India.

ACKNOWLEDGEMENT

The authors gratefully acknowledge Praffulit Bisht for his assistance in constructing the Indo-Gangetic Plains boundary.

REFERENCES

1. Aitchison, J.C., and A.M. Davis. 2007. "When and where did India and Asia collide?." *Journal of Geophysical Research*. 112: B05423.
2. Apte, J.S., and P. Pant, 2019. "Toward cleaner air for a billion Indians." *Proceedings of the National Academy of Sciences of the United States of America*. 116(22): 10614-16.
3. Burrard, S.J. 1915. "Origin of the Indo-Gangetic Trough, Commonly called the Himalayan Foredeep." *Proceedings of the Royal Society of London*. 91(628): 220-38.

4. Carre, J., N. Gatimel, J. Moreau, J. Parinaud, and R. Leandri. 2017. “Does air pollution play a role in infertility?: a systematic review.” *Environmental Health*. 16: 82.
5. Chen, Y., A. Ebenstein, M. Greenstone, and H. Li. 2013. “Evidence on the Impact of Sustained Exposure to Air Pollution on Life Expectancy from China’s Huai River Policy.” *Proceedings of the National Academy of Sciences of the United States of America*. 110(32): 12936-41.
6. Chowdhury, S., and S. Dey. 2016. “Cause-specific premature death from ambient PM2.5 exposure in India: estimate adjusted for baseline mortality.” *Environment International*. 91: 283–90.
7. Cropper, M.L., N.B. Simon, A. Alberini, and P.K. Sharma. 1997. “The Health Effects of Air Pollution in Delhi, India.” Policy Research Working Paper No. 1860, World Bank, Washington DC.
8. Ebenstein, A., M. Fan, M. Greenstone, G. He, and M. Zhou. 2017. “New evidence on the impact of sustained exposure to air pollution on life expectancy from China’s Huai River Policy.” *Proceedings of the National Academy of Sciences of the United States of America*. 114(39): 10384-89.
9. Global Burden of Disease Collaborative Network. 2017. *Global Burden of Disease Study 2016 Burden by Risk 1990-2016*. Seattle.
10. Greenstone, M., J. Nilekani, R. Pande, N. Ryan, A. Sudarshan, and A. Sugathan. 2015. “Lower Pollution, Longer Lives.” *Economic and Political Weekly*. L(8): 40-46.
11. Greenstone, M., and C.Q. Fan. 2018. “Introducing the Air Quality Life Index: Twelve Facts about Particulate Air Pollution, Human Health, and Global Policy.” Energy and Policy Institute, University of Chicago.

12. Guttikunda, S.K., and B.R. Gurjar. 2012. "Role of meteorology in seasonality of air pollution in megacity Delhi, India." *Environmental Monitoring and Assessment*. 184(5): 3199-3211.
13. Guttikunda, S.K., and P. Jawahar. 2012. "Application of SIM-air modeling tools to assess air quality in Indian cities." *Atmospheric Environment*. 62: 551–61.
14. Guttikunda, S.K., and R. Goel. 2013. "Health impacts of particulate pollution in a megacity- Delhi, India." *Environmental Development*. 6 8-20.
15. Hecht R., and R. Sinha. 2003. *Geomorphic Atlas of Indo-Gangetic Plains*. Available at <http://www.iitk.ac.in/gangetic/index.htm>. (2003)
16. Heft-Neal, S., J. Burney, E. Bendavid, and M. Burke. 2018. "Robust relationship between air quality and infant mortality in Africa." *Nature*. 559: 254-58.
17. Imbens, G., and J. Wooldridge, 2007. "Estimation of Average Treatment Effects Under Unconfoundedness." Lecture Notes, NBER, 30 July.
18. India State-Level Disease Burden Initiative Air Pollution Collaborators. 2019. "The impact of air pollution on deaths, disease burden, and life expectancy across the states of India: the Global Burden of Disease Study 2017." *Lancet Planetary Health*. 3: e26-e39.
19. Kandlikar, M., and G. Ramachandran. 2000. "The causes and consequences of particulate air pollution in urban India: a synthesis of the science." *Annual Review of Energy and the Environment*. 25: 629-84.
20. Kesarwani, R. 2015. "Estimation of Life Expectancy from Infant Mortality Rate at District Level." *International Research Journal of Social Sciences*. 4(6): 52-63.
21. Lavigne, E., R.T Burnett, D.M. Stieb, G.J. Evans, K.J.G. Pollitt, H. Chen, D. Rijswijk, and S. Weichenthal. 2018. "Fine Particulate Air Pollution and Adverse Birth

Outcomes: Effect Modification by Regional Nonvolatile Oxidative Potential.”

Environmental Health Perspectives. [CID:077012](#).

22. Lelieveld, J., A. Pozzer, U. Poschl, M. Fnais, A. Haines, and T. Munzel. 2020. “Loss of life expectancy from air pollution compared to other risk factors: a worldwide perspective.” *Cardiovascular Research*. <https://doi.org/10.1093/cvr/cvaa025>.
23. Lvovsky, K. 1998. “Economic Costs of Air Pollution with special reference to India.” Paper presented at National Conference on Health and Environment, Delhi, July 7-9.
24. Nema, P., and S.K. Goyal. 2010. “Estimation of health impacts due to PM10 in major Indian cities” In *Air Pollution: Health and Environmental Impacts*: Taylor & Francis Group, pp. 297-310.
25. Pope, C.A., T.B. Richard, and J.T. Michael. 2002. “Lung Cancer, Cardiopulmonary Mortality, and Long-Term Exposure to Fine Particulate Air Pollution.” *The Journal of the American Medical Association*. 287(9): 1132–41.
26. Shah J., and T. Nagpal. 1997a. “Urban AQMS in Asia, Greater Mumbai report.” World Bank Technical Paper No 381, World Bank, Washington, DC.
27. Tanaka, S. 2015. “Environmental Regulations on air pollution in China and their impact on infant mortality.” *Journal of Health Economics*. 42: 90-103.
28. Thistlethwaite, D.L., and D.T. Campbell. 1960. “Regression-discontinuity analysis: An alternative to the ex post facto experiment.” *Journal of Educational Psychology*. 51(6): 309-17.
29. Van Donkelaar, A., R.V. Martin, M. Brauer, N.C. Hsu, R.A. Kahn, R.C. Levy, A. Lyapustin, A.M. Sayer, and D.M. Winker. 2016. “Global Estimates of Fine Particulate Matter using a Combined Geophysical-Statistical Method with Information from Satellites, Models, and Monitors.” *Environmental Science & Technology* 50 (7): 3762-72.

30. World Health Organisation. 2016. "India takes steps to curb air pollution." *Bulletin of World Health Organisation*. 94: 487-88.
31. World Health Organisation. 2018. WHO Global Ambient Air Quality Database, Geneva.
32. Zhang, Y., J. Lang, S. Cheng, S. Li, Y. Zou, D. Chen, H. Zhang, and H. Wang. 2018. "Chemical composition and sources of PM1 and PM2.5 in Beijing in autumn." *Science of The Total Environment*. 630: 72-82.

TABLES

Table 1. Summary Statistics

Variable	North (1)	South (2)	Difference in means (3)	Adjusted difference in means (4)
I. Air pollution exposure				
<i>PM_{2.5}, $\mu\text{g}/\text{m}^3$</i>	74.52	47.92	26.60***	22.49***
II. Demographic characteristics				
<i>Literacy rate</i>	0.68	0.66	0.02	0.01
<i>Share, rural households</i>	0.77	0.79	-0.02	-0.03
<i>Share, treated tap water</i>	0.20	0.11	0.08***	0.10***
<i>Share, Clean Cooking Fuel</i>	0.33	0.25	0.08***	0.11***
<i>Share, Minority</i>	0.76	0.83	-0.07***	-0.06***
<i>Ln(Income)</i>	12.04	11.98	0.06	0.06
<i>Predicted Life Expectancy</i>	67.37	67.40	0.03	0.16
<i>Actual Life expectancy</i>	66.69	68.46	-1.77***	-1.82***

Note: n = 234. The results in column (4) are adjusted for a quadratic in degrees of latitude north of the Indo-Gangetic Plains boundary. Predicted life expectancy is calculated by OLS using all the demographic covariates shown.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table 2. Impact of PM_{2.5} on health outcomes using conventional strategy (OLS)

Dependent Variable	(1)	(2)
<i>Neo-Natal Mortality Rate</i>	0.09* (0.05)	0.17*** (0.04)
<i>Infant Mortality Rate</i>	0.20*** (0.06)	0.31*** (0.06)
<i>Under-Five Mortality Rate</i>	0.21*** (0.08)	0.37*** (0.07)
<i>Life Expectancy at Birth</i>	-0.04*** (0.01)	-0.05*** (0.01)
Number of observations	234	232
Demographic Controls	No	Yes

Note: Each cell in the table represents a coefficient from a separate regression, and heteroskedastic-consistent SEs are reported in parentheses.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table 3. Impact of being “North” of the boundary on listed variables, RDD

Main Independent Variable →	$N_j = 0, 1$		
Dependent Variables	(1)	(2)	(3)
$PM_{2.5}, \mu g/m^3$	15.12*** (2.61)	22.49*** (1.73)	23.33*** (2.01)
<i>Neo-Natal Mortality Rate</i>	3.51 (2.33)	1.80 (1.76)	5.64*** (1.69)
<i>Infant Mortality Rate</i>	1.47 (3.07)	1.18 (2.32)	6.45*** (2.31)
<i>Under-Five Mortality Rate</i>	1.33 (4.18)	-0.22 (3.24)	7.36** (3.14)
<i>Life Expectancy at Birth</i>	-1.26* (0.71)	-1.82*** (0.49)	-2.62*** (0.54)
Number of observations	232	234	232
Demographic Controls	Yes	No	Yes
Polynomial in latitude	Linear	Quadratic	Quadratic

Note: Each cell in the table represents a coefficient from a separate regression, and

heteroskedastic-consistent SEs are reported in parentheses. Models in column (1) are

estimated with a linear control for latitudinal difference and include demographic controls.

Models in column (2) include a quadratic in latitudinal difference and do not include demographic and socioeconomic controls. Models in column (3) include demographic and socioeconomic controls along with quadratic latitudinal difference.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table 4. Impact of PM_{2.5} exposure on health outcomes, 2SLS

Main Independent Variable →	<i>Fitted values of PM_{2.5} (μg/m³)</i>		
Dependent Variable	(1)	(2)	(3)
<i>Neo-natal Mortality Rate</i>	0.27* (0.16)	0.08 (0.08)	0.26*** (0.07)
<i>Infant Mortality Rate</i>	0.14 (0.20)	0.05 (0.10)	0.30*** (0.09)
<i>Under-Five Mortality Rate</i>	0.15 (0.27)	-0.01 (0.14)	0.36*** (0.13)
<i>Life Expectancy at Birth</i>	-0.09* (0.05)	-0.08*** (0.02)	-0.12*** (0.02)
Number of observations	232	234	232
Demographic Controls	Yes	No	Yes
Polynomial in latitude	Linear	Quadratic	Quadratic

Note: Each cell in the table represents a coefficient from a separate regression, and

heteroskedastic-consistent SEs are reported in parentheses. Models in column (1) are estimated with a linear control for latitude. Models in column (2) include a quadratic in latitude. Models in column (3) additionally include demographic controls. Two observations

are excluded in the second and fourth columns because of missing data on income in Palwal (Haryana) and Pratapgarh (Rajasthan).

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

FIGURES

Fig. 1. Lower boundary of the Indo-Gangetic Plains depicted in brown, used as the threshold for conducting Regression Discontinuity Analysis

Source: Plotted using ArcGIS

Fig. 2. Fitted values of PM_{2.5} exposure across the IGP boundary

Fig. 3. Fitted values of Life Expectancy at Birth across the IGP boundary

Fig. 4. Fitted values of Predicted Life Expectancy across the IGP Boundary

Appendix For:

**Health Effects of Sustained Exposure to Fine Particulate Matter:
Evidence from India**

Yashaswini Saraswat ¹ & Sangeeta Bansal ¹

This PDF file includes :

Definitions

Figure A1

Tables A1 to A7

¹ Centre for International Trade and Development, Jawaharlal Nehru University, Delhi-67

DEFINITIONS

1. Neonatal Mortality Rate (NNMR): Probability of dying during the first 28 days of life, expressed per 1,000 live births.

Source: Computed as per UNICEF definition from NFHS 4, 2015-16

2. Infant Mortality Rate (IMR): Probability of dying between birth and exactly 1 year of age, expressed per 1,000 live births.

Source Computed as per UNICEF definition from NFHS 4, 2015-16

3. Under-Five Mortality Rate (U5MR): Probability of dying between birth and exactly 5 years of age, expressed per 1,000 live births.

Source: Computed as per UNICEF definition from NFHS 4, 2015-16

4. Life Expectancy at Birth (LEB): Average number of years that a new-born is expected to live if current mortality rates continue to apply.

Source: Computed as per WHO definition from NFHS 4, 2015-16

5. Latitudinal difference: Latitudinal difference between a district and a corresponding district lying in the lower boundary of the Indo-Gangetic Plains.

6. Literacy Rate: Proportion of people aged 7 and above who can both read and write with understanding in any language.

Source: Census 2011

7. Access to treated tap water: Proportion of households who have access to treated tap water within their premises.

Source: Census 2011

8. Access to clean cooking fuel: Proportion of households who use LPG or electricity as the primary source of fuel for cooking purposes.

Source: NFHS 4, 2015-16

9. Share of rural households: Proportion of households not in urban areas. An urban area is defined as:

(a) all places with a Municipality, Corporation or Cantonment or Notified Town Area

(b) all other places which satisfied the following criteria:

(i) a minimum population of 5,000.

(ii) at least 75% of the male working population was non-agricultural.

(iii) a density of population of at least 400 sq. Km. (i.e. 1000 per sq. Mile)

Source: Census 2011

10. Consumption expenditure: Household Consumer Expenditure (HCE) is most easily understood as expenditure incurred by households on consumption goods and services, i.e., on goods and services used for the direct satisfaction of individual needs and wants or the collective needs of members of the community and not for further transformation in production.

Source: NSSO, Round 68, 2011-12

11. Share of minority population: Proportion of households belonging to Scheduled Caste, Scheduled Tribe or Other Backward Class.

Source: NFHS 4, 2015-16

Fig. A1. State-wise gains in life expectancy on meeting WHO standard for PM_{2.5}

Table A1. Life years saved on attaining the air quality standards

Region	Weighted Average PM _{2.5} (µg/m ³)	Life Years Saved: National Standard	Life Years Saved: WHO Standard
National	54.07	1.69	5.29
Indo-Gangetic Plains	83.70	5.24	8.84

Table A2. Inclusion of interaction variable for robustness check

Main Independent Variable →	$N_j = 0, 1$	Fitted values of $PM_{2.5}$ ($\mu g/m^3$)
Dependent Variable	(1)	(2)
$PM_{2.5}$, $\mu g/m^3$	22.76*** (2.26)	-
<i>Neo-natal Mortality Rate</i>	4.08** (1.84)	0.20** (0.08)
<i>Infant Mortality Rate</i>	3.77 (2.48)	0.19* (0.10)
<i>Under Five Mortality Rate</i>	3.98 (3.35)	0.21 (0.14)
<i>Life Expectancy at Birth</i>	-1.69*** (0.56)	-0.08*** (0.02)
Number of observations	232	232
Demographic Controls	Yes	Yes
Polynomial in latitude	Quadratic	Quadratic
Estimation method	RDD	2SLS

Note: Each cell in the table represents a coefficient from a separate regression, and heteroskedastic-consistent SEs are reported in parentheses. The estimates in column (1) denote the impact of being located in the IGP on the relevant dependent variables. The estimates in column (2) denote the change in the dependent variable due to a unit change in the $PM_{2.5}$ concentration level, ceteris paribus.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table A3. Inclusion of elevation for robustness check

Main Independent Variable →	$N_j = 0, 1$	Fitted values of $PM_{2.5}$ ($\mu g/m^3$)
Dependent Variable	(1)	(2)
$PM_{2.5}$, $\mu g/m^3$	20.90*** (2.40)	-
<i>Neo-natal Mortality Rate</i>	5.99*** (1.96)	0.31*** (0.10)
<i>Infant Mortality Rate</i>	8.78*** (2.74)	0.45*** (0.12)
<i>Under Five Mortality Rate</i>	9.70*** (3.70)	0.51*** (0.17)
<i>Life Expectancy at Birth</i>	-3.04*** (0.64)	-0.15*** (0.03)
Number of observations	228	228
Demographic Controls	Yes	Yes
Polynomial in latitude	Quadratic	Quadratic
Estimation method	RDD	2SLS

Note: Each cell in the table represents a coefficient from a separate regression, and heteroskedastic-consistent SEs are reported in parentheses. The estimates in column (1) denote the impact of being located in the IGP on the relevant dependent variables. The estimates in column (2) denote the change in the dependent variable due to a unit change in the $PM_{2.5}$ concentration level, ceteris paribus.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table A4. Inclusion of altitude for robustness check

Main Independent Variable →	$N_j = 0, 1$	Fitted values of $PM_{2.5}$ ($\mu g/m^3$)
Dependent Variable	(1)	(2)
$PM_{2.5}$, $\mu g/m^3$	19.46*** (2.56)	-
<i>Neo-natal Mortality Rate</i>	7.05*** (2.21)	0.39*** (0.12)
<i>Infant Mortality Rate</i>	10.02*** (3.12)	0.55*** (0.15)
<i>Under Five Mortality Rate</i>	11.46*** (4.25)	0.64*** (0.20)
<i>Life Expectancy at Birth</i>	-3.04*** (0.70)	-0.16*** (0.04)
Number of observations	232	232
Demographic Controls	Yes	Yes
Polynomial in latitude	Quadratic	Quadratic
Estimation method	RDD	2SLS

Note: Each cell in the table represents a coefficient from a separate regression, and heteroskedastic-consistent SEs are reported in parentheses. The estimates in column (1) denote the impact of being located in the IGP on the relevant dependent variables. The estimates in column (2) denote the change in the dependent variable due to a unit change in the $PM_{2.5}$ concentration level, ceteris paribus.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table A5. Impact of “North” on listed variables, RDD for different bandwidths

Main Independent Variable →	$N_j = 0, 1$		
Dependent Variable	(1)	(2)	(3)
<i>PM_{2.5}, $\mu\text{g}/\text{m}^3$</i>	22.52*** (2.03)	22.39*** (2.03)	22.38*** (2.11)
<i>Neo-natal Mortality Rate</i>	5.83*** (1.72)	5.92*** (1.73)	6.07*** (1.75)
<i>Infant Mortality Rate</i>	6.85*** (2.36)	6.84*** (2.37)	7.05*** (2.42)
<i>Under Five Mortality Rate</i>	7.58** (3.23)	7.68** (3.25)	8.16** (3.28)
<i>Life Expectancy at Birth</i>	-2.85*** (0.55)	-2.76*** (0.54)	-2.43*** (0.54)
Number of observations	218	211	192
Demographic Controls	Yes	Yes	Yes
Polynomial in latitude	Quadratic	Quadratic	Quadratic
Bandwidth	5°	4°	3°

Note: Each cell in the table represents a coefficient from a separate regression, and heteroskedastic-consistent SEs are reported in parentheses.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table A6. Impact of PM_{2.5} on listed variables, 2SLS for different bandwidths

Main Independent Variable →	<i>Fitted values of PM_{2.5} (μg/m³)</i>		
Dependent Variable	(1)	(2)	(3)
<i>Neo-natal Mortality Rate</i>	0.36*** (0.11)	0.36*** (0.12)	0.31** (0.12)
<i>Infant Mortality Rate</i>	0.52*** (0.15)	0.52*** (0.15)	0.42*** (0.16)
<i>Under Five Mortality Rate</i>	0.61*** (0.20)	0.61*** (0.21)	0.51** (0.22)
<i>Life Expectancy at Birth</i>	-0.15*** (0.04)	-0.14*** (0.04)	-0.13*** (0.04)
Number of observations	218	211	192
Demographic Controls	Yes	Yes	Yes
Polynomial in latitude	Quadratic	Quadratic	Quadratic
Bandwidth	5°	4°	3°

Note: Each cell in the table represents a coefficient from a separate regression, and heteroskedastic-consistent SEs are reported in parentheses.

*** Significant at 1% level, ** Significant at 5% level, * Significant at 10% level.

Table A7. 20 Most Populous Districts

District	Population (Millions)	PM _{2.5} Concentration, 2014 (µg/m ³)	PM _{2.5} Concentration, 1998 (µg/m ³)	Increase in Life Expectancy if district meets National Standard (40 µg/m ³)	Increase in Life Expectancy if district meets WHO Standard (10 µg/m ³)	Change in Life Expectancy Due to Change in PM _{2.5} , 1998-2014 (years)
Delhi	17	101.2	85.5	7.3	10.9	-1.9
Thane, Maharashtra	11	41.5	32.9	0.2	3.8	-1
North 24 Parganas, West Bengal	10	43.5	37.1	0.4	4	-0.8
Bangalore Urban, Karnataka	9.6	29.7	27.2	0	2.4	-0.3
Mumbai (Suburban), Maharashtra	9.4	45.2	36.3	0.6	4.2	-1.1
Pune, Maharashtra	9.4	44.8	33.3	0.6	4.2	-1.4

South 24 Parganas, West Bengal	8.2	41.8	37.8	0.2	3.8	-0.5
Bardhaman, West Bengal	7.7	50.7	49.6	1.3	4.9	-0.1
Ahmadabad, Gujarat	7.2	44.2	47.1	0.5	4.1	+0.3
Murshidabad, West Bengal	7.1	57.9	48.9	2.1	5.7	-1.1
Jaipur, Rajasthan	6.6	53.8	50	1.7	5.3	-0.5
Nashik, Maharashtra	6.1	36.8	27.9	0	3.2	-1.1
Surat, Gujarat	6.1	39.9	34	0	3.6	-0.7
Allahabad, Uttar Pradesh	6	72.8	65.4	3.9	7.5	-0.9
Paschim Medinipur, West Bengal	5.9	46.2	47.4	0.7	4.3	+0.1
Patna, Bihar	5.8	84.9	63.2	5.4	9	-2.6
Hugli, West Bengal	5.5	47.3	46.6	0.9	4.5	-0.1

Rangareddy, Telangana	5.3	34.6	29.8	0	3	-0.6
East Godavari, Andhra Pradesh	5.2	36.1	30.1	0	3.1	-0.7
Nadia, West Bengal	5.2	53.2	45.4	1.6	5.2	-0.9