



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search  
<http://ageconsearch.umn.edu>  
[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

## **YIELD RESPONSE OF SELECTED IRRIGATED RICE VARIETIES TO VARYING N, P AND K FERTILIZER REGIMES**

**J. Chirchir<sup>1</sup>, G. Chemining'wa<sup>2</sup>, F. Nzuve<sup>3</sup> and R. Murori<sup>4</sup>**

*Received 24 September 2018, Revised 3 December 2018, Accepted 26 December 2018, Published online 31 December 2018*

### **Abstract**

Rice production has been inconsistent in Mwea irrigation scheme due to poor crop management practices and reduced soil fertility. Proper rice nutrition conserves the environment, increases sustained crop production, farmer's crop yield and profits. A field experiment was conducted at MIAD Center, in Kirinyaga County, during 2016-17 and 2017-18 to determine the response of rice varieties to different N, P and K fertilizer treatments. The trial was conducted in randomized complete block design, with three replications of 13 N, P and K fertilizer regimes as the main plots treatments and varieties Basmati 370 and BW 196 as the sub-plot treatment. The N kg ha<sup>-1</sup>: P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>: K<sub>2</sub>O kg ha<sup>-1</sup> fertilizer treatment ratios used were: 00:00:00, 60:40:40, 80:60:60, 100:80:80, 60:40:00, 80:60:00, 100:80:00, 60:00:40, 80:00:60, 100:00:80, 00:40:40, 00:60:60 and 00:80:80. Plant height, number of tillers hill<sup>-1</sup>, panicle length, and grain yield responded positively to fertilizer application, but 1000-grain weight did not. Variety BW 196 recording shorter plants and panicles, more tillers hill<sup>-1</sup>, higher grain weight and yield than variety Basmati 370. 00:40:40, 00:60:60, 00:80:80, 60:40:00 and the no-fertilizer control treatments recorded the least number of tillers hill<sup>-1</sup>. 60:40:40, 80:60:60, 100:80:80, fertilizer treatments had longer panicles than the no fertilizer control and 00:40:40. Except for 100:80:00 and 00:40:40, in the first season and 60:40:00, 80:60:00, 100:80:00 in the second season, all the fertilizer regimes increased grain yield relative to the control. The highest grain yield was realized in fertilizer regime 80:60:60, 100:80:80, 80:00:60 and 100:00:80. 80:00:60 is the recommended fertilizer regime.

**Keywords:** Crop Productivity, Fertilizer Regimes, Varieties, Mwea

<sup>1,2&3</sup>Department of Plant Science and Crop Protection, Faculty of Agriculture, University of Nairobi, P.O. Box 29053, Nairobi, Kenya

<sup>4</sup>International Rice Research Institute (IRRI), Eastern and South Africa (ESA) Hub, P.O. Box 30709-00100, IRRI c/o ILRI, Nairobi, Kenya

\*Corresponding author's email: [jedida.chirchir@gmail.com](mailto:jedida.chirchir@gmail.com) (J. Chirchir)

Cite this article as: Chirchir, J., Chemining' wa, G., Nzuve, F. and Murori, R. (2018). Yield response of selected irrigated rice varieties to varying N, P and K fertilizer regimes. *Int. J. Agril. Res. Innov. Tech.* 8(2), 42-46.

### **Introduction**

Traditionally, rice crop was prominent only in Asia, but over the years, it has also gained prominence in Africa's farming systems and diet. In Africa, rice is ranked fourth after maize, sorghum and millet, in terms of acreage grown. Its production is ranked second after maize (FAOSTAT, 2012). Several studies have shown that Kenya has the capacity of producing over 9 t ha<sup>-1</sup> of rice (Saito *et al.*, 2013). In spite of that, the mean rice yields in Kenya between the year 2005 and 2009, was 2.5 t ha<sup>-1</sup> (Onyango, 2014). This translates to a yield gap of over 6.5 t ha<sup>-1</sup>.

According to reports from MOA (2009), the rice production sector in Kenya faces several setbacks, such as: poor infrastructure, high costs of farm inputs and machinery, uncoordinated marketing, inadequate rice crop management skills, high

disease incidences, land degradation and nutrient losses. In Mwea irrigation scheme, the average crop production has been fluctuating due to reduced soil fertility, soil degradation and use of inappropriate crop management practices by farmers (Nyamai *et al.*, 2012). Formerly, farmers in Mwea used to burn one part of their farm and spread ashes across the paddy fields. Because of the increasing demand for animal feed, straw is baled and sold; hence, organic matter is transported to a different location. Husks are also disposed elsewhere after dehulling (Muhunyu, 2012). The mass transportation of organic matter from the rice fields, combined with permanent water logging and rice monoculture reduces the inherent capacities of the soil to supply nutrients, and hence results in long-term detrimental effects on crop production.

Balanced application of inorganic fertilizers is important for increased yields and sustained production of quality rice. It aids in reducing injury from infestation of pests and diseases, and minimizes expenditures that are to be used in controlling them. Nutrient requirements vary with varying varieties of rice (Dobermann and Fairhurst, 2000). In Mwea, at least all farmers use inorganic fertilizers, with 86% using DAP (Di-ammonium Phosphate) as basal fertilizer, 93% using SA (Sulphate of Ammonia) as a top dresser. Only 2% use MOP (Muriate of Potash) as a basal fertilizer (Muhunyu, 2012). Irrespective of the soil nutrient status and rice variety planted, most farmers apply 50 kg acre<sup>-1</sup> of either DAP or SA, to their fields. Such application of blanket fertilizer could have significant negative effect on the performance and yield of rice. Good nutrition could lower the current rice yield gaps (6.5 t ha<sup>-1</sup>) in Kenya. Considering these perspectives, the current study was set up to monitor the response of selected rice varieties to varying nitrogen, phosphorus and potassium fertilizer regimes.

## Materials and Methods

The present experiment was conducted for two seasons; between the years 2016 and 2018. It was setup in MIAD center, in Kirinyaga County. MIAD is located to the northeastern region of Nairobi, Kenya's capital, and to the southeastern part of Mount Kenya (Ngige, 2004). Before setting up the trial plots, soil at the experimental site was collected from different homogenous units, at two depths (0-15 and 15-30 cm), and bulked into two samples. The soils were analyzed for macro and micronutrients, pH, electrical conductivity and organic carbon. Very low levels of potassium (68.00, 37.10 ppm) and phosphorus (3.06, 0.93 ppm), and considerably low levels of Boron (0.21, 0.33 ppm) and zinc (1.02, 1.12 ppm) were reported at both depths.

Meteorological data was collected from MIAD meteorological station, during the research period. The highest rainfall (481 mm) was recorded in November 2017 and the lowest (0.00 mm) in January 2017. The mean temperature ranged from 21.75 to 28.79°C. The highest relative humidity was 86.47% (November 2018) and the lowest was 69.74% (January 2017).

The experiment was conducted in randomized complete block design, with three replications of 13 nitrogen, phosphorus and potassium fertilizer regimes as the main plots treatments and varieties Basmati 370 and BW 196 as the sub-plot treatment. The N kg ha<sup>-1</sup>: P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>: K<sub>2</sub>O kg ha<sup>-1</sup> fertilizer treatment ratios used were: 00:00:00, 60:40:40, 80:60:60, 100:80:80, 60:40:00, 80:60:00, 100:80:00, 60:00:40, 80:00:60, 100:00:80, 00:40:40, 00:60:60 and 00:80:80. The sources of nitrogen, phosphorus and potassium fertilizer were Urea, Triple superphosphate (TSP) and Muriate of potash (MOP), respectively.

Each 9 m<sup>2</sup> nutrient plot was bunded and lined with a 500 gauge polythene bag (45 cm deep), to minimize mixing of nutrients and to maintain a uniform water depth. Plants in each plot had a 20 x 20 cm spacing.

To monitor the response of the crop to varying inorganic fertilizer regimes, data was collected from a sample of 10 plants (IRRI, 2014). Observations on plant height and the number of tillers hill<sup>-1</sup> were collected at 30 and 75 DAT. Plant height measurements were taken from the soil surface to the apex of the tallest panicle, excluding awns. The tiller count hill<sup>-1</sup> was counted on the ten marked hills plot<sup>-1</sup>. At maturity, paddy rice was harvested, and dried to 14% moisture content. Grain yield was weighed and the weight of 1000 well developed seeds were also recorded. Using the 15<sup>th</sup> edition of Genstat software, the collected data was subjected to analysis of variance. The Least Significant Difference (LSD) test (at P≤0.05) was used to compare means between treatments.

## Results and Discussion

A positive response of varieties Basmati 370 and BW 196 to nitrogen, phosphorus and potassium fertilizer, in plant height, number of tillers hill<sup>-1</sup>, panicle length and grain yield was recorded in this study. This could be due to the low levels of these plant nutrients in the soils in the study area. In the present study applying 80:60:60, 100:80:80 and 100:00:80 fertilizer treatments augmented plant height relative to the no-fertilizer control. N omission treatments (00:40:40, 00:60:60, 00:80:80) did not have a significant effect on plant height relative to the no fertilizer control. The increased plant height in fertilizer regime 80:60:60 and 100:80:80 could be explained by the sufficient supply of rice nutrients. The inadequate supply of N in 00:40:40, 00:60:60, 00:80:80 resulted in plant heights resembling those of the no-fertilizer control (Table 1); because nitrogen deficiency must have disrupted metabolic processes of the plants (Dobermann and Fairhurst, 2000).

Fertilizer regime: 00:40:40, 00:60:60, 00:80:80, 60:40:00 and 00:00:00 registered the least number of tillers hill<sup>-1</sup>. The poor performance of the tillers could be due to inadequate nitrogen and potassium fertilizer in the soils of study. The good performance in N, P, K (60:40:40, 80:60:60, 100:80:80), N, P (60:40:00, 80:60:00, 100:80:00) and N, K (100:00:80 and 80:00:60) fertilizer regime (Table 2) could be explained by the increased uptake of nitrogen, phosphorus and potassium from the soils of study. Nitrogen stimulates plant growth (Doberman and Fairhurst, 2000). Potassium increases number of tillers hill<sup>-1</sup> by improving the crops metabolic processes. Availability of phosphorus encourages development of roots and production of large healthy leaves (Plenet et al., 2000).

Table 1. Effect of varying fertilizer regimes on the average plant height (cm) of two rice varieties at 30 and 75 days after transplanting in first and second season at MIAD.

Fertilizers	First season						Second season					
	Height 30-DAT			Height 75-DAT			Height 30-DAT			Height 75-DAT		
N: P <sub>2</sub> O <sub>5</sub> : K <sub>2</sub> O	Bs	BW	Mean									
00:00:00	19	15	17	86	40	63	39	37	38	103	46	75
60:40:40	24	19	21	102	47	74	42	41	42	109	61	85
80:60:60	23	20	22	100	53	76	50	42	46	117	58	88
100:80:80	24	20	22	102	54	78	45	43	44	114	59	87
60:40:00	21	18	20	95	47	71	42	39	41	98	49	73
80:60:00	22	19	20	93	47	70	46	38	42	107	51	79
100:80:00	25	20	22	102	49	75	40	40	40	102	57	79
60:00:40	21	19	20	94	49	72	44	40	42	110	53	81
80:00:60	21	19	20	94	50	72	48	42	45	115	60	87
100:00:80	23	20	22	105	53	79	41	41	41	113	64	89
00:40:40	19	17	18	93	43	68	41	43	42	110	54	82
00:60:60	20	16	18	94	44	69	45	39	42	109	51	80
00:80:80	20	17	19	89	44	66	45	39	42	113	54	83
Mean	22	18	20	96	48	72	44	41	42	109	55	82
Fpr V	<.001			<.001			<.001			<.001		
Fpr F	<.001			<.001			0.02			<.001		
Fpr F*V	1.0			0.9			0.2			0.2		
LSD <sub>0.05</sub> V	0.7*			2.6*			1.5*			2.1*		
LSD <sub>0.05</sub> F	1.9*			6.5*			3.9*			5.5*		
LSD <sub>0.05</sub> F*V	NS			NS			NS			NS		
CV %	8.0			7.8			7.9			5.7		

Fpr- F probability, LSD- Least significant difference at  $P_{0.05}$ , V-variety, F-fertilizer regimes, F\*V- interaction between Variety and Fertilizer regimes, NS- Not significant at  $P_{0.05}$ , \*-significant at  $P_{0.05}$ , CV- co-efficient of variation Ratio of N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O refers to N kg ha<sup>-1</sup>: P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>: K<sub>2</sub>O kg ha<sup>-1</sup>

Table 2. Effects of varying fertilizer regimes on the mean of tiller numbers hill<sup>-1</sup> of two rice varieties at 30 and 75 days after transplanting in the first and the second season at MIAD.

Treatments	First season						Second season					
	Tillers 30			Tillers 75			Tillers 30			Tillers 75		
N: P <sub>2</sub> O <sub>5</sub> : K <sub>2</sub> O	Bs	BW	Mean	Bs	BW	Mean	Bs	BW	Mean	Bs	BW	Mean
00:00:00	3.6	4.8	4.2	12	24	18	4.8	6.9	5.8	14	23	18
60:40:40	4.6	6.1	5.4	17	35	26	5.5	6.7	6.1	19	29	24
80:60:60	5.0	6.9	6.0	22	27	25	8.9	9.7	9.3	18	28	23
100:80:80	5.7	7.7	6.7	18	36	27	7.3	7.4	7.4	19	29	24
60:40:00	4.2	6.7	5.5	16	28	22	5.4	7.1	6.3	19	23	21
80:60:00	4.9	8.7	6.8	17	39	28	6.5	7.2	6.9	17	26	21
100:80:00	5.7	7.6	6.7	20	33	26	4.4	6.9	5.7	17	29	23
60:00:40	3.7	5.7	4.7	17	30	23	6.2	8.1	7.2	16	26	21
80:00:60	4.1	6.7	5.4	17	35	26	7.1	7.5	7.3	20	27	23
100:00:80	4.7	5.9	5.3	18	32	25	4.3	5.7	5.0	19	30	24
00:40:40	3.7	5.2	4.4	14	27	20	5.2	7.9	6.6	13	23	18
00:60:60	4.8	6.6	5.7	15	24	19	6.4	6.7	6.6	14	25	20
00:80:80	5.5	6.8	6.2	13	24	19	5.9	6.2	6.1	16	24	20
Mean	4.6	6.6	5.6	17	30	23	6.0	7.2	6.6	17	26	22
Fpr V	<.001			<.001			<.001			<.001		
Fpr F	<.001			<.001			<.001			0		
Fpr F*V	0.7			0.1			0.8			0.6		
LSD <sub>0.05</sub> V	0.5*			1.9*			0.6*			1.3*		
LSD <sub>0.05</sub> F	1.1*			4.9*			1.6*			3.3*		
LSD <sub>0.05</sub> F*V	NS			NS			NS			NS		
CV %	17.5			17.9			20.2			13.4		

Fpr- F probability, LSD- Least significant difference at  $P_{0.05}$ , V-variety, F-fertilizer regimes, F\*V- interaction between Variety and Fertilizer regimes, NS- Not significant at  $P_{0.05}$ , \*-significant at  $P_{0.05}$ , CV- co-efficient of variation Ratio of N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O refers to N kg ha<sup>-1</sup>: P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>: K<sub>2</sub>O kg ha<sup>-1</sup>

60:40:40, 80:60:60, 100:80:80, N, P and K fertilizer regimes had significantly higher rice panicle lengths than the no fertilizer control and 00:40:40, in both seasons. Application of these fertilizer treatments may have improved the nutrient use efficiency of the rice crop; therefore increasing its chlorophyll content and enhancing proper assimilate translocation to reproductive parts, hence augmenting rice crop yield (Sarkar and Malik, 2001). Poor panicle development in 80:60:00, 100:80:00 could be have been due to the low (68 ppm) potassium levels in the study soils. Plots treated with fertilizer treatments:

00:00:00 and 00:40:40 could have been nutrient deficient resulting in poor development of panicles. Application of excess potassium and phosphorus where nitrogen was deficient (00:80:80) could have inhibited panicle development (Dobermann and Fairhurst, 2000). From the results of this study, fertilizer application had an insignificant effect on 1000-grain weight of rice (Table 3). A thousand grain weight is not prone to effects by environmental factors; it is controlled genetically (Yoshida, 1981). This explains why BW 196 had higher 1000-grain weight than Bs 370.

Table 3. Effect of different fertilizer regimes on yield parameters of two rice varieties planted in the first and the second season, between August 2016 and February 2018 at MIAD.

Treatments	First season						Second season					
	Panicle length (cm)			Grain weight (g)			Panicle length (cm)			Grain weight (g)		
N: P <sub>2</sub> O <sub>5</sub> : K <sub>2</sub> O	Bs 370	BW 196	Mean	Bs 370	BW 196	Mean	Bs 370	BW 196	Mean	Bs 370	BW 196	Mean
00:00:00	21.70	18.96	20.33	20.70	28.70	24.70	24.16	20.64	22.40	21.20	26.40	23.80
60:40:40	23.30	19.56	21.43	20.40	28.50	24.40	24.44	22.12	23.28	21.60	26.60	24.10
80:60:60	22.43	21.70	22.07	21.30	28.60	24.90	25.36	21.90	23.63	19.80	26.80	23.30
100:80:80	23.50	20.26	21.88	20.50	29.10	24.80	24.68	21.83	23.26	19.30	26.70	23.00
60:40:00	23.04	19.50	21.27	19.50	27.60	23.60	24.32	21.45	22.89	19.50	27.00	23.30
80:60:00	22.53	19.26	20.90	21.00	26.40	23.70	24.25	19.61	21.93	20.10	27.30	23.70
100:80:00	22.93	18.70	20.82	19.30	27.70	23.50	24.63	20.75	22.69	19.90	26.10	23.00
60:00:40	22.96	20.40	21.68	20.90	28.10	24.50	24.64	21.35	22.99	20.70	26.50	23.60
80:00:60	22.63	20.40	21.52	20.50	29.70	25.10	25.56	21.97	23.77	20.20	27.00	23.60
100:00:80	23.43	20.16	21.80	23.20	26.90	25.10	26.16	22.26	24.21	21.60	28.50	25.10
00:40:40	22.10	19.20	20.65	19.80	30.20	25.00	24.33	21.23	22.78	21.60	27.10	24.40
00:60:60	22.39	20.30	21.35	19.70	28.90	24.30	25.11	21.88	23.50	20.80	27.70	24.30
00:80:80	22.26	19.73	21.00	21.00	30.10	25.60	24.41	21.22	22.81	21.70	28.10	24.90
Mean	22.71	19.86	21.28	20.60	28.50	24.50	24.77	21.40	23.09	20.60	27.10	23.80
Fpr V		<.001			<.001			<.001			<.001	
Fpr F		0.03			0.5			0.03			0.5	
Fpr F*V		0.16			0.1			0.92			1.0	
LSD <sub>0.05</sub> V		0.40*			0.75*			0.47*			0.76*	
LSD <sub>0.05</sub> F		1.03*			NS			1.19*			NS	
LSD <sub>0.05</sub> F*V		NS			NS			NS			NS	
CV %		4.11			6.70			4.40			5.60	

Fpr- F probability, LSD- Least significant difference at  $P_{0.05}$ , V-variety, F-fertilizer regimes, F\*V- interaction between Variety and Fertilizer regimes, NS- Not significant at  $P_{0.05}$ , \* -significant at  $P_{0.05}$ , CV- co-efficient of variation Ratio of N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O refers to N kg ha<sup>-1</sup>: P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>: K<sub>2</sub>O kg ha<sup>-1</sup>

In the current study, higher (4.86 t ha<sup>-1</sup>) mean grain yield was registered in BW 196 than Basmati 370 (4.04 t ha<sup>-1</sup>). The yield registered for variety Bs 370 in the current study is in the range (4.1 to 6 t ha<sup>-1</sup>) reported by Ndiiri et al. (2013), from farmers' fields in Mwea. All the fertilizer treatments augmented grain yield relative to the control with an exception of 100:80:00 and 00:40:40, in the first season and 60:40:00, 80:60:00, 100:80:80 in the second season (Table 4). The highest grain yield was reported in fertilizer regime: 80:60:60, 80:00:60, 100:80:80 and 100:00:80 in both seasons. 80:00:60 is the

best fertilizer regime for rice production in Mwea, because it costs less than the rest that exhibit equal performance. For the advantage of sustainable land use, 80:60:60 and 100:80:80 fertilizer treatment would be suitable. To develop more precise soil recommendations, future research should seek to shed light on how long fertilizer regimes, 80:00:60 can consistently produce high rice yield; without phosphorus becoming deficient. The possibility of alternating 80:00:60 and 80:60:60 in different growing seasons should also be checked to minimize chances of phosphorus deficiency in rice fields.

Table 4. Effect of different fertilizer regimes on yield ( $t ha^{-1}$ ) of two rice varieties planted in the first and the second season, between August 2016 and February 2018 at MIAD.

Treatments	First season			Second season		
	Yield ( $t ha^{-1}$ )		Means	Yield ( $t ha^{-1}$ )		Means
N: P <sub>2</sub> O <sub>5</sub> : K <sub>2</sub> O	Bs 370	Bw 196		Bs 370	Bw 196	
00:00:00	2.42	3.27	2.85	3.96	4.44	4.2
60:40:40	3.95	3.83	3.89	4.74	5.98	5.36
80:60:60	4.36	5.7	5.03	4.71	6.12	5.41
100:80:80	4.19	4.22	4.21	4.82	6.13	5.48
60:40:00	3.72	4.06	3.89	3.54	4.68	4.11
80:60:00	3.6	4.33	3.96	3.93	4.69	4.31
100:80:00	3.7	3.41	3.56	3.63	5.13	4.38
60:00:40	3.71	4.12	3.92	4.7	5.18	4.94
80:00:60	4.16	5.82	4.99	4.97	6.26	5.62
100:00:80	3.67	5.09	4.38	4.57	6.08	5.33
00:40:40	3.1	3.46	3.28	4.1	5.91	5.01
00:60:60	4.27	3.34	3.81	4.3	5.72	5.01
00:80:80	3.79	3.82	3.81	4.4	5.5	4.95
Mean	3.74	4.19	3.97	4.34	5.52	4.93
Fpr V		0.02			<.001	
Fpr F		<.001			<.001	
Fpr F*V		0.17			0.34	
LSD 0.05 V		0.34*			0.21*	
LSD 0.05 F		0.85*			0.53*	
LSD 0.05 F*V		NS			NS	
CV %		18.53			9.23	

Fpr- F probability, LSD- Least significant difference at  $P_{0.05}$ , V-variety, F-fertilizer regimes, F\*V- interaction between Variety and Fertilizer regimes, NS- Not significant at  $P_{0.05}$ , \* -significant at  $P_{0.05}$ , CV- co-efficient of variation Ratio of N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O refers to N kg ha<sup>-1</sup>: P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>: K<sub>2</sub>O kg ha<sup>-1</sup>

## Conclusion

Increased fertilizer application resulted in increased plant height, number of tillers<sup>-1</sup> hill, length of the panicles and grain yield of rice. The highest grain yield was reported in fertilizer regimes: 80:60:60, 80:00:60, 100:00:80 and 100:80:80. 80:00:60 is the best fertilizer regime for rice production in Mwea.

## References

Dobermann, A. and Fairhurst, T. 2000. Rice: Nutrient disorders and nutrient management. Potash and Phosphate Institute (PPI), Potash Institute of Canada (PPIC) and International Rice Research, Singapore. pp. 13-82.

FAOSTAT. 2012. Food and Agriculture Organization Corporate Statistical Database. <http://Faostat.fao.org> (accessed on July 30<sup>th</sup>, 2017).

IRRI. 2014. Standard Evaluation System for rice (SES), 5<sup>th</sup> edition. Los Baños (Philippines): International Rice Research Institute. pp. 11-14.

MOA. 2009. National rice development strategy 2008-2018. Information and documentation services, Kenya Agricultural Research Institute, Ministry of Agriculture, Nairobi p.6.

Muhunyu, J.G. 2012. Is doubling rice production in Kenya by 2018 achievable? *J. Dev. Sust. Agric.* 7: 46-54.

Ndiiri, J.A. Mati, B.M., Home, P.G. and Odongo, B. 2013. Water productivity under the system of rice intensification (SRI) from experimental plots and farmer surveys in Mwea, Kenya. *Taiwan Water Conserv.* 61 (4): 63-75.

Ngige, K.J. 2004. An economic analysis of rice production in Mwea irrigation scheme. M.Sc. Thesis, School of Agriculture, University of Nairobi. 119p.

Nyamai, M.B., Mati, M., Home, P.G., Odongo, B., Wanjoqu, R. and Thuranira, E.G. 2012. Improving land and water productivity in basin rice cultivation in Kenya through System of Rice Intensification. *Agric. Eng. Int. CIGR J.* 14 (2): 1-9.

Onyango, A.O. 2014. Exploring options for improving rice production to reduce hunger and poverty in Kenya. *World Environ.* 4(4): 172-179.

Plenet, D. Mollier, A. and Pellerin, S. 2000. Growth analysis of maize field crops under phosphorus deficiency. Radiation use efficiency, biomass accumulation and yield components. *Plant and Soil.* 224: 259-272.

Saito, K. Nelson, A. Zwart, S.J. Niang, A., Sow, A. Yoshida, H. and Wopereis, M.C. 2013. Towards a Better Understanding. Realizing Africa's Rice Promise, Centre for Agriculture and Bioscience International. 188p.

Sarkar, R.K. and Malik, G.C. 2001. Effect of foliar spray of KNO<sub>3</sub> and Ca (NO<sub>3</sub>)<sub>2</sub> on grass pea (*Lathyrus sativus* L.) grown in rice fallows. *Lathyrus Lathyrism News.* 2: 47-48.

Yoshida, S. 1981. Fundamentals of rice crop science. International Rice Research Institute (IRRI), Manilla, Philippines. pp. 41-248.