



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.



Nutrition, cognition and social emotion among preschoolers in poor rural areas of South Central China: status and correlates

by Xinghua Liu, Kevin Chen, Chengfang Liu, Zimeiyi Wang, and Renfu Luo

Copyright 2021 by Xinghua Liu, Kevin Chen, Chengfang Liu, Zimeiyi Wang, and Renfu Luo. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

Nutrition, cognition and social emotion among preschoolers in poor rural areas of South Central China: status and correlates

Introduction

Although there is evidence that food assistance programs for school aged children have led to improved cognition abilities and greater educational achievement [1-2], little is known regarding whether such food assistance programs at the preschool level can achieve similar results. This has induced a heated discussion on whether the current free nutritious lunch program in rural China should be further expanded to the preschool level [3]. To settle this dispute, it calls for evidence on the prevalence of undernutrition among preschool children as well as the relationship between undernutrition and cognitive and socio-emotional development, which are important components of human capital [1]. Policymakers in China rely on this type of information to enhance the human capital foundation of the entire population to achieve its lofty goal of high quality development.

A large body of literature have examined the relationship between nutrition and child development [4-6], but little evidence comes from the poor, rural areas of China, where the majority of its vulnerable children reside. This is mainly due to the lack of data on both the nutrition status and developmental outcomes of children in those disadvantaged areas.

Although several existing studies focus on Chinese children, evidence for poor rural Chinese preschoolers is sparse. For example, Huang et al. (2013) find that the height-

for-age z score, as an indicator for postnatal growth is positively associated with IQ among over 8,000 Chinese children aged 4-7 years [7]. In addition, a couple of studies found Chinese preschoolers with higher hemoglobin (Hb) level tend to perform better in cognitive test and behavioral observations [8-9].

China provides an ideal setting for studying the linkage between child nutritional status and other developmental outcomes for two reasons. First, as the world's largest developing country, China has put great efforts in poverty reduction and enhancing its future growth by combating child malnutrition and investing in child development through interventions such as nutritious lunch program. Second, the prevalence of stunting, wasting, and anemia among children are still high, with children in poor, rural areas of China suffering disproportionately [10].

In this paper, we aim to provide additional evidence on the relationship between nutritional factors and developmental outcomes among preschoolers in poor rural China. To achieve this goal, we have three objectives. First, we will document the nutritional status among preschoolers in the study areas, thus better understanding the severity of undernutrition among preschoolers in poor areas of rural China. Second, we will document the levels of cognitive ability and social emotion performance among our sample children in order to assess how children in poor rural areas fare in terms of these measured outcomes. Finally, we will examine the linkages between nutritional indicators and preschoolers' developmental outcomes.

Methods

Sample Selection

We collected the data in September 2018 as part of the baseline survey of a preschool nutritious lunch pilot program launched by the Xiangxi prefecture government, with support from the World Food Program (WFP). The survey was conducted in two then nationally designated poverty counties in the Xiangxi Autonomous Prefecture, Hunan Province in the south central part of China. That said, our sample includes regions that were poor and populated by ethnic minorities, the subpopulations that are potentially at higher risk for childhood undernutrition. The survey was conducted prior to any intervention associated with the pilot program, and thus the intervention is not the focus of this study.

The sample included twenty-six preschools, which were randomly sampled from fifteen townships across the two project counties. Within each sample preschool, all children aged three or five years who attended the preschool on the survey day were included in the sample. In total, we surveyed 1,334 preschoolers. Primary caregivers of the children (mostly parents or grandparents) were interviewed one on one, face to face by trained enumerators. After excluding observations with missing data on cognition test, we have 1,293 children in our final sample for further analysis.

Data collection

For the purpose of this study, we draw on four types of information collected by the

survey team: 1) socioeconomic data from caregivers; 2) measures of child health, including height, weight and hemoglobin levels; 3) scores on a test of cognitive ability; and 4) scores on a test of socio-emotional skills.

Socioeconomic data. Administered by trained enumerators from local universities, the socioeconomic survey contains detailed information on the demographic and socioeconomic status of both sample children and their households. Questions were designed to learn each household's composition and economic status (measured as the possession status of a list of 13 durable assets). Caregiver's parenting style was evaluated using the Chinese version of the 27-item Parenting Style & Dimensions Questionnaire (PSDQ). The nutrition knowledge of caregiver was tested by asking them a battery of 11 questions regarding the sources and functions of macro- and micronutrients.

Health survey. We focus on three health indicators: height, weight and hemoglobin concentrations (Hb). Height and weight measurements were carried out by professional nurses on site following the WHO standard protocol [11]. The nursing team was trained to ensure that the weighing station was set up on level ground to ensure accuracy of the equipment. The children were measured in light clothing without shoes, hats or accessories. Height was recorded to the nearest 0.1cm with standard tape measure, and weight was recorded to the nearest 0.1kg using calibrated electronic scale. Children's Hb concentrations were measured on-site by finger-prick

blood test with HemoCue Hb 201+system (HemoCue, Inc., Angelholm, Sweden) in the field.

Cognitive ability. Cognitive ability was assessed using a battery of four sub-tests taken from the Mandarin-language version of the Wechsler Preschool and Primary Scale of Intelligence Fourth Edition (WPPSI-IV). As the latest version, WPPSI-IV was culturally adapted, translated and edited into simplified Chinese and validated for Chinese children. WPPSI-IV has two sub-versions, one is for preschoolers aged between 30 months and 47 months old (the younger cohort version) whereas the other for those aged between 48 months and 83 months (the elder cohort version). Since prior studies suggest that working memory and verbal comprehension are those areas of cognitive ability most likely to be affected by malnutrition [12-13], we focused on measuring these two outcomes. In WPPSI-IV, the working memory index (WMI) is assessed through two core subtests: Picture Memory and Zoo Locations. The verbal comprehension index (VCI) is assessed through two other subtests: Information and Picture Naming for the younger cohort whereas Information and Similarities for the elder cohort. Raw scores obtained from these core subtests were converted to age-scaled index scores using tables of norms from the official WPPSI-IV administration and scoring manual for China. Each of the sample preschoolers was individually administered for the four core sub-tests by trained examiners.

Socio-emotional skills. Children's socio-emotional skills were assessed through the caregiver-reported Mandarin Language Strengths and Difficulties Questionnaire

(SDQ). As a valid behavioral screening questionnaire [14], the SDQ has been extensively adopted by researchers and clinicians in Europe [15], Australia [16], USA [17] and China [18]. The questionnaire contains twenty-five items to assess children's emotions, behaviors and relationships. The SDQ includes five subscales: peer problems, hyperactivity/inattention symptoms, emotional problems, conduct problems and prosocial behavior. Each subscale includes five items.

Ethics Statement

This study received ethical approval from the International Food Policy Research Institute Institutional Review Boards (IRB) (DSG-18-0837). Written informed consent was obtained from all legal guardians of children and school staff involved in the study.

Statistical analysis

Height and weight were used to construct height-for-age z-scores (HAZ), weight-for-height z-scores (WHZ) and (BMI)-for-age z-score (BmiAZ) using the WHO Child Growth Standards for children under 5 years old and WHO Growth reference data for 5-19 year. We followed internationally recognized cutoffs [19] to consider children whose HAZ or WHZ (applies for children under 61 months) / BmiAZ (applies for children over 61 months) to fall more than two standard deviations below the international mean to be stunted and wasted, respectively.

Anemia status was determined based on finger prick blood analysis for hemoglobin (Hb). Following internationally accepted standards, anemia was defined as Hb<110 g/L for children under 5 years old and 115 g/L for children older than 5 years [20].

Raw scores obtained from the four core subtests of the WPPSI-IV were converted to aged-scaled index scores using tables of norms in the Mandarin version of the WPPSI-IV administration and scoring manual. Two index scores are considered for analysis: Working Memory Index (WMI) and Verbal Comprehension Index (VCI). Scores are then divided into internationally-recognized ranges. A score of 90–110 is considered “average”; a score of 80–89 is considered “low average”; a score of 70–79 is considered “borderline”; and a score of below 70 is considered “extremely low” and at risk for intellectual disabilities or mental retardation.

As mentioned earlier, SDQ contains five subscales to assess children’s socio-emotional risks. The score of each subscale ranges from 0 to 10, with higher scores indicating more problems, except for the prosocial behavior subscale, for which a lower score indicates more problems. The first four subscales aforementioned measure children’s behavioral problems and can be summed to generate a total difficulties score (ranged from 0-40). The fifth subscale was designed to evaluate children’s prosocial behavior attributes, such as the willingness to help others and to share. Total difficulties score above 16 and prosocial score below 5 were considered abnormal, following cutoff values described in an earlier study from China [21].

All statistical analyses were performed using STATA version 14.0. P-values below

0.05 were considered statistically significant. We conducted three types of statistical analyses. First, we started by comparing the subgroup population differences in descriptive statistics of nutritional, cognitive, and socio-emotional status along various dimensions using student's t-test. Second, a pairwise correlation analysis between nutritional status and developmental outcomes was conducted, which compared WPPSI and SDQ scores for anemic and non-anemic, stunted and non-stunted, wasted and non-wasted preschoolers, respectively. Values were presented as mean \pm SD or as percentage for categorical variables. Third, we examined relationship between nutritional indicators and key developmental outcome variables (WMI, VCI, prosocial score, and total difficulties score) by using STATA'S multiple linear regression model. Previous studies suggest that undernutrition is generally associated with many socioeconomic and biological disadvantages which can themselves affect children's development and thus may confound the nutrition-development link [4]. Thus, we included a vector of child, parent, caregiver and household-level characteristics as potential confounders in our correlational analysis. Definitions of key variables to be used in the rest of the paper are presented in Table A in S1 Table A.

Results

Table 1 presents descriptive statistics for the variables used in this analysis. Of the 1,293 children, 48.1 percent were girls. 40.2 percent and 48.6 percent of the children aged 3 and 5 respectively. Ethnic minorities accounted for 88.6 percent of the sample,

a ratio similar to that of Xiangxi prefecture. 863 or 70.9 percent children are defined as left-behind children. The majority of parents have an education level of junior high school or below.

Table 1. Descriptive statistics and prevalence of child undernutrition by characteristics.

		Full sample	Anemia no. (%)	Stunting no. (%)	Wasting no. (%)
Full sample		1293	444 (33.36)	141 (10.59)	23 (1.73)
Child characteristics					
Gender	Female	622	200 (32.21)	61 (9.82)	14 (2.25)
	Male	671	231 (34.48)	76 (11.33)	9 (1.34)
Ethnicity	Minority	1,145	372 (32.52)	119 (10.40)	22 (1.92)
	Han	147	59 (40.41)	18 (12.24)	1 (0.68)
Age	3 years old	520	159 (30.58) ^a	49 (9.42)	5 (0.96) ^a
	5 years old	628	237 (37.80) ^a	71 (11.31)	17 (2.71) ^a
Low birth weight	Yes (< 2.5 kg)	87	29 (32.22)	20 (22.22) ^a	5 (5.56) ^a
	No (≥ 2.5 kg)	1187	408 (33.42)	119 (9.74) ^a	17 (1.39) ^a
Breastfeeding	Inadequate (< 6 months)	439	162 (36)	48 (10.67)	10 (2.22)
	Optimal (≥ 6 months)	854	286 (32.01)	93 (10.54)	13 (1.47)
Left-behind status	Both parents around	354	126 (34.62)	37 (10.14)	5 (1.37)
	One parent out	248	75 (29.30)	32 (12.5)	5 (1.95)
	Both parents out	615	216 (34.18)	65 (10.28)	11 (1.74)
Parent characteristics					
Father's education	Junior high school or below	1050	377 (34.91) ^a	122 (11.29)	20 (1.85)
	Senior high school or above	243	67 (26.69) ^a	19 (7.57)	3 (1.20)
Mother's education	Junior high school or below	982	331 (32.80)	105 (10.41)	19 (1.88)
	Senior high school or above	311	113 (35.09)	36 (11.16)	4 (1.24)
Caregiver characteristics					
Caregiver's nutrition knowledge	Above average score	710	239 (32.69)	58 (11.34)	13 (1.78)
	Below average score	583	205 (34.17)	83 (9.67)	10 (1.67)
Caregiver's parenting style	Above average score	692	237 (33.24)	70 (9.82)	11 (1.54)
	Below average score	601	207 (33.50)	71 (11.47)	12 (1.94)
Household characteristics					
Household size	Small size (≤ 3 people)	5	2 (40)	0 (0)	0 (0)
	Medium size (4 people)	8	1 (11.11)	0 (0)	0 (0)
	Large size (≥ 5 people)	1280	441 (33.49)	141 (10.70)	23 (1.75)
Siblings	No sibling	396	138 (33.82)	38 (9.31)	4 (0.98)
	With siblings	897	306 (33.15)	103 (11.15)	19 (2.06)
Household durable asset	The lowest 1/3 quantile	432	151 (34.95)	57 (13.19)	8 (1.85)
	The middle 1/3 quantile	486	171 (35.26)	49 (10.10)	7 (1.44)
	The highest 1/3 quantile	375	109 (29.14)	31 (8.27) ^b	8 (2.13)

Source: authors' survey.

Note: ^a Significantly different from the other group; ^b Significantly different from the lowest quantile group.

Nutritional status

Table 1 also presents the prevalence of anemia, stunting and wasting among our sample children. The mean Hb level of the sample children was 115.8 ± 0.30 , with 33.4 percent children classified as anemic, which is the most prevalent types of undernutrition in our sample followed by stunting (10.6 percent) and wasting (1.7 percent). The mean HAZ was -0.8 ± 0.03 . The mean WHZ for children under 61 months was 0.02 ± 0.03 , whereas the mean BmiAZ for children older than 61 months was -0.18 ± 0.04 .

Child nutritional status differs significantly by child's age and birth weight, father's education, and households' durable assets ownership. Specially, lower prevalence of anemia and wasting was observed among children in the five years old age group. In addition, children with normal birth weight had lower incidence of being stunted or wasted. Moreover, children whose fathers had better education level had lower anemia rate, and children whose family possessed more durable assets had lower stunting rate (Table 1).

Developmental outcomes

Overall, 53.67 percent of the sample children had delay in at least one of the developmental domains measured in this study. As shown in Table 2, WMI and VCI scores of sample children averaged 90.43 ± 0.37 and 86.02 ± 0.35 , respectively. A total of 21.0 percent children had a WMI that was either extremely low or borderline low, whereas the figure was 30.5 percent for VCI.

The average prosocial score and total difficulties score were 6.83 ± 0.06 and 12.35 ± 0.13 , respectively. The prevalence of abnormal prosocial score was 15.1 percent. 21.0 percent children were categorized as abnormal in terms of total difficulties score.

Table 2 also indicates that preschoolers of certain characteristics were significantly more likely to have lower cognition or SDQ scores. First, for WMI, preschoolers with certain characteristics showed lower WMI score, including 3-year-old children, children with a birth weight lower than 2.5 kg, children whose father had lower education level, children whose caregivers followed a parenting style that was low in demands and responsiveness, children without siblings, as well as those whose family had fewer durable assets. In contrast, for VCI, breastfeeding duration shorter than six month was statistically correlated with poorer VCI performance. Unlike WMI, whether the preschoolers were only child or not did not seem to play a role for our sample preschoolers.

Second, in terms of prosocial score, preschoolers with the following characteristics exhibited more problems, including 3 year old, father's educational attainment and caregiver's parenting style. In addition, 5-year-old preschoolers and single-child preschoolers tended to perform significantly worse in terms of total difficulty scores compared to those 3 years old and those with siblings.

Table 2. Characteristics and distribution of sample children by development indicators.

WMI	VCI	Prosocial score	Total difficulties score
-----	-----	-----------------	--------------------------

Full sample		90.43±0.37	86.02±0.35	6.83±0.06	12.35±0.13
Child characteristics					
Gender	Female	90.31±0.53	86.23±0.52	7.11±0.09 ^a	12.11±0.19
	Male	90.55±0.53	85.82±0.49	6.57±0.08 ^a	12.57±0.18
Ethnicity	Minority	90.60±0.40	85.97±0.38	6.85±0.06	12.40±0.14
	Han	89.09±1.06	86.35±0.95	6.69±0.18	11.91±0.40
Age	3 years old	87.42±1.06 ^a	83.59±0.65 ^a	6.44±0.10 ^a	13.22±0.21 ^a
	5 years old	92.74±0.40 ^a	88.08±0.43 ^a	7.14±0.08 ^a	11.56±0.18 ^a
Low birth weight	Yes (< 2.5 kg)	86.52±1.28 ^a	82.05±1.40 ^a	6.69±0.25	12.37±0.47
	No (≥2.5 kg)	90.68±0.39 ^a	86.26±0.37 ^a	6.84±0.06	12.35±0.14
Breastfeeding	Inadequate (< 6 months)	89.47±0.65	84.99±0.62 ^a	6.76±0.10	12.50±0.23
	Optimal (≥6 months)	90.93±0.46	86.55±0.43 ^a	6.86±0.07	12.27±0.16
Left-behind status	Both parents around	91.03±0.70	87±0.68	6.79±0.11	12.33±0.25
	One parent out	90.33±0.94	85.13±0.81	6.84±0.14	12.74±0.30
	Both parents out	90.25±0.52	86.07±0.51	6.84±0.09	12.22±0.18
Parent characteristics					
Father's education	Junior high school or below	89.94±0.41 ^a	85.64±0.39 ^a	6.71±0.07 ^a	12.43±0.15
	Senior high school or above	92.54±0.93 ^a	87.63±0.82 ^a	7.34±0.12 ^a	12.01±0.29
Mother's education	Junior high school or below	90.31±0.42	85.89±0.40	6.87±0.07	12.25±0.15
	Senior high school or above	90.81±0.82	86.41±0.77	6.70±0.13	12.66±0.28
Caregiver characteristics					
Caregiver's nutrition knowledge	Above average score	90.61±0.53	86.03±0.50	6.94±0.08	12.26±0.17
	Below average score	90.22±0.52	86±0.50	6.70±0.09	12.45±0.20
Caregiver's parenting style	Above average score	91.13±0.51 ^a	86.89±0.48 ^a	7.30±0.08 ^a	11.97±0.18 ^a
	Below average score	89.63±0.54 ^a	85.01±0.53 ^a	6.29±0.09 ^a	12.78±0.19 ^a
Household characteristics					
Household size	Small size (≤3 people)	88.8±6.92	77.6±6.85	6.20±1.02	13.80±1.02
	Medium size (4 people)	90.88±3.48	83±3.63	7.00±0.71	12.50±1.30
	Large size (≥5 people)	90.44±0.38	86.07±0.36	6.83±0.06	12.34±0.13
Siblings	No sibling	89.23±0.67 ^a	86.17±0.67	6.92±0.11	12.74±0.23 ^a
	With siblings	90.96±0.45 ^a	85.95±0.42	6.79±0.07	12.18±0.16 ^a
Household durable asset	The lowest 1/3 quantile	88.74±0.61	84.59±0.60	7.00±0.11	12.45±0.23
	The middle 1/3 quantile	91.02±0.63 ^b	86.76±0.56 ^b	6.79±0.10	12.49±0.20
	The highest 1/3 quantile	91.61±0.70 ^b	86.70±0.69 ^b	6.68±0.12 ^b	12.05±0.25

Source: authors' survey.

Note: ^a Significantly different from the other group; ^b Significantly different from the lowest quantile group.

Association between nutrition and developmental outcomes

Table 3 shows cross-sectional correlation between nutritional status (anemia, stunting, wasting) and child developmental outcome variables. Anemia was negatively associated with WMI when controlling for age of the child (coefficient: -1.52; 95%

CI: -2.7, -0.34; P=0.013). Stunting was statistically correlated with a -2.88 (95% CI: -4.76, -0.99; P=0.004) and -3.84 (95% CI: -5.51, -2.17; P=<0.001) average difference in WMI and VCI, respectively. After adjusting for confounders, the magnitude of association between anemia and WMI was increased to -1.62 (95% CI: -2.98, -0.26; P=0.021). Conversely, for the correlation between stunting and VCI, the magnitude was reduced to -3.01 (95% CI: -4.71, -1.3; P=0.001). Noticeably, wasting was not statistically correlated with any developmental outcome variables, and none of the three nutritional indicators were statistically associated with socio-emotional performance.

Table 4 and Table 5 show correlation between continuous nutritional indicators and developmental outcome variables for children aged under and over 5 years, respectively. First, for children under 5 years old (Table 4), HAZ was positively associated with cognitive outcomes, and an increment of one unit in HAZ was associated with 1.81-point (95% CI: 0.73, 2.88; P=0.002) increase in WMI and 1.61-point (95% CI: 0.54, 2.68; P=0.005) increase in VCI. When adjusting for confounders, the magnitude of associations between HAZ and WMI and VCI remained similar and the significance remained unchanged (coefficient: 1.84; 95% CI: 0.64, 3.04; P=0.004, for WMI; coefficient: 1.55; 95% CI: 0.42, 2.68; P=0.009, for VCI). In addition, one unit increase in HAZ was associated with 0.16-point increase in prosocial score (95% CI: 0.02, 0.31; P=0.029). Note that neither Hb level nor WHZ showed significant association with any developmental outcome variables. Second, for children over 5 years old (Table 5), we did not find any significant correlation

between the three continuous nutritional status with the four developmental outcome variables.

Table 3. Cross-sectional associations between anemia, stunting, wasting, and child developmental outcomes, Values are coefficient [95% CI].

	WMI		VCI		Prosocial score		Total difficulties score	
	Age-adjusted	Multivariate-adjusted	Age-adjusted	Multivariate-adjusted	Age-adjusted	Multivariate-adjusted	Age-adjusted	Multivariate-adjusted
Anemia	-1.52** [-2.7,-0.34]	-1.62** [-2.98, -0.26]	-0.9 [-2.31,0.51]	-1.04 [-2.32, 0.23]	-0.06 [-0.33,0.21]	-0.06 [-0.31, 0.2]	0.14 [-0.45,0.73]	-0.04 [-0.73, 0.66]
Stunting	-2.88*** [-4.76,0.99]	-2.08 [-4.17, 0.02]	-3.84*** [-5.51,-2.17]	-3.01*** [-4.71, -1.3]	-0.13 [-0.53,0.28]	-0.01 [-0.42, 0.4]	0.01 [-1.11,1.12]	0.002 [-1.23, 1.23]
Wasting	3.95 [-1.71,9.61]	4.09 [-2.63, 10.8]	0.61 [-4.03,5.25]	0.88 [-5.18, 6.93]	-0.25 [-1.07,0.57]	-0.13 [-0.8, 0.54]	0 [-1.97,1.97]	0.24 [-1.64, 2.13]

Source: authors' survey.

Note: *, **, *** Statistically significant at the 10%, 5%, and 1% level, respectively.

Table 4. Cross-sectional associations between Hb, HAZ, WHZ, and child developmental outcomes for children younger than 61 months of age, Values are coefficient [95% CI].

	WMI		VCI		Prosocial score		Total difficulties score	
	Age-adjusted	Multivariate-adjusted	Age-adjusted	Multivariate-adjusted	Age-adjusted	Multivariate-adjusted	Age-adjusted	Multivariate-adjusted
Hb	0.06 [-0.07,0.19]	0.06 [-0.07,0.19]	0.06 [-0.07,0.19]	0.07 [-0.05, 0.20]	-0.004 [-0.03,0.02]	-0.01 [-0.02, 0.01]	0.02 [-0.02,0.06]	0.03 [-0.01, 0.06]
HAZ	1.81*** [0.73,2.88]	1.84*** [0.64,3.04]	1.61*** [0.54,2.68]	1.55*** [0.42, 2.68]	0.15 [-0.03,0.33]	0.16** [0.02, 0.31]	0.22 [-0.28,0.71]	0.06 [-0.44, 0.57]
WHZ	0.37 [-0.95,1.68]	0.23 [-1.15,1.6]	0.42 [-0.79,1.64]	0.41 [-0.79, 1.62]	0.09 [-0.14,0.31]	0.14 [-0.09, 0.37]	0.26 [-0.27,0.8]	0.22 [-0.33 0.78]

Source: authors' survey.

Note: *, **, *** Statistically significant at the 10%, 5%, and 1% level, respectively.

Table 5. Cross-sectional associations between Hb, HAZ, BmiAZ, and child developmental outcomes for children older than 61 months of age, Values are coefficient [95% CI].

	WMI		VCI		Prosocial score		Total difficulties score	
	Age-adjusted	Multivariate-adjusted	Age-adjusted	Coefficient [95% CI]	Age-adjusted	Coefficient [95% CI]	Age-adjusted	Coefficient [95% CI]
Hb	0.001 [-0.09,0.09]	0.004 [-0.09,0.1]	0.004 [-0.1,0.11]	0.001 [-0.11,0.11]	0.01 [-0.01,0.02]	0.01 [-0.004,0.02]	-0.03 [-0.06,0]	-0.03 [-0.07,0.01]
HAZ	0.96 [-0.07,1.99]	0.48 [-0.76,1.72]	1.51 [0.43,2.59]	0.91 [-0.53,2.35]	0.06 [-0.15,0.26]	-0.06 [-0.27,0.15]	0.07 [-0.39,0.53]	0.1 [-0.43,0.62]
BmiAZ	-0.27 [-1.12,0.59]	-0.14 [-1.23,0.94]	-0.44 [-1.1,0.22]	-0.39 [-0.92,0.14]	-0.07 [-0.26,0.12]	-0.1 [-0.27,0.08]	0.12 [-0.24,0.48]	0.08 [-0.3,0.46]

Source: authors' survey.

Discussion and conclusions

This study showed the status of undernutrition and retarded development as well as their relationships among 1,293 preschoolers in poor rural areas of South Central China. We observed that 33 percent of the sample children were anemic. Although this incidence of anemia was consistent with previous studies in rural China [22-23], it was more than 10 percent higher than the Chinese national average for children under 5 in 2016 (21.4%) [24]. In addition, compared to a stunting incidence of 28 percent in Guizhou in 2013 suggested by a prior study [25], our study indicated that the figure was 10.6 percent for our sampled preschoolers. Moreover, 53.67 percent of the sample children had delay in at least one of the developmental domains measured in this study.

Our study also found that the incidence of undernutrition and developmental delay differs along several important household- or individual-level characteristics, which provided further support for nutrition or development inequality among Chinese preschool children. Specially, children born with low birth weight were more likely to be affected by stunting, wasting as well as to be adversely affected by poor cognitive abilities. This aligns with the broad range of literature which indicate that low birth weight predicts compromised developmental outcomes [26-28]. Moreover, father's education matter for both child anemia status and WPPSI and prosocial scores. This is congruent with previous findings, suggesting that lower socioeconomic status is linked with lower IQ and executive functions [29-30]. Similarly, consistent with

previous findings [31-32], our study identified that parenting style which is low in demands and responsiveness is linked with poor cognitive and prosocial abilities. Additionally, we found that single child performed significantly worse in terms of both WMI and total difficulties score than their with-siblings peers. While this finding is in contrast with a couple of previous studies [33-34], it is consistent with a more recent study in China which suggests that children with siblings perform better in terms of cognition than their only-child counterparts [35]. Jointly, our findings provide suggestive evidence supporting that children from certain background tend to experience worse nutritional and developmental outcomes than their counterparts. Children of these characteristics tend to be more vulnerable and thus deserve special attention when designing and implementing policy interventions.

Our study further examined the association between three nutritional indicators and children's cognitive and socio-emotional performance. As expected, strong relationships were found between anemia and cognition development and between stunted growth and cognitive skills. What was unexpected, however, was our lack of identifying relationships between anemia and socio-emotional status and between wasting or WHZ and any developmental outcome variables.

First, consistent with previous findings [36-38], we found that anemic children were more likely to suffer suboptimal cognition development than non-anemic peers. However, unlike previous studies [39-40], our findings did not support strong associations of anemia with socio-emotional status. The lack of such associations is

partly because our findings were obtained after controlling for a wide range of socio-demographic and socioeconomic characteristics, suggesting that these characteristics may have a greater effect on child socio-emotional outcomes.

Second, in term of correlation between stunting and developmental outcomes, our results suggest that stunted growth was negatively associated with cognition development among preschoolers. These results align with previous studies which consider stunting as an important predictor for poor child cognitive skills [37-38, 41]. Moreover, consistent with previous studies [42-43], our study did not find any association between stunting and child social emotional performance.

Third, in term of correlation between wasting and developmental outcomes, we found no significant association of wasting or WHZ with any outcome variables. This is congruent with previous findings which suggested an inconsistent relationship between WHZ and child development indicators [44]. Nevertheless, some other studies suggested that WHZ was negatively associated with child's performance in cognitive tests [38] or developmental delay [45]. Thus, our findings contradicted with theirs. Nevertheless, we add to this strand of literature by providing additional evidence among preschoolers from poor rural China.

Our study is not without limitations. First, this study relies on cross-sectional data and uses simple statistical methods to identify associations; therefore, the results should be interpreted as correlation rather than causality. Second, due to the cross-sectional nature of our data, we are unable to identify the underlying channels behind the

linkages we have observed through this study. Investigating these potential channels could be fruitful avenues for future research. Nevertheless, this paper contributes to the understanding of the relationship between childhood nutrition and cognitive and socio-emotional achievement among preschool children. Our findings suggest that anemia and stunting are the strongest risks for cognitive performance among our sampled preschool-aged children. Undernutrition among preschoolers poses challenges to public health in China's underdeveloped areas, and there is a need to develop effective health interventions targeting preschool-aged children in these areas. More importantly, these results provide new suggestive evidence regarding whether school-level free nutritious lunch programs should be expanded to the preschool education level, and thus facilitating evidence-based decision making surrounding this important policy debate .

Supporting information

S1 Table A. Variable Definitions

S1 Table A. Variable Definitions

Variable	Description
<i>Nutritional indicators</i>	
Hb	Hemoglobin concentrations (g/L)
Anemic	Hb<110 g/L(<61 months) Hb<115 g/L(\geq 61 months)
HAZ	Height-for-age z-score
WHZ	Weight-for-height z-score
BMIAZ	BMI-for-age z-score
Stunting	More than two standard deviations below the WHO child growth reference data(1=yes, 0=no)
Wasting	More than two standard deviations below the WHO child growth reference data(1=yes, 0=no)
<i>Cognitive ability</i>	
Working memory index (WMI)	Standardized score on the working memory module of the WPPSI
Verbal comprehension index (VCI)	Standardized score on the verbal comprehension module of the WPPSI
<i>Socio-emotional performance</i>	
Total difficulties score	Total score of the first four subscale of SDQ (ranging from 0-40)
Prosocial score	Total score of the fifth subscale of SDQ (ranging from 0-10)
<i>Correlates</i>	
<i>Child characteristics</i>	
Female	Child is female (1=yes, 0=no)
Age	Age of child, years
Ethnicity	Child is non-Han ethnic minority (1=yes, 0=no)
Low birth weight	Birth weight was less than 2.5 kg (1=yes, 0=no)
Breastfeeding duration	Breastfeeding duration was less than 6 months (1=yes, 0=no)

Left-behind status	Children were defined as left-behind if one or both of their parents worked outside their hometown of household registration for at least 6 months
<i>Household characteristics</i>	
Household size	No. of family members (person)
Siblings	No. of siblings (person)
Pieces of durable assets	Pieces of durable assets owned by the household
<i>Parent characteristics</i>	
Mother educational attainment	Senior high school or above (1=yes, 0=no)
Father educational attainment	Senior high school or above (1=yes, 0=no)
<i>Caregiver characteristics</i>	
Primary caregiver's nutritional knowledge	Above the mean score (1=yes, 0=no)
Primary caregiver's authoritative parenting style score	Above the mean score (1=yes, 0=no)

References

1. Frisvold DE. Nutrition and cognitive achievement: An evaluation of the School Breakfast Program. *J Public Econ.* 2015; 124:91–104.
2. Meyers AF, Sampson AE, Weitzman M, Rogers BL, Kayne H. School Breakfast Program and school performance. *Am J Dis Child.* 1989;143(10):1234-9.
3. Xinhua Net. Nutritious lunch program should be expanded to preschool level.2016. http://www.xinhuanet.com/politics/2016-06/17/c_129069975.htm. Last accessed on December 23, 2019.
4. Alderman H, Fernald L. The Nexus Between Nutrition and Early Childhood Development. *Annu Rev Nutr.* 2017; 37: 447-476.
5. John CC, Black MM, Nelson CA 3rd. Neurodevelopment: The Impact of Nutrition and Inflammation During Early to Middle Childhood in Low-Resource Settings. *Pediatrics.* 2017; 139 (Suppl 1):S59-S71.
6. Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev.* 2014;72 (4):267–84.
7. Huang C, Martorell R, Ren A, Li Z. Cognition and behavioural development in early childhood: the role of birth weight and postnatal growth. *Int J Epidemiol.* 2013;42(1):160–71.
8. Ai Y, Zhao SR, Zhou G, Ma X, Liu J. Hemoglobin status associated with performance IQ but not verbal IQ in Chinese preschool children. *Pediatr Int.* 2012; 54(5):669–75.
9. Chang S, Wang L, Wang Y, Brouwer ID, Kok FJ, Lozoff B, Chen C. Iron-deficiency anemia in infancy and social emotional development in preschool-aged Chinese children. *Pediatrics.*

2011;127 (4):e927-33.

10. China National Bureau of Statistics. 2018 Chinese children development outline (2011-2020) statistical minitoring report.2019.
http://www.stats.gov.cn/tjsj/zxfb/201912/t20191206_1715751.html. Last accessed on December 23, 2019.
11. de Onis M, Blössner M, Borghi E, Morris R, Frongillo EA. Methodology for estimating regional and global trends of child malnutrition. *Int J Epidemiol.* 2004 Dec; 33(6):1260-70.
12. Liu J, Yang H, Li L, Chen T, Lynn R. An increase of intelligence measured by the WPPSI in China, 1984–2006. *Intelligence.* 2012;40(2):139–44.
13. Whaley SE, Sigman M, Espinosa MP, Neumann CG. Infant predictors of cognitive development in an undernourished Kenyan population. *J Dev Behav Pediatr.* 1998;19(3):169-77.
14. GOODMAN R. Psychometric Properties of the Strengths and Difficulties Questionnaire. *J Am Acad Child Adolesc Psychiatry.* 2001;40(11):1337–45.
15. Wiles NJ, Northstone K, Emmett P, Lewis G. 'Junk food' diet and childhood behavioural problems: results from the ALSPAC cohort. *Eur J Clin Nutr.* 2009; 63(4):491-8.
16. Renzaho AMN, Kumanyika S, Tucker KL. Family functioning, parental psychological distress, child behavioural problems, socio-economic disadvantage and fruit and vegetable consumption among 4–12 year-old Victorians, Australia. *Health Promot Int.* 2011; 26(3):263–75.
17. Deater-Deckard K, Wang Z, Chen N, Bell MA. Maternal executive function, harsh parenting, and child conduct problems. *J Child Psychol Psychiatry.* 2012; 53(10):1084-91.
18. Bao P, Jing J, Jin Y, Hu X, Liu B, Hu M. Trajectories and the influencing factors of behavior problems in preschool children: a longitudinal study in Guangzhou, China. *BMC Psychiatry.* 2016; 16:178.
19. World Health Organization. WHO Child Growth Standards. Geneva, Switzerland: World Health Organization.2006.
20. United Nations Children's Fund, World Health Organization, United Nations University. Iron Deficiency Anemia: Assessment, Prevention and Control, A Guide for Programme Managers. Geneva: World Health Organization. 2001.
21. Du Y, Kou J, Coghill D. The validity, reliability and normative scores of the parent, teacher and self report versions of the Strengths and Difficulties Questionnaire in China. *Child Adolesc Psychiatry Ment Health.* 2008; 2(1):8.
22. Hipgrave DB, Fu X, Zhou H, Jin Y, Wang X, Chang S, Scherbier RW, Wang Y, Guo S. Poor complementary feeding practices and high anaemia prevalence among infants and young children in rural central and western China. *Eur J Clin Nutr.* 2014; 68(8):916-24.
23. Luo R, Shi Y, Zhou H, Yue A, Zhang L, Sylvia S, et al. Micronutrient deficiencies and developmental delays among infants: evidence from a cross-sectional survey in rural China. *BMJ Open.* 2015; 5(10):e008400.

24. World Health Organization. Global health observatory data repository/world health statistics 2019. 2019.

25. Liu C, Luo R, Yi H, Zhang L, Li S, Bai Y, et al. Soil-Transmitted Helminths in Southwestern China: A Cross-Sectional Study of Links to Cognitive Ability, Nutrition, and School Performance among Children. *PLoS Negl Trop Dis.* 2015;9(6):e0003877–e0003877.

26. Hack M, Klein NK, Taylor HG. Long-term developmental outcomes of low birth weight infants. *Future Child.* 1995; 5(1):176-96.

27. Jefferis BJ, Power C, Hertzman C. Birth weight, childhood socioeconomic environment, and cognitive development in the 1958 British birth cohort study. *BMJ.* 2002; 325(7359):305.

28. Harding KL, Aguayo VM, Webb P. Birthweight and feeding practices are associated with child growth outcomes in South Asia. *Matern Child Nutr.* 2018;14(S4):e12650.

29. Noble KG, Houston SM, Brito NH, Bartsch H, Kan E, Kuperman JM, et al. Family income, parental education and brain structure in children and adolescents. *Nat Neurosci.* 2015;18(5):773–8.

30. Zhang H, Lee ZX, White T, Qiu A. Parental and social factors in relation to child psychopathology, behavior, and cognitive function. *Transl Psychiatry.* 2020;10(1):80.

31. Blair C, Raver CC, Berry DJ; Family Life Project Investigators. Two approaches to estimating the effect of parenting on the development of executive function in early childhood. *Dev Psychol.* 2014; 50(2):554-65.

32. Lucassen N, Kok R, Bakermans-Kranenburg MJ, Van IJzendoorn MH, Jaddoe VW, Hofman A, Verhulst FC, Lambregtse-Van den Berg MP, Tiemeier H. Executive functions in early childhood: the role of maternal and paternal parenting practices. *Br J Dev Psychol.* 2015;33(4):489-505.

33. Blake J. Family size and the quality of children. *Demography.* 1981; 18(4):421-42.

34. Blake J. Number of siblings and educational mobility. *Am Sociol Rev.* 1985;50(1):84-94.

35. Zhou H, Mo D, Luo R, Yue A, Rozelle S. Are Children with Siblings Really More Vulnerable Than Only Children in Health, Cognition and Non-cognitive Outcomes? Evidence from a Multi-province Dataset in China. *China World Econ.* 2016;24(3):3–17.

36. Ai Y, Zhao SR, Zhou G, Ma X, Liu J. Hemoglobin status associated with performance IQ but not verbal IQ in Chinese preschool children. *Pediatr Int.* 2012;54(5):669–75.

37. Kordas K, Lopez P, Rosado JL, García Vargas G, Alatorre Rico J, Ronquillo D, Cebrián ME, Stoltzfus RJ. Blood lead, anemia, and short stature are independently associated with cognitive performance in Mexican school children. *J Nutr.* 2004;134 (2):363-71.

38. Paxson C, Schady N. Cognitive Development Among Young Children In Ecuador : The Roles Of Wealth, Health And Parenting. Policy Research Working Papers. The World Bank; 2005. 34 p.

39. Liu J, Raine A. Nutritional status and social behavior in preschool children: the mediating effects of neurocognitive functioning. *Matern Child Nutr.* 2017; 13(2):e12321.

40. Lozoff B, Corapci F, Burden MJ, Kaciroti N, Angulo-Barroso R, Sazawal S, et al. Preschool-Aged Children with Iron Deficiency Anemia Show Altered Affect and Behavior. *J Nutr.* 2007;137(3):683–9.
41. Crookston BT, Dearden KA, Alder SC, Porucznik CA, Stanford JB, Merrill RM, et al. Impact of early and concurrent stunting on cognition. *Matern Child Nutr.* 2011;7(4):397–409.
42. Kang Y, Aguayo VM, Campbell RK, West Jr. KP. Association between stunting and early childhood development among children aged 36–59 months in South Asia. *Matern Child Nutr.* 2018; 14(S4):e12684.
43. Miller AC, Murray MB, Thomson DR, Arbour MC. How consistent are associations between stunting and child development? Evidence from a meta-analysis of associations between stunting and multidimensional child development in fifteen low- and middle-income countries. *Public Health Nutr.* 2016;19(8):1339–47.
44. Sudfeld CR, McCoy DC, Fink G, Muhihi A, Bellinger DC, Masanja H, et al. Malnutrition and Its Determinants Are Associated with Suboptimal Cognitive, Communication, and Motor Development in Tanzanian Children. *J Nutr.* 2015; 145(12):2705–14.
45. van den Heuvel M, Voskuijl W, Chidzalo K, Kerac M, Reijneveld SA, Bandsma R, Gladstone M. Developmental and behavioural problems in children with severe acute malnutrition in Malawi: A cross-sectional study. *J Glob Health.* 2017; 7(2):020416.