



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

JEL: O13, Q11, Q13, Q18

Olajide I. Sanusi¹, Samir K. Safi², Omotara Adeeko³, Mosab I. Tabash⁴

^{1,2}*United Arab Emirates University*

³*University of Pittsburgh*

⁴*Al Ain University*

^{1,2,4}*United Arab Emirates*

³*USA*

FORECASTING AGRICULTURAL COMMODITY PRICE USING DIFFERENT MODELS: A CASE STUDY OF WIDELY CONSUMED GRAINS IN NIGERIA

Purpose. This study highlights the specific and accurate methods for forecasting prices of commonly consumed grains or legumes in Nigeria based on data from January 2017 to June 2020.

Methodology / approach. Different models that include autoregressive integrated moving average (ARIMA), artificial neural networks (ANN), seasonal decomposition of time series by loess method (STLM), and a combination of these three models (hybrid model) were proposed to forecast the sample grain price data. This study uses price data on widely consumed grains, such as white maize, local rice, imported rice, and white beans, in Nigeria from January 2017 to June 2020.

Results. Our result indicates that ARIMA is the best applicable model for white maize and imported rice because it is well fitted to stationary data, as demonstrated in the sample period. The STLM is more appropriate in forecasting white beans. As white beans are highly seasonal in Nigeria, it further explains why the STLM model fits better in forecasting prices. The production of local rice is inconsistent in Nigeria because of erratic rainfall and stiff competition from the importation of rice from other countries. Therefore, and consistent with the analysis, the hybrid model is the best model applicable to local rice because it captures varying trends exhibited in the data.

Originality / scientific novelty. This study suggests most accurate forecasting techniques for specific agricultural commodities in sub-Saharan African countries. It considers forecasting prices of commonly consumed grains and legumes in Nigeria and traded worldwide, such as imported rice, local rice, beans, and maize.

Practical value / implications. The study highlights the importance of appropriate forecasts for policymakers, producers, and consumers to enhance better decision making and serve as an underlying incentive to guide the allocation of financial resources to the agricultural sector, which determines the structure and degree of sectoral growth.

Key words: grains, agriculture, forecasting, hybrid model, Nigeria.

Introduction and review of literature. Currently, price anticipating methods of agriculture sector can be segregated into two sorts i.e., qualitative and quantitative forecasting approaches. As an addition to the techniques of forecasting prices for agricultural products, the qualitative forecasting techniques generally do not dominate the usual position with small accuracy and big subjectivity. Depending on the distribution of time, quantitative forecasting techniques can be divided into

econometric technique, time series analysis technique, and intelligent forecasting technique. The econometric technique discovers the supportive economic theory to the research gap, then puts forward the hypothesis, and forms the econometric technique to affirm the hypothesis (Gogas et al., 2022; Khedr et al., 2021; Li & Leung, 2021; Sriboonchitta et al., 2013; Huang et al., 2012). Nevertheless, most empirical research does not verify that the anticipation effect of the classic econometric technique is better than that of the time series analysis technique (Martín-Rodríguez et al., 2012). Hence, the time series analysis technique was substituted with the econometric technique in the 1990s. Due to the perplexity and stalemate difficulty of price forecasting agricultural products, price volatility always reveals the features of repeated brunt, unpredictability, etc. The advantage of self-adaptation, self-learning, and self-organization possessed by the intellectual anticipation technique corresponds to the features of market price volatility of agricultural products. Therefore, in recent times, the intellectual anticipation technique has been progressively applied to the forecasting agricultural price. Intelligent prediction techniques usually comprise artificial neural network, chaos theory, entropy analysis, extreme learning machines, radial basis function, and support vector regression.

The agricultural sector contributes significantly to the Gross Domestic Product (GDP) in Nigeria, accounting for 20.85 %, 21.2 %, 21.91 % in 2017, 2018, and 2019, respectively (World bank, 2019). Agriculture is a common source of employment in sub-Saharan Africa, employing about 60 % population. Agriculture accounts for 90 % employment in the rural areas and a source of livelihood for 10–25 % of urban households (OECD-FAO, 2016). National census data in various countries around the world showed a gradual increase in the number of people employed in agriculture (Pattnaik et al., 2018; Huang & Yang, 2017; Lowder et al., 2016; Yeboah & Jayne, 2015). Development in the agricultural sector is determined by the prices of agricultural commodities, which indicates their scarcity or surplus. Besides, prices serve as incentives that drive the allocation of resources and fairly define the structure and extent of economic growth (Ferrara et al., 2022; McNerney et al., 2022; Akintunde et al., 2012). Developed nations and developing countries (e.g., Nigeria) benefit from agricultural commodity price forecasting because it helps in forecasting food security and alerting policymakers by detecting the warning signs of an impending crisis early in the crop marketing year (Wang et al., 2022; Xu & Hsu, 2021; Sabu & Kumar, 2020; Antonaci et al., 2014; Araujo et al., 2012). Hence, this study provides accurate forecasting techniques for specific agricultural commodities in sub-Saharan African countries. Price fluctuates all year round; therefore, understanding the trend of such fluctuations helps producers, consumers, and policy makers in better decision making. Generally, when price increases, a household will spend more of its disposable income on food compared with the income spent before the price increase (Van Wyk & Dlamini, 2018).

Recently, few studies have examined forecasting of agricultural commodity prices in Africa (Zhang et al., 2020; Tomek & Kaiser, 2014). This study addresses the

gap as it considers forecasting prices of widely consumed grains and legumes in Nigeria, such as imported rice, local rice, beans, and maize. These products are essential to overcome food deficit, increase income of households, reduce expenditures, and increase revenue for manufacturing industries. Nigeria is the most populated country in Africa, thereby making it the highest consumer of grains in tropical and Sub-Saharan Africa (USAID/MARKETS 2010). Commodities consumed in high proportion in Nigeria play a significant role in the prices of such commodities in Africa. The U.S. Department of Agriculture (USDA), Production, Supply and Distribution (PSD) production dataset for the years 2012/2013 and 2016/2017 produce marketing years indicate that the production of corn in Nigeria is between 7.0 and 7.8 million metric tons (PSD 2017). Maize is among the most significant crops and staple foods consumed in Nigeria. Nigeria is the 10th largest maize producer in the world and the major producer in Africa with a yearly production of more than six million metric tons (USAID/Markets, 2010). The demand for maize grain is increasing due to its enormous usage as raw material used by poultry farms, breweries, food, and beverage industries.

Beans are one of the largest produced legumes in Africa, with Niger and Nigeria producing more than 75 % of the total beans/cowpeas (Walker et al., 2014). Beans are an essential food legume and a ready source of protein for the masses, particularly in West Africa. Besides, it is a vital component of crop farming methods in sub-Saharan Africa because beans can be grown as a single crop, inter crop, or relay and combined with millet, sorghum, and maize (Boukar et al., 2011; Kamara et al., 2010; Manda et al., 2020). Beans are an important source of relatively low-cost protein that does not require cold storage, thus making it affordable for low-income households. It is popularly referred to as the 'poor man's meat' (Mishili et al., 2009). Nigeria produces the largest amount of beans in the world and is the largest importer and consumer in the Sub-Saharan Africa (Alene & Manyong, 2006; Mishili et al., 2009; Manda et al., 2020).

In Nigeria, rice is a prominent staple food among the agricultural commodities consumed. Rice cultivation is common mainly in the rice farms clustered in the northern to the middle belt in Nigeria. Rice consumption has increased by 10 % per year because of shifting consumer preferences (Akande, 2003). Ebuehi & Oyewole (2007) found that many Nigerians prefer imported rice brands rather than local rice varieties, mainly because local rice is not processed to meet international standards. The fluctuating prices of agricultural commodities have a substantial effect on the population's well-being and on the outputs and inputs of agricultural production. Therefore, this study underscores the specific and accurate methods for forecasting prices of commonly consumed grains or legumes in Nigeria. Accurate forecasting of grain prices will help agricultural policy makers, producers, and consumers to make informed decisions to ensure optimal profit, risk reduction, and build resistance against food insecurity, and farmers can decide the quantity to produce and the prices to set when cultivating crops (Dorosh & Haggblade, 2003; Vågsholm et al., 2020). Badmus & Ariyo (2011) showed that forecasting helps policy makers with regard to

production, price structure, and consumption of maize in Nigeria. Agricultural commodity prices can also help in predicting inflation. Tule et al. (2019) examined the predictability of agricultural commodity prices in Nigeria's inflation forecast. Their result indicated that agricultural commodity prices can individually predict food and headline inflation. Considering the use of various variables to estimate and forecast agricultural prices, appropriate models suitable for accurate forecast of individual commodities prices should be developed, because each commodity is likely to exhibit different characteristics or a trend during price determination. Hence, this study underscores forecasting of the most commonly consumed grain in Nigeria using the appropriate forecast models.

In forecasting agricultural commodity prices, studies have used various models with interesting results. Odior (2014) examined the effect of macroeconomic policy indicators on agricultural performance in Nigeria. The study employed a one-step dynamic forecast model to analyze this effect with annual time series data from 1970 to 2012. The findings showed that monetary aggregate, change in technology introduced over time, public spending on agriculture, rate of inflation, exchange rates, and nominal interest rates significantly influence the gross domestic product in Nigeria. Joshua et al. (2019) employed the Dicky-Fuller Test and simple exponential smoothing model to forecast beans prices in Adamawa state, Nigeria. The exponential smoothing model suggested an increase in beans prices. Agricultural production is risky in that it affects producers and consumers, hence the need for long-term strategies to mitigate these risks. Rashid & Jayne (2010) highlighted that farm income would increase by 30 % if effective risk management strategies are used. Appropriate statistical models should be used to understand these risks. Higgins et al. (2015) used the normalized difference vegetation index (NDVI) to identify and control for differences in productivity conditions for the prices of millet in three West African countries (Niger, Mali, and Burkina Faso). They found that NDVI information positively improves price forecast, which helps in the timely detection of food insecurity and with the planning and execution of a response. Nigeria is vulnerable to food insecurity because agriculture depends on rainfall and is a means of employment to most of its population. Zakari et al. (2012) forecasted production of two staple food grains (millet and sorghum) using the ARIMA model based on data for 1970–2010. They established that by 2030, the overall production of grains would be approximately 12678 thousand tons, with sorghum and millet production at nearly 1574.8 thousand tons and 4503 thousand tons, respectively. Such forecasting information will help policy makers to reduce the nation's vulnerability to food insecurity as regards price structure, production, and consumption.

The trend of food prices consumed locally and for export is an important budgetary tool for government agencies and food aid programs (Ackello-Ogutu, 2011; Schnepf, 2016; Sanusi, 2018; Kitenge & Morshed, 2019). Few among the existing studies analyze agricultural prices with specific techniques to forecasting commonly consumed grain prices in Africa. Chen et al. (2010) used the asset-pricing method to forecast world agricultural prices. They found that the indices of the

exchange rate and equity market of Australia, Canada, and New Zealand can forecast the changes of major food and agricultural commodity prices. Conversely, Taylor et al., (2006) forecasted crop prices for soybeans, corn, and sorghum in Kansas, United States, using historical averages augmented with the current market data. They found that this method improved the accuracy of forecasts based on post-harvest data. Zhang et al., (2020) used a model selection framework to forecast grain prices. Support vector regression (SVR), artificial neural network (ANN), and extreme learning machine (ELM) were used as prediction models. Their results suggested that less grain features are a feasible methodology to improve the model selection performance, and for grain produce, varying distributions of the time series characteristics are suitable for price forecasts. Forecast models for agricultural prices perform differently for each forecast period; therefore, the forecast period is essential for selecting the right forecast model. This issue remains underexplored in the literature on the model selection framework to forecast prices of agricultural commodities. To fill the gap in literature, this study suggests specific and suitable model selection framework comprising time series features for forecasting agricultural commodity prices for each grain considered. In this study we propose an appropriate forecasting model for each grain price over time with detailed insights to the procedure used in the selection.

This study is presented as follows: section 2 presents the purpose of the article, as well as materials and methods; the results and discussions are reported in section 3; the study is concluded in section 4.

The purpose of the article. This study highlights the specific and accurate methods for forecasting prices of commonly consumed grains or legumes in Nigeria based on data from January 2017 to June 2020.

Material and methods. Data on grain prices used in this study are obtained from the National Bureau of Statistics (NBS) in Nigeria; the data is available on request and also available on the bureau website. These prices were collected from the local governments across states, and it reflects the actual household prices. The average of commodity prices is collected every month and reported by the states, and then the country average is the combined average of all states. Data from respondents were gathered by more than 700 staff located in all states of the federation. These staff members are supported by supervisors, who are also monitored by internal and external observers. To maintain the standards of data collection, the NBS audit team conducts random selected verification of the collected prices. From these data, we selected the most widely consumed grains and legumes across the country: white maize, local rice, imported rice, and white beans. All grain prices consist of 42 periods from January 2017 to June 2020. In Figure 1, white beans and white maize show a relatively downward trend toward the end of 2019, after which both increase progressively. Local and imported rice show a similar pattern of a relatively stable trend for both until the third quarter and a gradual rise.




Figure 1. Time series plot of white beans, white maize, local rice, and imported rice

Source: authors' work.

Measures of forecasting accuracy. The best forecasting model was chosen based on three forecasting criteria: mean absolute error (MAE), root mean squared error (RMSE), and mean absolute percentage error (MAPE). Either MAE or RMSE is used to compare forecasting methods on a single data set, whereas MAPE is used for comparing the forecasting accuracy on data with varying time series with different measures. Thus, RMSE is an appropriate criterion if the data are free of extreme values, while MAE is superior in the presence of outliers (Pei & Li, 2019; Hyndman & Koehler, 2006). For example, based on MAE, the efficiency ratio of the suggested forecasting model relative to the benchmark model Ω , is defined as:

$$\Omega = \frac{MAE_p}{MAE_b}, \quad (1)$$

where MAE_b and MAE_p are from the benchmark and proposed models, respectively. A ratio less than 1 shows that the proposed forecasting model is more efficient than the benchmark model, and if Ω tends to 1, then the two forecasting models are nearly equivalent, or else, the proposed model works poorly (Safi & White, 2017).

1. ARIMA model. The typical $ARIMA(p,d,q)$ model is given by Box et al. (2015):

$$\phi(B)\nabla^d Y_i = \theta(B)\epsilon_i, \quad (2)$$

where, $d \geq 1$ is the degree of differencing; $\nabla = 1 - B$ is the differencing operator;

B, which is the *lag* operator, is defined as $BY_t = Y_{t-1}$, the operator that explains the prior value of the series. $\phi(B)$ And $\theta(B)$ are polynomials of degree p and q in B, respectively:

$$\phi(B) = 1 - \phi_1 B - \phi_2 B^2 - \dots - \phi_p B^p \quad (3)$$

And

$$\theta(B) = 1 - \theta_1 B - \theta_2 B^2 - \dots - \theta_q B^q \quad (4)$$

The best fit ARIMA model is selected according to AIC, AICc, or BIC value.

2. *STLM*. Seasonal and trend decomposition using loess method (STLM), developed by Cleveland et al. (1990), is a versatile and robust method for decomposing time series. Loess is a method for estimating nonlinear relationships. STLM can be robust to extreme values; consequently, these values will not affect the estimates of the trend cycle and seasonal components. In addition, STLM can handle any type of seasonality.

3. *Artificial neural network*. The *nnetar* function is used in fitting the ANNs. This function is described as feed-forward neural networks with one concealed layer and lagged inputs for forecasting univariate time series. This function fits Neural Network Autoregressive models NNAR (p, P, k). For the nonseasonal time series, the default is the optimum number of lags, which give the AIC, for a linear autoregressive (p) model (Hyndman, 2006).

4. *Hybrid model*. The hybrid model fits numerous individual model specifications to allow easy creation of collective forecasts. The hybrid model consists of a combination of three models: ARIMA, ANN, and STLM. Each component of the hybrid model captures its specific trend inbuilt in the model. For example, STLM is used for highly seasonal data, ARIMA is used for stationary data that is linear, ANN is used for nonlinear data.

Results and discussion. This section presents the empirical results of the models used for forecasting the price of legume and grains (white beans, white maize, local rice, and imported rice) using four approaches: the ARIMA, STLM, ANN models, and the hybrid combination of the three models.

In this study, the Anderson-Darling (AD) normality test was used to confirm if the residuals after approximation for the four models followed a normal distribution. The normality test yielded the following p -values for the following legumes and grains for the four models, Hybrid, ARIMA, STLM, and ANN, respectively: (1) white beans residuals: 0.5787, 0.3629, 0.06149, and 0.01706, (2) white maize residuals: 0.0549, 0.02835, 0.01366, and 0.0006516, (3) local rice residuals: 0.4596, 0.002722, 0.3162, and 0.09366, and (4) imported rice residuals: 0.5856, 0.01944, 0.5379, and 0.005839. The tests for normality for the residuals of the four data sets indicate that the normality assumption was not satisfied for all models. Given that the data are not normally distributed (without loss of generality), to compare the performance of the models through the four datasets, forecasting accuracy measure MAE was used over the forecasting period for each model. Smaller values of MAE indicate higher forecasting accuracy. Therefore, the ratios of the MAE of the hybrid

model to those of the ARIMA, STLM, and ANN models were analyzed.

Table 1 lists the complete empirical results for MAEs using the actual and predicted values of ARIMA, STLM, ANN, and Hybrid models. The forecasting model is chosen based on the forecasting criterion MAE. The efficiency ratio defined in equation (1) is used for this selection.

Table 1

MAEs of ARIMA, STLM, ANN, and Hybrid models

Dataset	Statistics	ARIMA	STLM	ANN	Hybrid
White Beans	RMSE	1127.85	1039.65	11020.39	3191.14
	MAE	30.35	27.67	100.26	52.60
	MAPE	11.93	10.93	39.04	20.57
White Maize	RMSE	376.98	880.48	518.45	542.09
	MAE	16.36	27.15	18.64	20.72
	MAPE	9.99	16.99	11.33	12.77
Local Rice	RMSE	1346.17	1904.68	5628.13	147.75
	MAE	35.15	41.98	73.27	9.15
	MAPE	10.28	12.30	21.77	2.76
Imported Rice	RMSE	5720.83	6124.17	9598.73	7039.68
	MAE	74.44	77.15	97.01	82.86
	MAPE	16.22	16.82	21.16	18.06

Source: authors' work.

The ANN model is applied to white beans with an average of 1,000 networks, which is a 2–25–1 network, with 101 weights and an estimated noise variance of 3.302. The result shows that the best fit model was the ARIMA (0,1,0) and an estimated noise variance of 127.8 (with AIC = 255.72, AICc = 255.84, and BIC = 257.21). For the STLM model, simple exponential smoothing with multiplicative errors is fitted. The estimated values of the model smoothing parameters are $\hat{\alpha} = 0.7796$, with the initial level $l_0 = 412.015$, and the estimated noise variance equal to 0.00072 (with AIC = 281.4043, AICc = 282.2043, and BIC = 285.9834). The corresponding MAEs of ARIMA, STLM, ANN, and Hybrid models equal 30.35, 27.67, 100.26, and 52.60, respectively. This result shows that the relative efficiencies of STLM model to the ANN, ARIMA, and Hybrid models equal $\Omega = 0.2759$, 0.9115, and 0.5260, respectively. Therefore, the STLM model is more efficient compared with ARIMA and is superior to the ANN and Hybrid models for white beans data. However, given the second choice in this case, the ARIMA model can be used because it is almost as efficient as the STLM model, although it is not a perfect substitute.

The ANN model is applied for white maize with an average of 1,000 networks, which is a 3–25–1 network, with 126 weights and an estimated noise variance of 0.00008. For the ARIMA model, the result shows that the best fit model was the ARIMA (0,1,1) and an estimated noise variance of 210.3 (with AIC = 273.52, AICc = 273.92, and BIC = 276.52). For the STLM, simple exponential smoothing with multiplicative errors is fitted. The estimated values of the model smoothing parameters are $\hat{\alpha} = 0.5636$, with the initial level $l_0 = 168.3865$, and the estimated noise

variance equals 0.00736 (with AIC = 306.3208, AICc = 307.1208, and BIC = 310.8999).

The corresponding MAEs equal 16.36, 27.15, 18.64, and 20.72 for ARIMA, STLM, ANN, and Hybrid models, respectively. This result shows that the relative efficiencies of the ARIMA model to the STLM, ANN, and Hybrid models equal $\Omega = 0.6026$, 0.8777, and 0.7897, respectively. Therefore, the ARIMA model is more efficient than ANN and is superior to the STLM and Hybrid models for white maize data. However, as a second choice for this grain, the ANN model could be considered. The ANN model is applied for local rice with an average of 1,000 networks, each of which is a 2–25–1 network, with 101 weights and an estimated noise variance of 3.85. For the ARIMA model, the result shows that the best fit model was the ARIMA (0,1,0) and estimated noise variance of 81.32 (with AIC = 240.8, AICc = 240.92, and BIC = 242.29). For the STLM, simple exponential smoothing with multiplicative errors is fitted. The estimated values of the model smoothing parameters are $\hat{\alpha} = 0.9999$, with the initial level $l_0 = 299.4986$, and the estimated noise variance equals 0.000807 (with AIC = 266.9505, AICc = 267.7505, and BIC = 271.5296).

The corresponding MAEs for these models equal 35.15, 41.98, 73.27, and 9.15 for ARIMA, STLM, ANN, and Hybrid models, respectively. This result indicates that the relative efficiencies of the Hybrid model to the STLM, ARIMA, and ANN models equal $\Omega = 0.2181$, 0.2605, and 0.1249, respectively. Therefore, the Hybrid model is superior to the ARIMA, STLM, and ANN models for local rice data. The ANN model is applied for imported rice with an average of 1,000 networks, each of which is a 2–25–1 network, with 101 weights and an estimated noise variance of 3.221. For the ARIMA model, the result shows that the best fit model was the ARIMA (0,1,0) and an estimated noise variance of 127.8 (with AIC = 255.72, AICc = 255.84, BIC = 257.21). For the STLM, simple exponential smoothing with multiplicative errors is fitted. The estimated values of the model smoothing parameters are $\hat{\alpha} = 0.7796$, with the initial level $l_0 = 412.015$, and the estimated noise variance equals 0.00072 (with AIC = 281.4043, AICc = 282.2043, and BIC = 285.9834). The MAEs equal 74.44, 77.15, 97.01, and 82.86 for ARIMA, STLM, ANN, and Hybrid models, respectively. This result indicates that the relative efficiencies of the ARIMA model to the STLM, ANN, and Hybrid models equal $\Omega = 0.9649$, 0.7674, and 0.8983, respectively. Therefore, the ARIMA model performs more efficiently than STLM, and is superior to the ANN and Hybrid models for imported rice data. However, as a second choice, the STLM model should be considered.

Figure 2 illustrates a comparison of the forecast for the grains using the best forecasting model with the actual values. It shows that the predicted values are close to the actual values; therefore, it substantiates the valid use of the suggested models. After feeding the model with data from November 2019 to June 2020 and repeating the procedure the forecasts for the four products for the following 8 months, that is,

July 2020 to February 2021 is shown in Table 2 and Figure 3.




Figure 2. Forecasts and actual values for grains

Source: authors' work.

In our forecast, using the best model for each grain, white beans and local rice show a declining trend in prices, whereas white maize and imported rice show a relatively stable trend for the forecasts. The prediction intervals are preserved from the individual component models and use the most extreme values from each individual model, producing conservative estimates for the performance of the combination of the models.

Table 2

Forecast for all grains using the full data

Date	White Beans (STLM)			White Maize (ARIMA)			Local Rice (Hybrid)			Imported Rice (ARIMA)		
	Point	Lo	Hi	Point	Lo	Hi	Point	Lo	Hi	Point	Lo	Hi
Jul-20	256.0	234.2	277.9	170.2	143.4	197.0	346.4	322.3	373.1	479.7	449.2	510.3
Aug-20	251.1	220.4	281.9	170.2	140.4	200.0	348.3	314.0	385.9	479.7	436.5	523.0
Sep-20	262.1	224.5	299.7	170.2	137.7	202.7	340.8	295.5	392.9	479.7	426.8	532.7
Oct-20	248.7	205.3	292.1	170.2	135.2	205.1	323.7	272.4	396.1	479.7	418.6	540.9
Nov-20	244.3	195.8	292.8	170.2	132.9	207.5	328.6	281.3	404.8	479.7	411.3	548.1
Dec-20	248.7	195.6	301.9	170.2	130.7	209.7	328.0	278.8	415.3	479.7	404.8	554.7
Jan-21	241.6	184.2	298.9	170.2	128.6	211.7	326.5	277.2	412.6	479.7	398.8	560.7
Feb-21	241.5	180.1	302.8	170.2	126.6	213.7	325.5	273.7	418.0	479.7	393.2	566.3

Source: authors' own calculations.

This study used different forecasting models to exploit the capabilities of the ARIMA, ANN, STLM, and the hybrid model that combines these three models in time series forecasting of grain prices in Nigeria. The forecasting performance was compared on the basis of the residuals for these models using data for some of the

most widely consumed grains in the region. The study further enumerated, explained, and discussed the various forecasting approaches and the criteria used for choosing a forecasting technique to provide the best result for each grain price. These results show that there is no universally suitable technique for all grains; rather, the forecast of each grain performs better with a specific model.




Figure 3. Forecasts for the four grains

Source: authors' work.

The study finds that ARIMA is the best applicable model for white maize and imported rice because it is well fitted to stationary data, as demonstrated in the sample period. Imported rice shows a relatively stable trend because specific quotas are allowed into the country every year, which provides some stability in its price, which also explains why the ARIMA model is the most suitable forecasting model. However, the STLM is more appropriate in forecasting white beans. As white beans are highly seasonal in Nigeria, it further explains why the STLM model fits better in forecasting prices. The production of local rice is inconsistent in Nigeria because of erratic rainfall and stiff competition from the importation of rice from other countries. Therefore, and consistent with the analysis, the hybrid model is the best model applicable to local rice because it captures varying trends exhibited in the data. For future research, we will answer questions such as; if the model decided for each grain is consistent for countries with similar characteristics for a wider range of commonly consumed grains in Africa. In addition, it will be important to test whether the importation and exportation of grains play a significant role in the forecasting methods that are chosen to forecast agricultural commodity prices. Consequently, we may apply the hybrid model for analyzing longitudinal data for African countries.

Conclusion. This study uses price data on widely consumed grains, such as white maize, local rice, imported rice, and white beans, in Nigeria from January 2017 to June 2020 to forecast grain prices. Different models that include autoregressive

integrated moving average, artificial neural networks, seasonal decomposition of time series by loess method, and a combination of these three models (hybrid model) is proposed to forecast the sample grain price data. This study contributes to the literature on accurate forecasting of agricultural commodity prices, and the analysis underscores the importance of providing the appropriate forecasts for policy makers, producers, and consumers for better decision making. Accurate agricultural price forecasts serve as a basic incentive to guide in the allocation of financial resources to the agricultural sector, which determines the structure and degree of sectorial growth. As prices fluctuate all year round, having information on future agricultural prices will improve planning and enhance food security.

The study finds that ARIMA is the best applicable model for white maize and imported rice because it is well fitted to stationary data, as demonstrated in the sample period. The STLM is more appropriate in forecasting white beans. As white beans are highly seasonal in Nigeria, it further explains why the STLM model fits better in forecasting prices. The production of local rice is inconsistent in Nigeria because of erratic rainfall and stiff competition from the importation of rice from other countries. Therefore, and consistent with the analysis, the hybrid model is the best model applicable to local rice because it captures varying trends exhibited in the data.

For future studies, a multivariate analysis can be performed to explore variables that explain the movement of grain prices. Also, a mixed data analysis approach could be used to see the effect of daily agricultural stock prices on future prices of grain commodities given different frequencies in available data.

Acknowledgments. We would like to express the deepest appreciation to the United Arab Emirates University library for proof reading, editing, comments and suggestions to improve the quality for this paper. Furthermore, we would like to add that there is no funding source for this research.

References

1. Ackello-Ogut, C. (2011). Managing food security implications of food price shocks in Africa. *Journal of African Economies*, 20(1), i100–i141. <https://doi.org/10.1093/jae/ejr010>.
2. Akande, T. (2003). The rice sector in Nigeria, United Nation Crop Project (UNCP) Country Agricultural Project on trade liberalization in agricultural sector and the environment. *African crop science journal*, 15(4), 223–235. Available at: <http://www.ifmaonline.org/pdf/congress/07Umeh&Ataborh.pdf>.
3. Akintunde, O. K., Akinremi, T. B., & Nwauwa, L. O. E. (2012). Food grain marketing in Osun State, Nigeria: a study of long-run price integration. *Continental journal of agricultural economics*, 6(1), 1–9. <https://doi.org/10.5707/cjae.2012.6.1.1.9>.
4. Alene, A., & Manyong, V. (2006). Endogenous technology adoption and household food security: the case of improved cowpea varieties in northern Nigeria. *Quarterly Journal of International Agriculture*, 45(3), 211–230. Available at: <https://hdl.handle.net/10568/91759>.

5. Antonaci, L., Demeke, M., & Vezzani, A. (2014). The challenges of managing agricultural price and production risks in sub-Saharan Africa. *ESA Working Paper No. 14-09*. Rome, FAO. <https://doi.org/10.22004/ag.econ.288979>.
6. Araujo, C., Araujo-Bonjean, C., & Brunelin, S. (2012). Alert at Maradi: preventing food crises by using price signals. *World Development*, 40(9), 1882–1894. <https://doi.org/10.1016/j.worlddev.2012.04.018>.
7. Badmus, M. A., & Ariyo, O. S. (2011). Forecasting cultivated areas and production of maize in Nigerian using Arima Model. *Asian Journal of Agricultural Sciences*, 3(3), 171–176. Available at: <https://maxwellsci.com/print/ajas/v3-171-176.pdf>.
8. Boukar, O., Massawe, F., Muranaka, S., Franco, J., Maziya-Dixon, B., Singh, B., & Fatokun, C. (2011). Evaluation of cowpea germplasm lines for protein & mineral concentrations in grains. *Plant Genetic Resources*, 9(4), 515–522. <https://doi.org/10.1017/S1479262111000815>.
9. Box, G. E., Jenkins, G. M., Reinsel, G. C., & Ljung, G. M. (2015). *Time series analysis: forecasting and control*. Hoboken, John Wiley & Sons.
10. Chen, Y., & Rogoff, K. S., & Rossi, B. (2010). Can exchange rates forecast commodity prices? *Quarterly Journal of Economics*, 125(3), 1145–1194. <https://doi.org/10.1162/qjec.2010.125.3.1145>.
11. Cleveland, R. B., Cleveland, W. S., McRae, J. E., & Terpenning, I. (1990). STL: seasonal-trend decomposition. *Journal of Official Statistics*, 6(1), 3–73. Available at: <https://www.wessa.net/download/stl.pdf>.
12. Dorosh, P., & Haggblade, S. (2003). Growth linkages, price effects and income distribution in Sub-Saharan Africa. *Journal of African Economies*, 12(2), 207–235. <https://doi.org/10.1093/jae/12.2.207>.
13. Ebuehi, O. A. T., & Oyewole, A. C. (2007). Effect of cooking and soaking on physical characteristics, nutrient composition and sensory evaluation of indigenous and foreign rice varieties in Nigeria. *African Journal of Biotechnology*, 6(8), 1016–1020. Available at: <https://scirp.org/reference/referencespapers.aspx?referenceid=2104917>.
14. Ferrara, L., Karadimitropoulou, A., & Triantafyllou, A. (2022). Commodity price uncertainty comovement: does it matter for global economic growth? Available at: <https://www.bundesbank.de/resource/blob/624356/68adedd2d53283e808f0a5563a218764/mL/2022-02-25-ferrara-data.pdf>.
15. Gogas, P., Papadimitriou, T., & Sofianos, E. (2022). Forecasting unemployment in the euro area with machine learning. *Journal of Forecasting*, 41(3), 551–566. <https://doi.org/10.1002/for.2824>.
16. Higgins, N., Hintermann, B., & Brown, M. E. (2015). A model of west african millet prices in rural markets. *Food Policy*, 52, 33–43. <https://doi.org/10.1016/j.foodpol.2014.09.011>.
17. Huang, W., Huang, Z., Matei, M., & Wang, T. (2012). Price volatility forecast for agricultural commodity futures: the role of high frequency data.

Romanian Journal of Economic Forecasting, 15(4), 83–103. Available at: https://ipe.ro/rjef/rjef4_12/rjef4_2012p83-103.pdf.

18. Huang, J., & Yang, G. (2017). Understanding recent challenges and new food policy in China. *Global Food Security*, 12, 119–126. <https://doi.org/10.1016/j.gfs.2016.10.002>.

19. Hyndman, R. J., & Koehler, A. B. (2006). Another look at measures of forecast accuracy. *International Journal of Forecasting*, 22(4), 679–688. <https://doi.org/10.1016/j.ijforecast.2006.03.001>.

20. Joshua, T., Zalkuwi, J., & Audu, M. M. (2019). Analysis of cost and return in cowpea production: a case study Mubi south local government area of Adamawa State, Nigeria. *Agricultural Science and Technology*, 11(2), 144–147. <https://doi.org/10.15547/ast.2019.02.023>.

21. Kamara, A. Y., Ellis-Jones, J., Ekeleme, F., Omoigui, L., Amaza, P., Chikoye, D. & Dugje, I. Y. (2010). A participatory evaluation of improved cowpea cultivars in the Guinea and Sudan savanna zones of North East Nigeria. *Archives of Agronomy and Soil Science*, 56(3), 355–370. <https://doi.org/10.1080/03650340903099692>.

22. Khedr, A. M., Arif, I., Raj, P., El-Bannany, M., Alhashmi, S. M., & Sreedharan, M. (2021). Cryptocurrency price prediction using traditional statistical and machine-learning techniques: a survey. *Intelligent Systems in Accounting, Finance and Management*, 28(1), 3–34. <https://doi.org/10.1002/isaf.1488>.

23. Kitenge, E. M., & Morshed, A. M. (2019). Price convergence among Indian cities: the role of linguistic differences, topography, and aggregation. *Journal of Asian Economics*, 61, 34–50. <https://doi.org/10.1016/j.asieco.2019.02.002>.

24. Li, R., & Leung, G. C. (2021). The relationship between energy prices, economic growth and renewable energy consumption: evidence from Europe. *Energy Reports*, 7, 1712–1719. <https://doi.org/10.1016/j.egyr.2021.03.030>.

25. Lowder, S. K., Skoet, J., & Raney, T. (2016). The number, size, and distribution of farms, smallholder farms, and family farms worldwide. *World Development*, 87, 16–29. <https://doi.org/10.1016/j.worlddev.2015.10.041>.

26. Manda, J., Alene, A. D., Tufa, A. H., Abdoulaye, T., Kamara, A. Y., Olufajo, O., ... & Manyong, V. M. (2020). Adoption and ex-post impacts of improved cowpea varieties on productivity and net returns in Nigeria. *Journal of Agricultural Economics*, 71(1), 165–183. <https://doi.org/10.1111/1477-9552.12331>.

27. Mishili, F. J., Fulton, J., Shehu, M., Kushwaha, S., Marfo, K., Jamal, M. ... & Lowenberg-DeBoer, J. (2009). Consumer preferences for quality characteristics along the cowpea value chain in Nigeria, Ghana, and Mali. *Agribusiness*, 25(1), 16–35. <https://doi.org/10.1002/agr.20184>.

28. Martín-Rodríguez, G., & Cáceres-Hernández, J. J. (2012). Forecasting pseudo-periodic seasonal patterns in agricultural prices. *Agricultural Economics*, 43(5), 531–544. <https://doi.org/10.1111/j.1574-0862.2012.00601.x>.

29. McNerney, J., Savoie, C., Caravelli, F., Carvalho, V. M., & Farmer, J. D. (2022). How production networks amplify economic growth. *Proceedings of the*

National Academy of Sciences, 119(1), e2106031118.
<https://doi.org/10.1073/pnas.2106031118>.

30. Odior, E. S. (2014). The macroeconomic policy effect on Nigerian agricultural performance: one-step dynamic forecasting analysis. *International Journal of Economics and Finance*, 6(9), 190–198.
<https://doi.org/10.5539/ijef.v6n9p190>.

31. OECD/Food and Agriculture Organization of the United Nations (2016). Agriculture in Sub-Saharan Africa: prospects and challenges for the next decade. OECD-FAO Agricultural Outlook 2016-2025, OECD Publishing, Paris.
https://doi.org/10.1787/agr_outlook-2016-5-en.

32. Pattnaik, I., Lahiri-Dutt, K., Lockie, S., & Pritchard, B. (2018). The feminization of agriculture or the feminization of agrarian distress? Tracking the trajectory of women in agriculture in India. *Journal of the Asia Pacific Economy*, 23(1), 138–155. <https://doi.org/10.1080/13547860.2017.1394569>.

33. Pei, S., & Li, Y. (2019). Wind turbine power curve modeling with a hybrid machine learning technique. *Applied Sciences*, 9(22), 4930.
<https://doi.org/10.3390/app9224930>.

34. Rashid, S., & Jayne, T. S. (2010). *Risks management in African agriculture: a review of experiences*. Paper presented at African agricultural markets programme's seminar – agricultural risks management in Africa: taking stock of what has and hasn't worked. Available at: <http://aec.msu.edu/fs2/aamp/index.htm>.

35. Sabu, K. M., & Kumar, T. M. (2020). Predictive analytics in agriculture: forecasting prices of arecanuts in Kerala. *Procedia Computer Science*, 171, 699–708.
<https://doi.org/10.1016/j.procs.2020.04.076>.

36. Safi, S. K., & White, A. K. (2017). Short and long-term forecasting using artificial neural networks for stock prices in Palestine: a comparative study. *Electronic Journal of Applied Statistical Analysis*, 10: 14–28.
<https://doi.org/10.1285/i20705948v10n1p14>.

37. Sanusi, O. I. (2018). The role of supply shocks and information dissemination on price dispersion in Nigeria. *Southern Illinois University at Carbondale*. Available at: <https://opensiuc.lib.siu.edu/dissertations/1599>.

38. Sriboonchitta, S., Nguyen, H. T., Wiboonpongse, A., & Liu, J. (2013). Modeling volatility and dependency of agricultural price and production indices of Thailand: static versus time-varying copulas. *International Journal of Approximate Reasoning*, 54(6), 793–808. <https://doi.org/10.1016/j.ijar.2013.01.004>.

39. Schnepf, R. (2016). US international food aid programs: background and issues, vol. 9. Washington DC, Congressional Research Service. Available at: <https://sgp.fas.org/crs/misc/R41072.pdf>.

40. Taylor, M. R., Dhuyvetter, K. C., & Kastens, T. L. (2006). forecasting crop basis using historical averages supplemented with current market information. *Journal of Agricultural and Resource Economics*, 31(3), 549–567.
<https://www.jstor.org/stable/40987335>.

41. Tule, M. K., Salisu, A. A., & Chiemeké, C. C. (2019). Can agricultural

commodity prices predict Nigeria's inflation? *Journal of Commodity Markets*, 16, 100087. <https://doi.org/10.1016/j.jcomm.2019.02.002>.

42. Tomek, W. G., & Kaiser, H. M. (2014). Agricultural product prices. In *Agricultural Product Prices*, 5th ed. Cornell University Press.

43. United States Agency for International Development (USAID) and Maximizing Agricultural Revenues and Key Enterprises in Targeted sites (MARKETS) (2010). *Package of Practices for Maize Production in Nigeria*. Available at: <https://data.nal.usda.gov/dataset/usda-foreign-agricultural-service-production-supply-and-distribution-database>.

44. USDA Foreign Agricultural Service (2018). *USDA foreign agricultural service production, supply, and distribution database*. Foreign Agricultural Service, Department of Agriculture. Available at: <https://data.nal.usda.gov/dataset/usda-foreign-agricultural-service-production-supply-and-distribution-database>.

45. Vågsholm, I., Arzoomand, N. S., & Boqvist, S. (2020). Food security, safety, and sustainability – getting the trade-offs right. *Frontiers in Sustainable Food Systems*, 4, 16. <https://doi.org/10.3389/fsufs.2020.00016>.

46. Van Wyk, R. B., & Dlamini, C. S. (2018). The impact of food prices on the welfare of households in South Africa. *South African Journal of Economic and Management Sciences*, 21(1), a1979. <https://doi.org/10.4102/sajems.v21i1.1979>.

47. Walker, T., Alene, A., Ndjeunga, J., Labarta, R., Yigezu, Y., Diagne, A., ... & Pandey, S. (2014). *Measuring the effectiveness of crop improvement research in sub-Saharan Africa from the perspectives of varietal output, adoption, and change: 20 crops, 30 countries, and 1150 cultivars in farmers' fields*. Synthesis Report for Objectives 1 and 2 of Bill & Melinda Gates Foundation's Diffusion and Impact of Improved Varieties in Africa (DIIVA) Project.

48. Wang, J., Wang, Z., Li, X., & Zhou, H. (2022). Artificial bee colony-based combination approach to forecasting agricultural commodity prices. *International Journal of Forecasting*, 38(1), 21–34. <https://doi.org/10.1016/j.ijforecast.2019.08.006>.

49. White, A. & Safi, S. K. (2016). The efficiency of artificial neural networks for forecasting in the presence of autocorrelated disturbances. *International Journal of Statistics and Probability*, 5(2), 51–58. <https://doi.org/10.5539/ijsp.v5n2p51>.

50. World Bank (2019), *World Development Indicators*. Available at: <http://data.worldbank.org/data-catalog/world-development-indicators>.

51. Xu, J. L., & Hsu, Y. L. (2022). The impact of news sentiment indicators on agricultural product prices. *Computational Economics*, 59(4), 1645–1657. <https://doi.org/10.1007/s10614-021-10189-4>.

52. Yeboah, K., & Jayne, T. S. (2015). *Employment Trends in Sub-Saharan Africa: The Evolving Role of Agriculture*. International Development Working Paper, Michigan State University, East Lan Gss-2009.

53. Zakari, S., Ying, L., & Province, H. (2012). Forecasting of Niger grain production & harvested area. *Asian Journal of Agricultural Sciences*, 4(4), 308–313. Available at: <https://maxwellsci.com/print/ajas/v4-308-313.pdf>.

54. Zhang, D., Chen, S., Liwen, L., & Xia, Q. (2020). Forecasting agricultural commodity prices using model selection framework with time series features and forecast horizons. *IEEE Access*, 8, 28197–28209. <https://doi.org/10.1109/ACCESS.2020.2971591>.

Citation:

Стиль – ДСТУ:

Sanusi O. I., Safi S. K., Adeeko O., Tabash M. I. Forecasting agricultural commodity price using different models: a case study of widely consumed grains in Nigeria. *Agricultural and Resource Economics*. 2022. Vol. 8. No. 2. Pp. 124–140. <https://doi.org/10.51599/are.2022.08.02.07>.

Style – APA:

Sanusi, O. I., Safi, S. K., Adeeko, O., & Tabash, M. I. (2022). Forecasting agricultural commodity price using different models: a case study of widely consumed grains in Nigeria. *Agricultural and Resource Economics*, 8(2), 124–140. <https://doi.org/10.51599/are.2022.08.02.07>.