



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

DECISION SUPPORT FOR FREEZE PROTECTION USING ARTIFICIAL NEURAL NETWORKS

Ronald W. McClendon, Professor
Gerrit Hoogenboom, Professor
Biological & Agricultural Engineering, University of Georgia

Introduction

In rainfed agricultural production systems, up to 90% of the variability in the yield can be attributed to weather differences. Fruit crops such as blueberries and peaches are particularly susceptible to low air temperatures. Temperatures near but above freezing might slow plant growth and development but such conditions do not typically cause significant damage if the plants are exposed for a short duration. However, once the temperature drops below freezing, the plants are easily damaged, with the severity of damage being determined by the duration of low temperature as well as the temperature itself.

As an example, during the Spring of 2002, a large area in blueberry and peach production in South Georgia was damaged due to an unusually severe and unexpected late frost. Irrigation is the most widely practiced frost protection measure for southeastern U.S. crops including peaches and blueberries. Irrigation results in the formation of a layer of ice that keeps the temperature of the flower near freezing, preventing it from dropping to below freezing temperatures. Farmers need information about when to start irrigation, as the process has to be started before the temperature drops to freezing. In addition to the expected low temperatures, farmers also need information about local wind speed, dew point or vapor pressure deficit to determine the point to initiate the freeze protection measures. Thus there is a need for accurate local weather information and short-term weather forecasts.

The goal of frost/freeze protection is to prevent plant parts (particularly flower and fruit) from being damaged by temperatures that drop below a critical level. This critical temperature is a function of crop variety and growth stage. There are several factors that can affect the degree of injury due to low temperature including the following: 1) type of plant, 2) stage of development of the crop, 3) the amount of leaf cover over the blossoms and fruit, 4) severity and duration of the freeze, and 5) wind speed. The primary prevention mechanism used by horticultural growers is irrigation. However, there are some problems associated with irrigating to prevent frost and freeze damage. If the humidity is low and the wind speed is high during a freeze, insufficient irrigation may cause evaporative cooling, which may damage crops more than non-irrigated fields. There are a number of recommendations that have been published by the extension service regarding frost/freeze damage:

- 1) If relative humidity is greater than 90%, there is little likelihood of worsening freeze damage with irrigation.
- 2) Open blueberry flowers cannot survive temperature less than 28°F. At the petal fall, the developing berry can be damaged by temperatures less than 30°F.

- 3) Overhead irrigation can be effective in preventing damage to blueberries if there is very little wind when the temperature drops below freezing.
- 4) If the humidity is low, irrigation should begin when temperature drops to 37°F.
- 5) If the temperature drops to the low 20s or below, or if wind is higher than 5 miles an hour, irrigation can cause increased damage.
- 6) If wind rises to 5 miles an hour, an application rate of 0.5 inches per hour will be needed to give some protection from frost or freeze damage.
- 7) At wind speeds above 5 miles an hour, more damage may be done by irrigating than by non-irrigating.

The dewpoint temperature is an important factor in decision making related to frost/freeze protection. Producers need an accurate estimate of temperature, wind speed, relative humidity or dew point temperature during critical times of the freeze. The dewpoint temperature is useful for estimating moisture content of air during freezes. With little wind, the dewpoint temperature changes only slightly between noon and the following sunrise, which makes it a good indicator of humidity. Relative humidity, however, may have large fluctuations during such periods.

Traditionally the role of providing weather forecasts has been the responsibility of the National Weather Service (NWS). However, due to changes in the laws, the NWS no longer provides data for agricultural applications. The NWS collects data from urban centers, thus the data are less useful for rural areas where farming is mostly done. In response to this need, the University of Georgia initiated the Georgia Automated Environmental Monitoring Network (AEMN) (Hoogenboom, 1996; 2000a; 2000b; Hoogenboom et al., 2000). This is a network of automated weather stations that are mainly located in the rural areas of Georgia (www.Georgiaweather.net). These weather stations collect data every second on the following variables: air temperature, relative humidity, soil temperature at 5 cm, 10 cm and 20 cm depths, wind speed, wind direction, solar radiation, vapor pressure deficit and soil moisture. The averages, or totals depending on the variable, are calculated every fifteen minutes and stored in the data logger. In addition, daily summaries are also calculated. The 15-minute and daily summary data are downloaded automatically to a central computer located in Griffin. The website disseminates this information as well as simple calculators that can dynamically provide degree days, chilling hours or water balance for management of irrigation (Georgiev and Hoogenboom, 1998; 1999; Hoogenboom et al., 1998).

An Artificial Neural Network (ANN) is a computational intelligence technique that mimics the behavior of neurons in the brain. The basic components of an ANN are its nodes or neurons and the connections between the nodes. A node is primarily a computational unit. It receives inputs, calculates a weighted sum and presents the sum to an activation function. The nodes are generally arranged in layers. In the back propagation ANN architecture, the input layer nodes receive the inputs and pass the results of their computations to the nodes in the hidden layer by means of the connections. The hidden nodes sum these weighted values as inputs, calculate an activation level and pass the results to the output nodes.

For an ANN to be useful it should be able to learn or capture the complex relationships between inputs and outputs. This is done by searching for an optimal set of the weights of the

connections between the nodes. It is achieved by first sending one set of inputs in the feed forward mode through the ANN. The error between the ANN output and the expected output is calculated. The error is then used to adjust the weights of the connections by using the method of gradient descent. The data that are used for this process are called the training data. The process is repeated until the error on another set of data called the testing data reaches a minimum. The testing data set is used in the feed forward mode only and are not used to adjust the weights. The training data and testing data comprise the model development data set. Once the training is complete, the ANN is then used with a separate model evaluation data set to determine its accuracy.

The overall goal of the research reported herein is to develop an ANN based Decision Support System (DSS) for predicting the occurrence of frost by predicting hourly temperatures during the subsequent twelve hour period. The most important inputs needed for the temperature predictions will be determined.

Methodology

Weather data for this study were obtained from the Georgia AEMN for the fruit producing areas of Georgia, including sites in peach producing areas (Fort Valley and Blairsville) as well as a blueberry producing area (Alma). The data from each location were divided into a model development set and model evaluation set. Data prior to 2001 were used for model development and data from years 2001 and 2002 were used for model evaluation. A final evaluation of the models was performed with data for 2003 since the 2001 and 2002 data, while not directly used for model development, were used to select the best configuration of hidden nodes. Only data from the months of January through April were used in the study. It is during this period that air temperatures vary between freezing and non-freezing and the crops are susceptible to freeze conditions.

The weather data consisted of observations of temperature, relative humidity, wind speed, solar radiation and rainfall. The current values of these variables were used as inputs along with corresponding prior values of these variables. The change in value of the weather variables from the prior values to current value (Δ) were calculated and also used as inputs.

To identify which weather variables are important, experiments were conducted with the hourly data from Fort Valley. The duration of prior data was held constant at four hours and the period of prediction was kept constant at four hours. The first set of experiments was performed to determine the best architecture when only temperature and its related inputs i.e. four hours of prior temperatures as well as their respective Δ values were used along with time of day. This set of inputs was called the core set of input variables. It was assumed that the current and prior air temperatures as well as time of day were important input variables. Subsequently, experiments were conducted for determining the best architecture when two weather variables are used as inputs. For this set of experiments the core set of inputs was used in conjunction with each of the other weather variables (relative humidity, solar radiation, rainfall and wind speed) one at a time to serve as the inputs. These experiments were then ranked based on the accuracy of their best architectures. The combination ranked first then became the new core set of inputs and

subsequent experiments were conducted by adding a third weather variable. This process was carried out until the most important input variables were determined in order of importance.

Results

A study was performed to determine the most important input variables. When using only temperature and its associated Δ values as the only inputs, it was found that the lowest MAE was 1.41°C when predicting four hours in the future. Each of the remaining weather variables and associated Δ values were included one at a time to determine the next most important input variables. Using this approach it was thus determined that the order of importance of the weather variable inputs considered was temperature, relative humidity, wind speed and solar radiation. The addition of rainfall slightly reduced the accuracy of the ANNs and it was thus excluded from all subsequent model development.

As expected, the accuracy of the ANNs decreased as the prediction period increased. Using the hourly format the MAE for predicting the temperature one hour in the future was 0.56°C and was 2.36°C for predicting temperature twelve hours in the future. A plot of predicted temperatures for a one hour period of prediction vs. the observed temperatures using the hourly data format for Fort Valley, GA, is shown. A linear regression line fit to this data gives an R^2 of 0.989. A similar plot of predicted temperatures for a twelve hour period is shown. As expected, the scatter is greater for the 12 hour period of prediction and the R^2 is 0.818.

The overall goal of this project was to develop ANN models for predicting temperatures which could be incorporated into a real time frost warning system. As such, all twelve ANNs predicting hourly temperatures for a given location were used to generate a simulated temperature forecast for the subsequent twelve hours. The outputs from the twelve networks that were developed were combined into a Decision Support System for frost prediction. Sample periods of data were selected for the simulation which included a freeze event. A Decision Point (DP) is defined as that point in time 't' when all networks are presented with the latest values of the input variables. For the simulation, the first decision point (DP1) was selected to be approximately ten hours prior to a freeze event. The subsequent decision points follow on an hourly basis.

Five day plots of predicted temperature and observed temperature versus time are shown using the eight hour prediction model. The data are for Ft. Valley, GA for February 24-March 1, 2002 and March 2-March 7, 2002. The model predicted the occurrences and durations of freezing conditions with reasonable accuracy. An animation of an actual twelve hour temperature prediction for Alma, GA in Bacon county is shown for the period January 5 through January 7, 2004. For this plot the prediction is updated every 45 minutes, although it could be performed every 15 minutes.

As an indication of the value of this frost/freeze decision support system, a quote from a UGA College of Agricultural and Environmental Sciences (CAES) county agent is included as an appendix. Robert T. Boland, Jr., county agent of Brantley County, Georgia, included this impact statement on the CAES website regarding the value of the AEMN weather network.

Acknowledgements

This work was funded in part by the USDA Federal Crop Insurance Corporation Risk Management Agency and by State and Federal Funds allocated to Georgia Agricultural Experiment Stations Hatch projects GEO00877 and GEO00895.

References

- Georgiev, G. and G. Hoogenboom. 1998. Crop growth and yield estimation using current weather forecasts and weather observations, *p 69-72*, In: *Preprints of the 23rd Conf. On Agricultural and Forest Meteorology*, American Meteorological Society, Boston, Massachusetts.
- Georgiev, G. A. and G. Hoogenboom. 1999. Near real-time agricultural simulations on the web, *Simulation 73(1): 22-28*.
- Hoogenboom, G. 1996. The Georgia Automated Environmental Monitoring Network, In: *Preprints of the 22nd Conf. On Agricultural and Forest Meteorology 343-346*, Boston, Mass., American Meteorological Society.
- Hoogenboom, G. 2000a. Contribution of agrometeorology to the simulation of crop production and its applications, *Agricultural and Forest Meteorology 103(1-2): 137-157*.
- Hoogenboom, G. 2000b. The Georgia Automated Environmental Monitoring Network, *p. 24-25*, In: *Preprints of the 24th Conference On Agricultural and Forest Meteorology*, American Meteorological Society, Boston, Massachusetts.
- Hoogenboom, G., G. Georgiev, D. Clarke, D.D. Gresham and G. Harbers. 1998. Internet Tools for delivery of weather data and applications, *p. 251-253*, In: *Preprints of the 23rd Conf. On Agricultural and Forest Meteorology*, American Meteorological Society, Boston, Massachusetts.
- Hoogenboom, G., G. A. Georgiev, and D.D. Gresham. 2000. Development of weather based products for agricultural and environmental applications, *p. 66-67*, In: *Preprints of the 24th Conf. On Agricultural and Forest Meteorology*, Boston, Mass., American Meteorological Society.

Appendix

Freeze Protection for Southern Highbush Blueberries

Robert T. Boland Jr.
Brantley County

Situation

Brantley County's growth in Southern Highbush Blueberries acreage is on the increase. Southern Highbush Blueberries are harvested around April 15th in Brantley County. This is a time when Florida is finishing their blueberry harvest, and the price for early Georgia Southern Highbush Blueberries start around \$36.00 a flat (\$10.00/lb).

However, producing the early maturing Southern Highbush Blueberry is not without risk. Late winter freezes can wipe out an entire crop. To reduce this risk of late winter freezes growers must use irrigation to protect the fruit and blooms. This method of irrigation involves a process of water melting and freezing, which keeps the fruit and blooms at a temperature around 32°. The key variable with freeze protection is knowing when to start irrigating. This requires knowing temperature, relative humidity, and dew points. Brantley County Blueberry growers were getting their weather information from the Weather Channel web site and from NOAA Weather Station in Jacksonville, Florida. However, this data is not generated in Brantley County. Due to the lack of good weather information ZBLU Farms experience a loss of about 80% of the crop, which includes the early Southern Highbush Blueberry.

Response

To address this problem for future Blueberry crops, the Brantley County Extension Coordinator contacted Dr. Gerrit Hoogenboom, Coordinator of the UGA's College of Agriculture and Environmental Sciences/Georgia Automated Environmental Monitoring Network about locating a weather station in Brantley County. This weather station would give growers in Brantley County local weather information. The request was granted and a weather station was located in Brantley County at ZBLU Farms on March 22, 2002. ZBLU Farms provide the local phone line allowing Dr. Hoogenboom to collect weather data and post it on the web.

Results

Brantley County Southern Highbush Blueberry growers have an early market window during which they can receive higher prices, around \$10.00 a pound. However, with early maturing varieties they face the risk of late freezes. Losses from the February 28, 2002 freeze were a whopping 80%, an estimated \$115,000.00 loss to the Brantley County Blueberry crop to the early maturing varieties. The Georgia Automated Environmental Monitoring Network is, and will continue to be a management tool which Brantley County Southern Highbush Blueberry growers are using to monitor weather condition in the county. This weather station will allow growers to use current and local weather data to guide them on when to start freeze protection for their blueberries.

Decision Support for Freeze Protection Using Artificial Neural Networks

R.W. McClendon, G. Hoogenboom
A. Jain, R. Ramyaa, B.A. Smith

Biological & Agricultural Engineering,
University of Georgia,
Athens and Griffin Campuses

Artificial Intelligence Center

Agricultural Outlook Forum 2005

Discussion

- 1) Frost/Freeze Problem
- 2) Recommendations
- 3) Weather Monitoring Systems

1) Frost / Freeze Problem

Goal

- Goal of Frost/Freeze Protection is to prevent plant parts (particularly flower and fruit) from being damaged by temperatures that drop below a critical level.
- Critical temperature is a function of crop, variety and growth stage.

Factors in degree of injury due to low temperature

- 1) Type of plant
- 2) Stage of development of the crop
- 3) Amount of leaf cover over the blossoms and fruit
- 4) Severity and duration of freeze
- 5) Wind speed

Problem associated with irrigation

If humidity is low and wind speed is high during freeze, insufficient irrigation may cause evaporative cooling which may damage crops more than non-irrigated fields.

2) Recommendations

Recommendations

- **If RH > 90%, little likelihood of worsening freeze damage with irrigation**
- **Open blueberry flowers cannot survive $T < 28$ °F. After petal fall, the developing berry can be damaged by $T < 30$ °F.**
- **Overhead irrigation can be very effective in preventing damage to blueberries if there is little wind when temp drops below freezing.**

Recommendations

- If humidity is low, start irrigation at 37 °F.
- If temp. drops to low 20's or below or if wind is higher than 5 mph, sprinkling can cause increased damage.

Recommendations

- If wind rises to 5 mph, an application rate of 0.5 in. / hr. will be needed to give some protection.
- At wind speeds above 5 mph, more damage may be done by irrigating than not irrigating

Dew point temperature (T_{DP})

- Producers need estimates of temperature, wind speed, relative humidity (or T_{DP}) during critical times of the freeze.
- T_{DP} is useful for estimating moisture content of air during freezes.
- With little wind, the T_{DP} changes little between noon and the following sunrise.

3) Weather Monitoring Systems


AEMN

The University of Georgia's
Automated
Environmental
Monitoring
Network




- Dr. Gerrit Hoogenboom
- 58 automated weather stations in rural areas of Georgia
- <http://www.griffin.peachnet.edu/bae/>

AEMN Stations




Equipment


- Campbell Scientific
 - CR10/CR10X Measurement and Control Module



Air Temperature & Relative Humidity



Rainfall



CR10 Power Supply




Solar Radiation



Wind Speed



Wind Direction



Soil Temperature



Communications

➤ Telephone

- Modem
- Voice modem

- Dedicated land lines
- Cellular phone/modem combination



Data Recording

- Collect data every second on:
 - Air temperature
 - Relative humidity
 - Soil temperature (4 depths)
 - Wind speed
 - Wind direction
 - Solar radiation
 - Vapor pressure deficit
 - Soil moisture

Data Recording

- Sensors are scanned at a *one-second* frequency
- Data are stored at 15-minute intervals
 - 15-Minute Average
 - 15-Minute Total
 - Maximum wind speed
 - Time of maximum wind speed
 - First rainfall occurrence

Automated Processing

Communications

- Call *all* stations, at least, every hour
- Call local stations at 15 minute intervals
- Loggernet (Campbell Scientific) software for data downloading

The screenshot shows the 'Status of Communication' window with a table of station data. The table has columns for 'Device Map', 'On-line', 'Error Rate', 'Last Collection Attempt', 'Collect', and 'Status'. The 'JONESB' station is highlighted in blue.

Device Map	On-line	Error Rate	Last Collection Attempt	Collect	Status
COM2		0%			
L Modem		16% (2)			
-ALPHARET		16% (2)	02/05/04 15:06:18	Area1 0/0	02/05/04 15:21:18
-ATLANTA		0%	02/05/04 15:04:46	Area1 18/18	02/05/04 15:15:30
-BLEDSOE		0%	02/05/04 15:02:46	Area1 23/23	02/05/04 15:15:30
-COVING		0%	02/05/04 15:03:20	Area1 24/24	02/05/04 15:15:30
-DALLAS		0%	02/05/04 15:03:54	Area1 24/24	02/05/04 15:15:30
-DEMPESEY		0%	02/05/04 15:04:19	Area1 24/24	02/05/04 15:15:30
-DULUTH		5%	02/05/04 15:01:19	Area1 26/26	02/05/04 15:15:30
-DUNWOODY		15% (2)	02/05/04 15:07:02	Area1 0/0	02/05/04 15:22:02
-GAINES		0%	02/05/04 15:05:13	Area1 22/22	02/05/04 15:15:30
-GRIFFIN		0%	02/05/04 15:00:47	Area1 30/30	02/05/04 15:15:25
-JONESB		0%	02/05/04 15:01:46	Area1 22/22	02/05/04 15:15:30
-WANSLEY		0%	02/05/04 15:02:20	Area1 22/22	02/05/04 15:15:30
L Temp		0%	10/07/03 08:22:54		
COM1	T	0%			
L Modem1	O	7%			
-CUMMINGS		9% (1)	02/05/04 15:01:30	Area1 0/0	02/05/04 16:00:25
-SNEADS		3%	02/05/04 15:02:36	Area1 744/744	02/05/04 16:00:25
-ALAPAHA		14% (1)	02/05/04 15:07:53	Area1 0/0	02/05/04 15:22:53
-ALMA		0%	02/05/04 14:13:12	Area1 69/69	02/05/04 15:00:30
-ARLINGT		3%	02/05/04 14:13:50	Area1 66/66	02/05/04 15:00:30
-ATTAPUL		0%	02/05/04 14:31:09	Area1 120/120	02/05/04 15:00:30
-BLAIRSVI		15% (2)	02/05/04 14:48:14	Area1 0/0	02/05/04 15:03:14
-BOWEN		6%	02/05/04 14:41:47	Area1 96/96	02/05/04 15:00:30
-BRUNSW		4%	02/05/04 15:09:38	Area1 104/104	02/05/04 16:00:30
-BYRON		0%	02/05/04 14:32:59	Area1 135/135	02/05/04 15:00:30
-CAIRO		3%	02/05/04 15:04:53	Area1 44/44	02/05/04 15:30:29
-CALHOUN		0%	02/05/04 14:33:45	Area1 125/125	02/05/04 15:00:30
-CALLAWAY		0%	02/05/04 15:10:31	Area1 96/96	02/05/04 16:00:30
-CAMILLA		0%	02/05/04 14:12:23	Area2 36/36	02/05/04 15:00:30
-CORDELE		0%	02/05/04 14:16:57	Area1 66/66	02/05/04 15:00:30
-DAHLON		5%	02/05/04 15:05:39	Area1 52/52	02/05/04 15:30:29
-DAWSON		1%	02/05/04 14:16:19	Area1 66/66	02/05/04 15:00:30
-DEARING		0%	02/05/04 14:20:09	Area2 99/99	02/05/04 15:00:30
-DIXIE		0%	02/05/04 14:23:28	Area1 120/120	02/05/04 15:00:30
-DUBLIN		3%	02/05/04 14:22:47	Area1 44/44	02/05/04 15:00:30
-EATONTON		0%	02/05/04 14:24:08	Area1 120/120	02/05/04 15:00:30
-ELBERTON		8%	02/05/04 14:34:39	Area1 192/192	02/05/04 15:00:30
-ELLIJAY		0%	02/05/04 14:24:48	Area1 120/120	02/05/04 15:00:30
-FLOYD		8%	02/05/04 14:44:44	Area1 78/78	02/05/04 15:00:30
-FTVALLEY		9%	02/05/04 14:51:39	Area1 220/220	02/05/04 15:00:30
-GEORGETO		1%	02/05/04 14:14:28	Area1 44/44	02/05/04 15:00:30
-HOMERV		0%	02/05/04 15:04:03	Area1 44/44	02/05/04 15:30:28

Collecting From: 57 nodes Toggle On/Off Collect Data Comm SWF Messages Low Level I/O
Disk Space: 1023.9 Mb View Messages View I/O
 Show Stations Only Reset Retries Stop Collect Log Status Msgs Log I/O To Disk



Georgia
Weather



Expand All | Collapse
All

☐ Stations (A-F)

☐ Stations
(G-W)

☐ Drought

☐ Fruits

☐ Cotton

☐ Peanut

☐ Weather Maps

☐ Daily

☐ Summaries

☐ Climate Maps

☐ Calculators

☐ Background

☐ Info

☐ General

☐ Others

☐ News

Supporters



Georgia Automated
Environmental
Monitoring Network
www.GeorgiaWeather.net



The Georgia Automated Environmental Monitoring Network was established in 1991 by the College of Agricultural and Environmental Sciences of the University of Georgia. The objective of the AEMN is to collect reliable weather information for agricultural and environmental applications. Each station monitors air temperature, relative humidity, rainfall, solar radiation, wind speed, wind direction, soil temperature at 2, 4, and 8 inch depths, atmospheric pressure, and soil moisture every 1 second. Data are summarized at 15 minute intervals and at midnight a daily summary is calculated. A microcomputer at the Georgia Experiment Station initiates telephone calls to each station periodically and downloads the recorded data. The data are processed immediately and disseminated via the world wide web (www.Georgiaweather.net). Click on a site below to obtain local weather and related information.

Warning! Please note that the web address <http://www.griffin.peachnet.edu/bae> will be discontinued. Please check your bookmark and change to www.Georgiaweather.net or www.griffin.uga.edu/aemn if needed.

24-hour graphical display for current conditions. See example for [Covington, Georgia](#) **L DATE!**


[Frances spawned highest winds and Jeanne, the most rain](#) **new**

[Hurricane highlights documented](#) **new** ...more news

Position: [Agrometeorologist/Agroclimatologist](#), [Hydroclimatologist/Hydrologist](#), [Crop Modeler](#)

Recent new stations: [Tiger](#), [Sasser](#), [Clarks Hill](#) **new**

Select a Site on the Map



Simple Applications & Calculators

- Heat Index/Wind Chill (humans & livestock)
- *WBGT – Wet Bulb Globe Temperature = Heat stress index* (humans & livestock)
- Degree Days
- Chilling Hours
- Water Balance Calculator
- Heating Degree Days
- Cooling Degree Days
- Rainfall Calculator
- Average Temperature Calculator
- Soil Temperature Calculator



Georgia
Weather



[AEMN Home](#)

Stations

- [Alapaha](#)
- [Albany](#)
- [Alma](#)
- [Alpharetta](#)
- [Arlington](#)
- [Atlanta](#)
- [Attapulgus](#)
- [Blairsville](#)
- [Brunswick](#)
- [Byron](#)
- [Cairo](#)
- [Calhoun](#)
- [Camilla](#)
- [Clarks Hill](#) NEW
- [Cordele](#)
- [Covington](#)
- [Dahlonega](#)
- [Dallas](#)
- [Dawson](#)
- [Dawson-HHERC](#)
- [Dearing](#)
- [Dixie](#)




**Georgia Automated
Environmental
Monitoring Network**
www.GeorgiaWeather.net





Georgia Experiment Station The University of Georgia Griffin, Spalding County, GA

Warning: Please note that the web address www.griffin.peachnet.edu/bae will be discontinued. Please check your bookmark and change to www.Georgiaweather.net or www.griffin.uga.edu/aemn if needed.

- | | |
|--|---|
| Current Conditions UPDATE! | Background Information NEW |
| Graph Current Conditions UPDATE! | Rainfall Calculator |
| Yesterday's Conditions | Average Temperature Calculator |
| 31-Day Summary | Soil Temperature Calculator |
| Historical Data | Degree Day Calculator |
| Climate Data | Chilling Hours Calculator |
| Today's local forecast | Water Balance Calculator |
| Temperature Prediction | Heating Degree Day Calculator |
| Minimum Temperature Estimator | Cooling Degree Day Calculator |
| Sunrise and Sunset Times and Moon Phases | Simulation Model Application |



All units will be in US system. Click the button to use units




FAWN



DATABASE : CLIMATE : TOOLS : FOCUS : NEWS : TOUR

- Latest Observations
- Graph Weather Data
- Archived Weather Data
- Management Tools



FAWN Focus
Our Sponsors

***** BELLE GLADE station will be offline for repairs until mid-January *****

Click for [location info...](#)
Toll-free in-state telephone access [\(866\)754-5732](#)

We site suggestions, questions? [Email FAWN's staff!](#)
Read the [Terms of use](#) under which this service is provided to you.
Copyright© 2004 UF/IFAS Last Modified: 01/04/2005 16:18:30





[DATABASE](#) [CLIMATE](#) [TOOLS](#) [FOCUS](#) [NEWS](#) [TOUR](#)

Latest Observations

Select Stations


----- Temperature (F) -----

Location	Datetime (Eastern Standard Time)	Temperature (F)					RH (pct)	Rain (in)	Wind		Solar Rad. (Wm2)
		2ft	Air 6ft	30ft	Dew* 30ft	Soil 4in			Dir. (deg)	Speed (mph)	
ALACHUA	TUE 1/4/2005 4:15 PM	75.5	74	74.2	55.8	63.1	53	0	171	4.27	233
APOPKA	TUE 1/4/2005 4:15 PM	76.4	75.2	75.3	56.9	69.9	53	0	75	5.82	117
AVALON	TUE 1/4/2005 4:15 PM	74.4	74.3	74.2	57.2	72.9	55	0	72	4.56	79
BALM	TUE 1/4/2005 4:15 PM	79	77.8	76.8	59.6	80.5	54	0	82	7.78	241
BELLE GLADE2	WED 11/17/2004 7:45 AM	63.1	62.4	62.3	62.1	68.9	99	0	0	0	0
BRADENTON	TUE 1/4/2005 4:15 PM	76.6	76.8	76.2	58.8	66.7	54	0	134	4.58	133
BRONSON	TUE 1/4/2005 4:15 PM	79.4	76.1	74.8	57	68.3	52	0	92	6.17	230
BROOKSVILLE	TUE 1/4/2005 4:15 PM	76.8	76	75.3	56	62.2	50	0	76	5.46	195
CARRABELLE	TUE 1/4/2005 4:15 PM	73.2	69.6	69.6	59.3	61.4	70	0	116	1.96	141
CITRA	TUE 1/4/2005 4:15 PM	73.4	73.6	72.9	59	69.8	60	0	21	3.04	66
DOVER	TUE 1/4/2005 4:15 PM	79	77.2	76.9	59.3	62.7	54	0	102	3.29	272
FORT LAUDERDALE	TUE 1/4/2005 4:15 PM	74.7	75	74.3	62.1	72	64	0	95	11.65	144
FT PIERCE	TUE 1/4/2005 4:15 PM	72.7	72.5	72.6	64.5	68.8	76	0	105	8.15	98
HASTINGS	TUE 1/4/2005 4:00 PM	71.7	71.1	71.1	61.2	65	71	0	84	6.76	131
HOMESTEAD	TUE 1/4/2005 4:15 PM	74.9	74.8	73.6	62.4	75.4	65	0	105	10.98	166

Items with an asterisk(*) are calculated values. They are not measured directly by FAWN.

Station: Parameter: Range: Units:
 Temp at 2 Ft Temp at 6 Ft Temp at 30 Ft Soil Temp Dew Point Wet Bulb Temp

Latest Readings
 Temp at 2Ft: 76.44 Temp at 6Ft: 75.18 Temp at 30Ft: 75.31 Soil Temp: 69.93 Dew Point: 56.91
 For the displayed period, the Maximum Temp at 2Ft is 80.37 and the Minimum is 76.08.




Copyright © 2003 UF/IFAS
 Read the [Terms of use](#) under which this service is provided to you.
 Web site suggestions, questions? [Email FAWN's staff!](#)
 Last Modified: 01/04/2005 16:23:20

[Click for Location Info...](#)
 Toll-free in-state telephone access
 (866) 754-5732

Approach

- Develop web-based, frost-warning Decision Support System (DSS) which utilizes current and prior weather data
- Use Artificial Neural Networks (ANNs) to develop the algorithms used in the DSS

Standard Three-Layer Back-Propagation Artificial Neural Network




Pattern	Input 1	Input 2	Input 3	...	Input n	Output
1	-	-	-	...	-	-
2	-	-	-	...	-	-
⋮						

Temperature	Rel. Humidity	Wind	Solar Radiation	Rainfall	MAE °C *
X					1.41
X	X				1.19
X		X			1.34
X			X		1.35
X				X	1.44
X	X	X			1.15
X	X		X		1.17
X	X			X	1.21
X	X	X	X		1.12
X	X	X		X	1.16
X	X	X	X	X	1.14

* Ft. Valley evaluation data set, Four-hour temperature prediction

ANN architecture for 12-hour prediction



Two General Categories of ANNs

- 1) Estimation of a numerical value
- 2) Classification

We have pursued both types of ANNs in our research project.

A. Jain, M.S. Artificial Intelligence, UGA

- Developed estimation ANNs to predict air temperature in hourly increments from one to twelve hours in the future based on current and prior weather data.

R. Ramyaa, M.S. Artificial Intelligence, UGA

- Developed classification ANNs to predict if frost occurs in a subsequent four, eight, or twelve-hour period.

B.A. Smith, M.S. Artificial Intelligence, UGA
(expected 2005)

- Developed software to allow the automatic training of large numbers of temperature-predicting ANNs using larger data sets.


Initial Study Locations

- Blairsville – North Georgia Mountains
- Fort Valley – Central Georgia
- Alma – South Georgia


Weather Data

- Model Development: 1993-2000
- Model Evaluation: 2001-2003

**Predicted and observed one-hour temperature predictions
for Fort Valley, 2001 and 2002 data.**




Predicted and observed twelve-hour temperature predictions for Fort Valley, 2001 and 2002 data.




A Predicted Freeze Event

March 4, 2002


1200 hrs to 2200hrs




Observed temperatures on
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs



Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs



Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs




Predictions made from the Decision Points (DP),
March 4, 2002, 1200hrs to 2200hrs


A Predicted Freeze Event

January 8, 2001


1200 hrs to 2400hrs




Observed temperatures on,
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs



Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs




Predictions made from the Decision Points (DP),
January 8, 2001, 1200hrs to 2400hrs

Fort Valley-8 hour prediction February 24-March 1, 2002



Fort Valley-8 hour prediction March 2-7, 2002



Implementation

During Spring 2004, the temperature prediction ANN was implemented on the website.



**Georgia Automated
Environmental
Monitoring Network**
www.GeorgiaWeather.net




Alma - Bacon County Airport Alma, Bacon County, GA

Temperature Prediction

Current Air Temperature: 8:30 PM, January 5 through 8:30 PM, January 6

Predicted Temperature: 8:30 PM, January 6 through 8:30 AM, January 7



Robert T. Boland Jr., Brantley County Agent

“Brantley County Southern Highbush Blueberry growers have an early market window during which they can receive higher prices, around \$10 per pound. However, with early maturing varieties they face the risk of late freezes. Losses from the Feb. 28, 2002 freeze were a whopping 80%, an estimated \$115,000 loss to the Brantley County Blueberry crop’s early maturing varieties. The Georgia [AEMN] is, and will continue to be, a management tool which Brantley County Southern Highbush Blueberry growers are using to monitor weather conditions... [It] will allow growers to use current and local weather data to guide them on when to start freeze protection for their blueberries.”

Acknowledgement

Risk Management Agency of the USDA
Federal Crop Insurance Corporation



Decision Support for Freeze Protection Using Artificial Neural Networks

**Questions and
Comments**