



**AgEcon** SEARCH  
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

# Estimating Nitrous Oxide (N<sub>2</sub>O) emissions from the application of N fertilizers to soils at higher resolution in Ireland

Carlos Alberto Francisco-Cruz<sup>1</sup>, Cathal Buckley<sup>2</sup>, James Breen<sup>1</sup>, Gary Lanigan<sup>3</sup>

<sup>1</sup> University College Dublin, School of Agricultural and Food Science, Dublin, Ireland.

<sup>2</sup> Teagasc Rural Economy and Development Programme, Athenry, Galway, Ireland.

<sup>3</sup> Teagasc Environmental Research Centre, Johnstown Castle, Wexford, Ireland.

Contact Information: [carlos.franciscocruz@ucdconnect.ie](mailto:carlos.franciscocruz@ucdconnect.ie)



*Extended abstract prepared for presentation at the 181st seminar of the EAAE  
“Greenhouse gas emissions in the EU agriculture and food sector: potential and limits  
of climate mitigation policies and pricing instruments”*

*Berlin, Germany*

*October 5-7, 2022*

*Copyright 2022 by C.A. Francisco-Cruz, C. Buckley, J. Breen, G. Lanigan. All rights reserved.  
Readers may make verbatim copies of this document for non-commercial purposes by any  
means, provided that this copyright notice appears on all such copies.*

## **Introduction**

This paper aims to define a high-resolution model to estimate nitrous oxide (N<sub>2</sub>O) emissions from agricultural soils in the Republic of Ireland. N<sub>2</sub>O emissions from the management of agricultural soils represented 10% of the total national GHG emissions in 2020 (Duffy et al., 2022). The current methodology for quantifying N<sub>2</sub>O emissions uses a general equation based on the Guidelines for National Greenhouse Gas Inventories (IPCC, 2006) and national emissions factors differentiated by fertilizer type (Tier 2). Although this method approximates the sector's emissions, it does not consider two crucial environmental factors, namely soil characteristics (pH, texture, drainage) and climate (precipitation and temperature). By taking into account these factors, N<sub>2</sub>O emissions can be quantified with increased granularity.

## **Background**

This research follows the methodology proposed by the Intergovernmental Panel on Climate Change (IPCC) for agricultural soils (category 3.D.1). N<sub>2</sub>O is primarily emitted from ecosystems as a by-product of nitrification and denitrification. Nitrification is the aerobic microbial oxidation of ammonium to nitrate, and denitrification is the anaerobic microbial reduction of nitrate to nitrogen gas. N<sub>2</sub>O is a gaseous intermediary in the reaction sequence (IPCC, 2006). The methodology measures direct and indirect N<sub>2</sub>O emissions, where the direct N<sub>2</sub>O emissions take into account the nitrogen inputs from synthetic fertilizer, animal manure, sewage sludge, crop residues, mineral soils, urine and dung deposited by grazing animals, and managed/drained organic soils, and the indirect N<sub>2</sub>O emissions includes emissions from atmospheric deposition and nitrogen leaching and run-off. N<sub>2</sub>O emissions are calculated by accounting for the amounts of each input and multiplying them by their respective emission factor (Duffy et al., 2022).

## **High-resolution model to quantify N<sub>2</sub>O emissions**

The high-resolution model modifies the emissions factor of the direct N<sub>2</sub>O emissions equations by adding soil characteristics and climate (environmental factors) by using indices. The soil characteristics indices are soil pH, texture, drainage, soil type and soil moisture, and they are defined considering the results obtained in previous research in Ireland and the EU (Hafner et al., 2019; Harty et al., 2016; Krol et al., 2016; Murphy et al., 2022; Rahman et al., 2021; Roche et al., 2016). Regarding climate, there are two indices, one for the annual precipitation and the other for air temperature. The precipitation index

is a linear function that affects emissions directly. The air temperature index is the difference between the highest and average temperature of the year. The assumptions of the climate indices are defined according to the information registered from 2010 to 2019 by the Irish Meteorological Service (Met, 2022).

In order to compare the high-resolution model, a baseline model was estimated to follow the current methodology in Ireland. This model was proposed in the 2020 Sustainability Report to show the GHG emissions at farm level in Ireland (Buckley, C. & Donnellan, T., 2021).

## **Data**

This paper draws on the Teagasc National Farm Survey (NFS) 2018, representing the agricultural sector by type of farm and region in Ireland. The NFS is part of the EU Farm Accountancy Data Network and provides highly detailed activity data for 900 farmers annually, and farms can be spatially referenced. In this survey, it is possible to identify the primary type of soil and its main characteristics for each farm according to the soil classification suggested by Gardiner and Radford (1980). Also, georeferencing allows overlaying the sample farms on climate data rasters at high-resolution (2.5 km horizontal grid) obtained from the Met Éireann (Gleeson et al., 2017).

## **Results**

Results indicate that the high-resolution model quantified 2.2% less N<sub>2</sub>O emissions compared to the baseline model. This reduction in the quantification is mainly due to a decrease in the estimated fertilizer emissions of 3.9% and a drop of 1.5% in urine and dung emissions when environmental factors are included. The differences between the baseline and high-resolution models are heterogeneous around the country due to the soil characteristics and climate conditions. The variance by farms ranges from -33% to 38%. Farms with a reduction in the quantification of N<sub>2</sub>O emissions have fewer quality soils, lower soil moisture and less variability in temperature and precipitation. In contrast, farms with an increase in the quantification of N<sub>2</sub>O emissions have a better quality of soils, high soil moisture and more variability in temperature and precipitation.

We confirm that the difference is statistically significant through a quantile regression of the difference between the baseline and high-resolution model. Furthermore, climate conditions have a statistically significant impact on N<sub>2</sub>O emissions in the high-resolution model. The results of the environmental factors show that the climate has a higher effect on N<sub>2</sub>O emissions than soil characteristics, and this impact is different throughout all the

distribution of N<sub>2</sub>O emissions at farm level. The total effect of the climate is 21%, while soil characteristics are 11%. The temperature and soil moisture variables are the ones with the most significant variability throughout the country, and they are the ones that best explain the differences in the high-resolution model.

## Conclusions

This paper's results contribute to improving the accuracy of N<sub>2</sub>O emission measurement on agricultural soils. In addition, it shows the importance of having disaggregated data on agricultural activity and climate. The environmental factors have a statistically significant effect on N<sub>2</sub>O emissions, mainly soil moisture and temperature. The evidence from this study suggests that climate conditions have a higher impact than soil characteristics on N<sub>2</sub>O emissions. This is important because the soil characteristics would be constant over time. In contrast, the climate conditions will show far greater variability.

The design of a high-resolution emissions estimation process allows analysis of different agricultural practices and could assist in targeting appropriate mitigation measures spatially. For instance, to define the cost of carbon at the farm level, it would be essential to consider the emission levels incorporating environmental factors by type of farm.

## Reference

- Buckley, C., & Donnellan, T. (2021). *Sustainability Report 2020*. Teagasc, Agricultural Economics and Farm Surveys Department.  
<https://www.teagasc.ie/publications/2021/national-farm-survey---2020-sustainability-report.php>
- Duffy, P., Black, K., Fahey, D., Hyde, B., Kehoe, A., Monaghan, S., Murphy, J., Ryan, A. M., & Ponzi, J. (2022). *Ireland National Inventory Report 2022* (p. 503).
- Gardiner, M. J., & Radford, T. (1980). *Ireland, general soil map* (2nd ed) [Map]. National Soil Survey.
- Gleeson, E., Whelan, E., & Hanley, J. (2017). Met Éireann high resolution reanalysis for Ireland. *Advances in Science and Research*, 14, 49–61.  
<https://doi.org/10.5194/asr-14-49-2017>
- Hafner, S. D., Pacholski, A., Bittman, S., Carozzi, M., Chantigny, M., Géniermont, S., Häni, C., Hansen, M. N., Huijsmans, J., Kupper, T., Misselbrook, T., Neftel, A., Nyord, T., & Sommer, S. G. (2019). A flexible semi-empirical model for estimating ammonia volatilization from field-applied slurry. *Atmospheric Environment*, 199, 474–484.  
<https://doi.org/10.1016/j.atmosenv.2018.11.034>

- Harty, M. A., Forrester, P. J., Watson, C. J., McGeough, K. L., Carolan, R., Elliot, C., Krol, D., Laughlin, R. J., Richards, K. G., & Lanigan, G. J. (2016). Reducing nitrous oxide emissions by changing N fertilizer use from calcium ammonium nitrate (CAN) to urea based formulations. *Science of The Total Environment*, 563–564, 576–586. <https://doi.org/10.1016/j.scitotenv.2016.04.120>
- IPCC. (2006). *Guidelines for National Greenhouse Gas Inventories*.
- Krol, D. J., Carolan, R., Minet, E., McGeough, K. L., Watson, C. J., Forrester, P. J., Lanigan, G. J., & Richards, K. G. (2016). Improving and disaggregating N<sub>2</sub>O emission factors for ruminant excreta on temperate pasture soils. *Science of The Total Environment*, 568, 327–338. <https://doi.org/10.1016/j.scitotenv.2016.06.016>
- Met. (2022). *The Irish Meteorological Service*. The Irish Meteorological Service. <https://www.met.ie/climate/available-data/mera>
- Murphy, R. M., Saunders, M., Richards, K. G., Krol, D. J., Gebremichael, A. W., Rambaud, J., Cowan, N., & Lanigan, G. J. (2022). Nitrous oxide emission factors from an intensively grazed temperate grassland: A comparison of cumulative emissions determined by eddy covariance and static chamber methods. *Agriculture, Ecosystems & Environment*, 324, 107725. <https://doi.org/10.1016/j.agee.2021.107725>
- Rahman, N., Richards, K. G., Harty, M. A., Watson, C. J., Carolan, R., Krol, D., Lanigan, G. J., & Forrester, P. J. (2021). Differing effects of increasing calcium ammonium nitrate, urea and urea + NBPT fertilizer rates on nitrous oxide emission factors at six temperate grassland sites in Ireland. *Agriculture, Ecosystems & Environment*, 313, 107382. <https://doi.org/10.1016/j.agee.2021.107382>
- Roche, L., Forrester, P. J., Lanigan, G. J., Richards, K. G., Shaw, L. J., & Wall, D. P. (2016). Impact of fertilizer nitrogen formulation, and N stabilizers on nitrous oxide emissions in spring barley. *Agriculture, Ecosystems & Environment*, 233, 229–237. <https://doi.org/10.1016/j.agee.2016.08.031>