



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

## **BLACK TURMERIC- *CURCUMA CAESIA* ROXB, AND ITS MEDICINAL PROPERTIES: A REVIEW**

Sapan Tamang

PhD Research scholar, Department of Plantation Spices,  
Medicinal and Aromatic crops, Faculty of Horticulture, Bidhan Chandra  
Krishi Viswavidyalaya, Mohanpur- 741252, Nadia, West Bengal

**DOI: <https://doi.org/10.51193/IJAER.2022.8111>**

Received: 05 Feb. 2022 / Accepted: 12 Feb. 2022 / Published: 28 Feb. 2022

### **ABSTRACT**

*Curcuma caesia* Roxb., commonly known as black turmeric is an important medicinal plant belonging to the family Zingiberaceae. It is a perennial crop with bluish-black rhizome. The rhizome in inner part is bluish-black in colour and emits a characteristic sweet smell due to the presence of essential oil. It contains: alkaloids, terpenes, amino acids, carbohydrates, tannins, flavones, flavonoids, steroids, reducing sugars, proteins, anthraquinones, glycosides, cardiac glycosides. The rhizomes have a high value because of its putative medicinal properties. The rhizomes are used in carminative properties (Gantait *et al.*, 2011), treatment of smooth muscle relaxant activity, haemorrhoids, leprosy, asthma, cancer, epilepsy, fever, wound, vomiting, menstrual disorder, anthelmintic, aphrodisiac, gonorrhoeal discharges and inflammation (Arulmozhi *et al.*, 2006 and Sasikumar, 2005).

**Keywords:** alkaloids, *Curcuma caesia*, essential oil, economical importance, flavonoids

### **INTRODUCTION**

*Curcuma caesia* Roxb., commonly known as black turmeric is an important medicinal plant belonging to the family Zingiberaceae. *C. caesia* or Black Turmeric is a perennial crop with bluish-black rhizome. The rhizome is bluish-black in colour in inner part and emits a characteristic pungent smell due to the presence of essential oil (Pandey *et al.*, 2003; Das *et al.*, 2013). It is native to North East India and central India, relatively unexplored medicinal plant valued all over Asia for its therapeutic efficacy. It flourishes good at moist deciduous forest areas with rich humid and clayey soils. In India, it thrives well states like-Chhattisgarh, Madhya Pradesh, Odisha, Uttar Pradesh and West Bengal (Nadkarni, 1976). The rhizomes have a high

value because of its putative medicinal properties. The rhizomes are used in carminative properties (Gantait *et al.*, 2011), treatment of smooth muscle relaxant activity, haemorrhoids, leprosy, asthma, cancer, epilepsy, fever, wound, vomiting, menstrual disorder, anthelmintic, aphrodisiac, gonorrhoeal discharges and inflammation (Arulmozhi *et al.*, 2006 and Sasikumar, 2005). The rhizomes are aromatic and contain an essential oil. A paste made from the crop is used to cure blood dysentery and as poultice in rheumatic pain. The leaves both fresh and dry are used as a wrapping material and fuel respectively by Bengalis in Bangladesh (Yusuf *et al.*, 2002). Northern tribes use Black Turmeric as a talisman to keep the evil spirits away, while in West Bengal it is an important place in traditional system of medicine and is also used as a substitute for turmeric in state of Madhya Pradesh also the plant is regarded as very auspicious and it is stated that a person who possesses it will never encounter the shortage of cereals and food (Pandey and Chowdhury, 2003). This clear that this crop is not only important medicinally but also important socially as well as spiritually. The National Medicinal Plant Board of India has listed more than 112 plants as crucially important species circulated by Ministry of Environment in year 1997 and according to this, export of these species is illegal without the permission of the legal competent authorities and this notification includes *Curcuma caesia* (Black Turmeric) as well. These plants is in great demand in Central India and due to indiscriminate exploitation the plant has been categorized as an endangered species (Kumar *et al.*, 1998). While turmeric species have been growing all over the tropics, some species have also been reported from Australia, China, and South Pacific. More than 110 species of turmeric have been reported from the tropical Asia with greatest diversity from India, Myanmar and Thailand. About 40 species are indigenous to India, and black turmeric *Curcuma caesia* is one of the economically important species along with *C. aromatica*, *C. amada*, *C. aeruginosa*, *C. longa*, and *C. zanthorrhiza* (Ravindran 2007; Nair 2013).

## **BOTANICAL DESCRIPTION**

Kingdom: Plantae

Subkingdom: Viridaplantae

Phylum: Tracheophyta Sinnott

Subphylum: Euphyllophytina

Class: Magnoliopsida “monocotyledons” “commelinids”

Order: Zingiberales

Family: Zingiberaceae

Subfamily: Zingiberoideae

Tribe: Hedychieae

Genus: *Curcuma*

Species: *C. caesia* Roxb

Vernacular Names in different parts of India *C. caesia* is known by different names

Hindi: Kali Haldi, Nar Kachura Krishna Kedar

Bengali: Kala Haldi

Mizo: Aihang, Ailaihang

Telugu: Nalla Pasupu

Assamese: Kala Haladhi

Nepalese: Kaal Hardi



**Fig 1: Black Turmeric**

The plant height ranging from 0.5 to 1.0 m with normally erects. It is divided into underground large ovoid tuberous rhizome, often called rootstock, and aerial shoot along with the leaves. The leaves are usually in the groups of 10–20, each leaf is broad oblong lanceolate and glabrous. A deep farraginous purple colour is present in the middle region of the lamina the petiole is ivory in colour. The variation is parallel in nature. The flowering bracts are green with a ferruginous tinge. Flower petals may be deep pink or red in colour.



**Fig 2: Flower of Black Turmeric**

## CONSTITUENTS

It contains: amino acids, anthraquinones, alkaloids, carbohydrates, flavones, flavonoids, terpenes, tannins, steroids, reducing sugars, proteins, glycosides, cardiac glycosides.

The volatile rhizomes oil of *Curcuma caesia* contains of 30 components, representing 97.48% of the oil, with camphor (28.3%), ar-turmerone (12.3%), (Z) ocimene (8.2%), 1,8 cineole (5.3%), elemene (4.8%), borneol (4.4%), bornylacetate (3.3%) and curcumen (2.82%), ar- curcumene (6.8%) as the major constituents.





**Fig 3: Black Turmeric rhizomes**

## MEDICINAL USES

1. The multiple phyto-constituents like curcuminoids, flavonoids, phenolics, amino acids, protein and high alkaloids, found in the rhizome are responsible for the antimicrobial, antioxidant activities, anti-tumor, anxiolytic, anti-inflammatory, anti-ulcer, and CNS depressant (Devi *et al.*, 2015; Karmakar *et al.*, 2013; Vineela *et al.*, 2017).

### 2. Bronchodilating activity

The bronchodilating activity of extracts of Black turmeric . Bronchodilator activity of the extract was studied on the histamine aerosol induced Bronchospasm and preconvulsion dyspnoea in guinea pigs was reported by Pritesh Paliwal *et al.* (2011)

### 3. Anthelmintic Activity

Gill Randeep *et al.* (2011) studies two popular species of genus *Curcuma*, *C. amada* and *C. caesia* were proved for their anthelmintic activity. In this study, 4 extracts viz. aqueous extract of rhizomes, Dichloromethane, ethanol and Petroleum ether of *Curcuma amada* and *Curcuma caesia* were investigated for anthelmintic activity at three different concentrations. 50 mg/ml, 100 mg/ml and 150 mg/ml, three concentrations of each extract were studied the determination of paralysis time and time of death of earthworms The results proved that ethanol extract (150 mg/ml) of Black turmeric was most effective in causing paralysis of earthworms, while the

Dichloromethane extract (150 mg/ml) and ethanol extract (150 mg/ml) of both *C. caesia* and *C. amada* were very effective in causing death of earthworms.

#### 4. Anti-bacterial Activity

Angel Gabriel Rajamma *et al.* (2012) investigated antibacterial and antioxidant of oleoresins isolated from nine *Curcuma* species. Oleoresins were extracted from rhizomes of nine *Curcuma* species such as *C. aeruginosa*, *C. amada*, *C. aromatica*, *C. brog*, *Curcuma caesia*, *Curcuma malabarica*, *C. rakthakanta*, *C. sylvatica* and *Curcuma zedoaria*

#### 5. Anti-ulcerogenic activity

The anti-ulcer activity of the ethanolic extract of the rhizome of *Curcuma caesia* was experimented on four groups of albino rats and revealed that there is significant reduction of gastric acid volume, pepsin, ulcer index, free and total acidity along with increased production of gastric mucus. Pranab KR Bordoloi *et al.* (2012)

6. Smooth Muscle Relaxant and Anti-asthmatic Activity: Arulmozhi *et al.* (2006) evaluated anti-asthmatic property of *Curcuma caesia*.

#### 7. Anti-fungal Activity

Banerjee and Nigam, 1976 reported that the antifungal activity in *C. caesia* rhizomes. Essential oil of rhizomes of *Curcuma caesia* has been known for its antifungal activity.

8. Black turmeric, used in traditional and folk medicine, seems to be a promising source of active therapeutic agents. Further studies on isolation of active principle agents may play vital role in increasing its pharmaceutical and industrial significance. *Curcuma*, known for its antimicrobial potential since immortal, it's an important source of plant based antimicrobials. Such species provide safer alternates to the microbe-based antimicrobials which are increasingly reported for their side effects and drug resistance (Pandey *et al.*, 2015)

### REFERENCES

- [1] Angel, G.R, Vimala, B. 2012. Antioxidant and anti-inflammatory activities of proteins isolated from eight *Curcuma* species. *Journal of Phytopharmacology*. 4(1):96-105.
- [2] Arulmozhi, D.K, Sridhar N. 2006. Preliminary mechanistic studies on the smooth muscle relaxant effect of hydroalcoholic extract of *Curcuma caesia*. *Journal of Herb Pharmacotherapy*.6:117-24

- [3] Arulmozhi, D.K., Sridhar N. Veeranjanyulu, A. and Arora, K.S. 2006. Preliminary mechanistic studies on the smooth muscle relaxant effect of hydroalcoholic extract of *Curcuma caesia*. *Journal of Herbal Pharmacotherapy*.6: 3 – 4
- [4] Banerjee, A. Nigam, S.S.1976. Antifungal activity of the essential oil of *Curcuma caesia* Roxb. *Indian Journal of Medicinal Research*. 64:1318-21.
- [5] Das, S, Bordoloi, P.K, Phukan, D, Singh S. 2012. Study of the anti-ulcerogenic activity of the ethanolic extracts of rhizome of *Curcuma caesia* against gastric ulcers in experimental animals. *Asian Journal of Pharmacy Clinical Research*. 5:200-3
- [6] Das, S., Mondal, P., Zaman., Md. K .2013. *Curcuma caesia* Roxb. and its medicinal uses: a review. *International Journal of Research in Pharmacy and Chemistry*, 3(2): 370-375.
- [7] Devi, H.P, Mazumder, P.B, Devi, L.P .2015.Antioxidant and antimutagenic activity of *Curcuma caesia* Roxb. rhizome extracts. *Toxicology Reports*: 2: 423–428.
- [8] Gantait, A, Barman, T, Mukherjee, P.2011. Validated method for determination of curcumin in turmeric powder. *Indian Journal of Traditional Knowledge*.10:247-250.
- [9] Gill, R, Kalsi, V.2013. Photochemical investigation and evaluation of anthelmintic activity of *Curcuma amada* and *Curcuma caesia*-a comparative study of Ethnopharmacology; 2(1):412-11.
- [10] Karmakar, I, Dolai, N Suresh R.B.K, Kar, B, Roy S.N, Halder, P.K.2013. Antitumor activity and antioxidant property of *Curcuma caesia* against Ehrlich's ascites carcinoma bearing mice. *Pharmaceutical Biology*: 51(6): 753–759.
- [11] Kumar, S. Singh, J, Shah. N.C, Ranjan V.1998. Indian Medicinal Plants Facing Genetic Erosion. CIMA, Lucknow.
- [12] Nadkarni, K.M.1976. Indian Material Medical. *Popular Prakashan*, Bombay.
- [13] Nair, K.P.P .2013. The agronomy and economy of turmeric and ginger, the invaluable medicinal spice crops. *Elsevier Scientific Publishing Company*.
- [14] Pandey, A. Agnihotri, V.2015. Antimicrobials from medicinal plants: research initiatives, challenges and the future prospects. In: biotechnology of bioactive compounds: sources and applications in food and pharmaceuticals: (VK Gupta, M G Tuohy, A O'Donovan, M Lohani, Eds.), John Wiley & Sons, Ltd., 123-150.
- [15] Pandey, A.K, Chowdhury, A.R.2003. Volatile constituents of the rhizome oil of *Curcuma caesia* Roxb. from central India. *Flavour and Fragrance Journal*: 18: 463-465.
- [16] Pritesh, Paliwal, S S Pancholi.2011. Pharmacognostic parameters for evaluation of the rhizomes of *Curcuma caesia*. *Journal of advanced pharmaceutical technology and research*:2(1):56-6



- [17] Ravindran, P.N. 2007. Turmeric the golden spice of life. In: turmeric the genus *Curcuma*: medicinal and aromatic plants- industrial profiles (PN Ravindran, K. Nirmal Babu, K. Sivaraman, Eds.), *CRC Press*, 45: 1-13.
- [18] Sasikumar, B. 2005. Genetic resource of *Curcuma*: diversity, characterization and utilization. *Plant Genet Resource*: 3:230-251
- [19] Vineela, C, Soundarya B. 2017. Study of antihelmintic activity of *Curcuma caesia*: 7(7): 248-252.
- [20] Yusuf, M., Rahma, M. A., Chowdhury, J. U. and Begum J.2002. Indigenous knowledge about the use of Zingibers in Bangladesh. *Journal of Economic Taxonomy Botany*: 26: 566–570.