



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*



Ag 84 Te
Reserve
United States
Department of
Agriculture

CAT/8a

Agricultural
Research
Service

Technical
Bulletin
Number 1761

Evaluation of the USDA Wild Soybean Germplasm Collection: Maturity Groups 000 to IV (PI 65.549 to PI 483. 464)

ABSTRACT

Juvik, Gail A., Richard L. Bernard, Ruzhen Chang, and James F. Cavins. 1989. Evaluation of the USDA Wild Soybean Germplasm Collection: Maturity Groups 000 to IV (PI 65.549 to PI 483.464). U.S. Department of Agriculture, Technical Bulletin No. 1761, 25 p.

This report contains data on the origin, descriptive characteristics, agronomic performance, and seed composition for 200 wild soybean (Glycine soja Siebold & Zucc.) accessions in maturity groups 000 to IV from the USDA Wild Soybean Germplasm Collection. These accessions (PI 65.549 to PI 483.464) were introduced from China, Japan, South Korea, and the Soviet Union into the United States from 1925 to 1984. A total of 38 categories of data is presented for each entry. These accessions were evaluated at Urbana, IL, in 1984 and 1985.

KEYWORDS: Evaluation, germplasm, Glycine soja Siebold & Zucc., origin, seed composition, wild soybean.

Trade names are used in this publication solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture or an endorsement by the Department over other products not mentioned.

Copies of this publication may be purchased from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.

ARS has no additional copies for free distribution.

CONTENTS

Introduction	1
Maturity groups 000 to IV	6
Table 1. Identification and origin information for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464	
Table 2. Descriptive data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464	
Table 3. Agronomic data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464	
Table 4. Seed composition data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464	

EVALUATION OF THE USDA WILD SOYBEAN GERMPLASM COLLECTION:
MATURITY GROUPS 000 TO IV (PI 65.549 TO PI 483.464)

Gail A. Juvik, Richard L. Bernard, Ruzhen Chang, and
James F. Cavins

This publication contains information on the origin, descriptive characteristics, agronomic performance, and seed composition of wild soybean (*Glycine soja* Siebold & Zucc.) germplasm accessions for PI 65.549 to PI 483.464 in maturity groups 000 through IV. These data can also be obtained through the Germplasm Resources Information Network (GRIN), Database Management Unit, USDA-ARS, BARC-West, Beltsville, MD 20705.

The accessions were tested on the Agronomy-Plant Pathology South Farm, University of Illinois, Urbana (40°8' N. lat.), with one replication per year. Seeds were planted on May 14 and 15, 1984, and on May 20, 1985. Plots consisted of one row with four hills per row, 60 cm between hills, and 1 m between rows. In 1984, plots for maturity groups III and IV were two rows wide, two hills per row, 1 m between hills, and 1 m between rows. Data were collected on all plants in a plot. The accessions were blocked by maturity group in the field, but the data are reported numerically by PI number.

The growth of wild soybeans in central Illinois is highly variable. Seeds of wild soybeans, which are normally very impermeable, were scarified in concentrated sulfuric acid for 5 minutes, rinsed thoroughly in water, and air-dried before planting. Despite this treatment, hard seed coats still resulted in some delayed emergence and poor stands. The number of plants per plot ranged from 2 to 20. Most wild soybeans are very susceptible to soybean mosaic virus and leafhopper feeding, and these pests caused considerable yellowing, stunting, and leaf rugosity. The insecticide Pydrin was applied in July both years to help reduce insect feeding.

Some accessions in the collection appear to be of hybrid origin with *Glycine max* and are identified as semiwild. Although their seed and plant size are larger and more vigorous than those of true wild soybeans, these semiwild accessions exhibit characteristics, such as excessive twining and shattering, which make them easier to maintain

G.A. Juvik, USDA-ARS, agronomist, GRIN, Beltsville, MD (formerly Department of Agronomy, University of Illinois, Urbana); R.L. Bernard, professor of plant genetics, Department of Agronomy, University of Illinois, Urbana (formerly USDA-ARS, research geneticist, University of Illinois, Urbana); Ruzhen Chang, Chinese Academy of Agricultural Sciences, Beijing, People's Republic of China; and J.F. Cavins, USDA-ARS, supervisory research chemist, USDA Northern Regional Research Center, Peoria, IL.

with other wild soybeans than with cultivated soybeans. These accessions are more similar to wild soybeans than those designated "semiwild" in the cultivated soybean collection.

The data in tables 3 and 4 are 2-year means (except data for PI numbers greater than PI 468.919, which are based on 1984 values only). An asterisk (*) following a mean indicates that the difference between the values for the 2 years exceeded a specified limit. The limits for the traits were as follows:

Twining date	> 7 days	Flowering date	> 7 days
Maturity date	> 7 days	Height	> 30 cm
Leaf length	> 15 mm	Leaf shape	> .15 unit
Seed weight	> 0.5 cg/seed	Pod length	> 4 mm

This system was used because of the possibility of misinterpreting the mean of only two observations when the difference between the individual values was large.

To obtain protein and oil percentages of an accession, approximately 7 g of seeds was placed in a beaker and dried in a Thelco forced air oven for 3 hours at 130°C. The seeds were then transferred to a 50-g bottle, sealed, and allowed to cool for 1 hour. A sample was then ground in a Varco electric dry-food grinder and returned to the 50-g bottle. The ground meal was analyzed by near-infrared reflectance in a Pacific-Scientific feed-grain analyzer. The analyzer calibration was checked with two sealed standards (a wheatmeal and a soymeal) and a freshly ground soymeal standard before each batch of 100 samples was processed.

Fatty acid composition was obtained by gas-liquid chromatography of the methyl esters. Seeds were ground in a small food grinder and stored at -20°C until analyzed. Approximately 200 mg was placed in a 25-ml vial, and 5 ml of sodium methoxide was added in two 2.5-ml aliquots with an automatic syringe in such a way as to ensure mixing. The sodium methoxide solution was prepared daily by adding 1 g of sodium metal to 100 ml of reagent grade methanol. The suspension of ground sample in sodium methoxide was allowed to stand for 45 minutes, after which 1 ml of 10% acetic acid solution was added followed immediately by 10 ml of heptane. The sample was completely mixed after each reagent addition. After the final mixing the sample was allowed to stand for several minutes so that the layers could separate. The heptane layer was analyzed with a Varian gas chromatograph equipped with two autoinjectors and flame detectors. Columns were 2 m by 2 mm and packed with 100/200 mesh Gas-Chrom Q coated with 5% LAC-2R-446. Analysis was made isothermally at 180°C, with the injector at 230°C and the detector at 240°C. Gas flow rates for helium, hydrogen, and air were 25, 25, and 250 ml per minute, respectively. The autoinjectors were set to inject 0.5 μ l. Total analysis time per injection was 10 minutes. Integration, peak identification, data storage, and report printing were done by computer.

Explanation of the data categories and abbreviations used are as follows:

Table 1:

PI number:

Serial numbers assigned by the Plant Introduction Office, Germplasm Resources Laboratory, USDA-ARS, BARC-West, Beltsville, MD 20705.

Accession name and foreign or collector number:

Accession names and foreign or collector numbers are reported as received. An attempt was made to standardize transliterations and to correct obvious errors. When heterogeneous introductions were received, two or more sublines were preserved and are distinguished by a letter (A, B, C, etc.) suffixed to the PI number, except PI 407.018 to PI 407.307, to which different PI numbers were assigned after sublining.

Country of acquisition:

This is the country from which the seeds were obtained.

Country of origin:

This is the country from which the accession originated, based on information received from the country of acquisition.

Year introduced:

This is the year the introduction was assigned a PI number.

Maturity group:

Classification of relative time of maturity based on data previously collected at Urbana, IL. Maturity groups include 000 (earliest), 00, 0, I, II, III, and IV.

Table 2:

Flower color:

P = purple, W = white

Pubescence color:

Ng = near gray, T = tawny

Pubescence form:

A = appressed (most hairs flat on leaf surface)

E = erect on leaf surface

Sa = semiappressed (between erect and appressed)

Va = very appressed (all hairs flat on leaf surface)

Pubescence density:

N = normal, Sp = sparse

Pod color:

Bl = black, Br = brown, Dbr = dark brown

Seed coat luster:

B = bloom (heavy coating of a powdery substance adhering to the seed coat)

D = dull (trace amounts of bloom)

Db = dense bloom (heavy bloom)

I = intermediate (between dull and shiny)

Lb = light bloom (slight bloom)

S = shiny (absence of bloom)

Seed coat and hilum color:

Bl = black, Br = brown, Gn = green

Seed shape:

Nr = near round

Ob = oblong (near rectangular in lateral outline)

Ov = oval (elliptical in lateral outline)

Other seed traits:

Fleck = brown flecks on black seed coat

Other leaf traits:

Na = narrow leaflet

Other plant traits:

4sd = four-seeded pods

Sw = semiwild (visual classification based on plant and seed characteristics, between soybean and wild soybean)

Table 3:

Twining date:

Date that 50% of the plants have begun to twine, expressed as month (mm) and day (dd).

Flower date:

Date that 50% of the plants have begun to flower, expressed as month (mm) and day (dd).

Maturity date:

Date that 95% of the pods have reached final color, expressed as month (mm) and day (dd).

Height:

Length of stem from ground to tip, in centimeters, at maturity.

Leaflet length:

Length of a typical terminal leaflet from the upper third and from the lower third of plant, in millimeters.

Leaflet shape:

Ratio of width to length of a typical terminal leaflet from the upper third and from the lower third of plant.

Pod length:

Length of a typical three-seeded pod, in millimeters.

Seed weight:

Centigrams per seed based on a 200-seed sample.

SMV:

Reaction to soybean mosaic virus based on leaf symptoms.

Scored 1 (moderately resistant) to 5 (highly susceptible).

Data are for 1985 only.

Table 4:

Seed composition:

Protein and oil: Percentage of dry weight of seeds.

Fatty acids (palmitic, stearic, oleic, linoleic, linolenic, other): Percentage of total fatty acids.

Table 1

Identification and origin information for USDA wild soybean germplasm
in maturity groups 000 to IV, PI 65.549 to PI 483.464

PI No.	Accession name	Foreign or collector No.	Country of acquisition	Country of origin	Year introduced	Maturity group
65.549			China	China	1925	II
81.762			Japan	Soviet Union	1929	II
101.404A		T106-2	China	China	1932	II
101.404B		T106-6	China	China	1932	II
135.624			China	China	1940	II
326.581		K-5820	Soviet Union	Soviet Union	1968	II
326.582A			Soviet Union	Soviet Union	1968	II
326.582B			Soviet Union	Soviet Union	1968	I
339.732			South Korea	South Korea	1969	IV
339.735A			South Korea	South Korea	1969	IV
339.735B			South Korea	South Korea	1969	IV
342.618A			Soviet Union	Soviet Union	1969	II
342.618B			Soviet Union	Soviet Union	1969	I
342.619A			Soviet Union	Soviet Union	1969	0
342.619B			Soviet Union	Soviet Union	1969	00
342.620A			Soviet Union	Soviet Union	1969	I
342.620B			Soviet Union	Soviet Union	1969	II
342.621A			Soviet Union	Soviet Union	1969	00
342.621B			Soviet Union	Soviet Union	1969	00
342.621C			Soviet Union	Soviet Union	1969	00
342.622A			Soviet Union	Soviet Union	1969	I
342.622B			Soviet Union	Soviet Union	1969	I
366.120		A-2	Japan	Japan	1971	IV
366.121		A-4	Japan	Japan	1971	IV
366.122			Japan	Japan	1971	IV
366.123		A-20	Japan	Japan	1971	IV
378.692		A-21	Japan	Japan	1973	IV
378.702			Japan	Japan	1973	IV
391.587	Ye sheng tou		China	China	1974	II
406.684	Hidaka 1		Japan	Japan	1976	III
407.029		J8	Japan	Japan	1976	IV
407.162		K1	South Korea	South Korea	1976	IV
407.166		K2	South Korea	South Korea	1976	IV
407.171		K2	South Korea	South Korea	1976	IV
407.175		K4	South Korea	South Korea	1976	IV
407.176		K4	South Korea	South Korea	1976	IV
407.182		K8	South Korea	South Korea	1976	IV
407.184		K29	South Korea	South Korea	1976	IV
407.195		K12	South Korea	South Korea	1976	IV
407.196		K12	South Korea	South Korea	1976	IV
407.200		K14	South Korea	South Korea	1976	IV
407.205		K17	South Korea	South Korea	1976	IV
407.209		K21	South Korea	South Korea	1976	IV
407.217		K24	South Korea	South Korea	1976	IV
407.275		K101	South Korea	South Korea	1976	IV
407.277		K101	South Korea	South Korea	1976	IV
407.278		K102	South Korea	South Korea	1976	IV

Table 2

Descriptive data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Matur-	ity	Flower	<u>Pubescence</u>			Pod	<u>Seed coat</u>		Hilum	Seed	<u>Other traits</u>			
				group	color	Color		color	Luster			color	shape	Seed	Leaf
65.549	II	P	T	Sa	N		B1	Db	B1	B1	Nr				
81.762	II	P	T	Sa	N		B1	Db	B1	B1	Ov				
101.404A	II	P	T	Sa	N		B1	Db	B1	B1	Nr				
101.404B	II	P	T	Sa	N		B1	Db	B1	B1	Nr	Fleck			
135.624	II	P	T	Sa	N		B1	B	B1	B1	Ov				
326.581	II	P	T	Sa	N		B1	Db	B1	B1	Ob				
326.582A	II	P	T	Va	N		B1	Lb	B1	B1	Ov	Fleck			
326.582B	I	P	T	Va	N		B1	B	B1	B1	Ob				
339.732	IV	P	T	Va	N		B1	B	B1	B1	Ob	Fleck			
339.735A	IV	P	T	A	N		B1	Lb	B1	B1	Ov	Fleck			
339.735B	IV	P	T	A	N		B1	B	B1	B1	Ov	Fleck			
342.618A	II	P	T	Va	N		B1	Lb	B1	B1	Ob	Fleck			
342.618B	I	P	T	Va	N		B1	B	B1	B1	Ov				
342.619A	0	P	T	A	N		Br	Db	B1	B1	Ob	Fleck	Sw		
342.619B	00	P	T	Va	N		Br	Db	B1	B1	Ob	Fleck	Sw		
342.620A	I	P	T	Va	N		B1	Db	B1	B1	Ob				
342.620B	II	P	T	Va	N		B1	B	B1	B1	Ov	Fleck			
342.621A	00	P	T	Va	N		B1	Db	B1	B1	Ob				
342.621B	00	P	T	Va	Sp		B1	Db	B1	B1	Ob				
342.621C	00	P	T	Va	N		B1	B	B1	B1	Ob				
342.622A	I	P	T	Va	N		B1	Db	B1	B1	Ov	Fleck			
342.622B	I	P	T	Va	N		B1	Db	B1	B1	Ov	Fleck			
366.120	IV	P	T	A	N		B1	D	B1	B1	Ob	Fleck			
366.121	IV	P	T	Sa	N		B1	D	B1	B1	Ob	Fleck			
366.122	IV	P	T	Sa	N		B1	D	B1	B1	Ob	Fleck			
366.123	IV	P	T	Va	N		B1	B	B1	B1	Ob	Fleck			
378.692	IV	P	T	Va	N		B1	B	B1	B1	Ob	Fleck			
378.702	IV	P	T	Sa	N		B1	B	B1	B1	Ob	Fleck			
391.587	II	P	T	A	N		B1	Db	B1	B1	Ov	Fleck			
406.684	III	P	T	Va	N		B1	Db	B1	B1	Ob				
407.029	IV	P	T	A	N		B1	Lb	B1	B1	Ob	Fleck			
407.162	IV	P	T	Va	N		B1	B	B1	B1	Ob	Fleck			
407.166	IV	P	T	Va	N		B1	Lb	B1	B1	Ov	Fleck			
407.171	IV	P	T	A	N		B1	B	B1	B1	Ov	Fleck			
407.175	IV	P	T	Va	N		B1	B	B1	B1	Ov	Fleck			
407.176	IV	P	T	Va	N		B1	Db	B1	B1	Ov	Fleck			
407.182	IV	P	T	Va	N		B1	B	B1	B1	Nr	Fleck			
407.184	IV	P	T	A	N		B1	Db	B1	B1	Ov	Fleck			
407.195	IV	P	T	A	N		B1	B	B1	B1	Ob	Fleck			
407.196	IV	P	T	A	N		B1	Lb	B1	B1	Ob	Fleck			
407.200	IV	P	T	Va	N		B1	B	B1	B1	Ob	Fleck			
407.205	IV	P	T	A	N		B1	Lb	B1	B1	Ov	Fleck			
407.209	IV	P	T	Va	N		B1	Lb	B1	B1	Ob	Fleck			
407.217	IV	P	T	A	N		B1	B	B1	B1	Ov	Fleck			
407.275	IV	P	T	Va	N		B1	B	B1	B1	Ov	Fleck			
407.277	IV	P	T	A	N		B1	Lb	B1	B1	Ob	Fleck			
407.278	IV	P	T	A	N		B1	B	B1	B1	Ov	Fleck			

Table 3

Agronomic data for USDA wild soybean germplasm in maturity groups
000 to IV, PI 65.549 to PI 483.464

Entry	Twining	Flower	Maturity	Leaflet length				Leaflet shape		Pod	Seed
	date (mmdd)	date (mmdd)	date (mmdd)	Height (cm)	lower (mm)	upper (mm)	lower (mm)	upper (mm)	length (mm)	weight (cg/sd)	SMV
65.549	627	729	914	100*	50	45*	0.44	0.40	21	1.4	4
81.762	626	724	913	110	50	46	0.47*	0.37	21	1.4	3
101.404A	628*	727	914	100*	53*	46*	0.50	0.40	20	1.3	2
101.404B	618	724	913	102*	48	49*	0.49	0.35*	22	1.5	3
135.624	623	725	913	98*	43	41*	0.60	0.33	23	1.4	2
326.581	620	720	905	123*	40	44*	0.54*	0.40	24	1.4	3
326.582A	619	726	911	77	43	33	0.54*	0.42*	24	1.7	4
326.582B	625	727	912	89*	48*	45*	0.46*	0.37	23	1.5	3
339.732	624	817	1002*	55	34	29	0.65	0.47	26	1.6*	5
339.735A	625	818	1011	129	53	41	0.57	0.50	24	1.8*	3
339.735B	622	819	1009	81	50	44	0.50	0.41	27*	1.6	5
342.618A	701*	728	910	67	48	39	0.50*	0.39	24	1.6	4
342.618B	624	725	912	79*	44	41	0.46*	0.34	22	1.4	3
342.619A	622	626	821*	83*	61	65*	0.50	0.53*	35*	6.1*	1
342.619B	623	625	821*	58*	66	68*	0.56	0.41*	39*	6.7*	2
342.620A	624	725	909	75	55	42	0.40*	0.28	23	1.3	3
342.620B	619	724	912	71	54	49*	0.47*	0.32	22	1.6	4
342.621A	627*	625	813*	27	40	40	0.46	0.28	26	1.7	5
342.621B	626	627	811	45*	43	44*	0.37	0.25	24	2.0*	4
342.621C	623	627	812	55*	53*	61*	0.38	0.29	26*	2.1*	4
342.622A	624	717*	830	79*	40	43*	0.44	0.31	22	1.3	4
342.622B	620	709	828	54	33	23	0.39	0.30	24	1.1	4
366.120	619	821*	1012*	126	57	48	0.55*	0.38*	28	2.0	4
366.121	619	817	1012*	146	61*	46	0.49	0.39	27	1.8*	4
366.122	618	825*	1011*	135*	52	41	0.46	0.40	28	1.7*	4
366.123	625	817	1005*	117*	52	41	0.47*	0.37	24	1.5*	4
378.692	621	816	1006*	117*	45*	44	0.50	0.36	25	1.5	4
378.702	623	818	1009*	94*	55	45	0.58	0.34	24	2.2	4
391.587	622	730	912	66*	43	43	0.53*	0.37	22	1.4	4
406.684	625	805*	1002	139*	65*	52*	0.42	0.43	33	2.3*	3
407.029	623	815	1007*	79	47	38	0.50*	0.37	27	2.0	4
407.162	705	820	1009	91	45	40	0.66*	0.39	27	2.4	3
407.166	704	818	1007*	113*	53	43	0.56	0.51	26	1.9	3
407.171	704	821	1009	73	57	45	0.58	0.48	26	1.7	4
407.175	705	821	1008*	60	45	44	0.51*	0.33	23	1.7	4
407.176	629	820	1008*	73*	51	47*	0.57	0.38	25	1.9	3
407.182	629	818	1010	98*	61	46	0.44	0.40	23	1.6	3
407.184	630*	819	1008*	80*	49	42	0.50	0.34	26	1.9	4
407.195	625	814	1007*	99	57	44	0.65*	0.37	26	2.1	4
407.196	623	814	1007*	86*	51	35	0.48	0.45	29	1.9	4
407.200	706	818	1008*	68*	45	35	0.61	0.53	24	2.1*	3
407.205	623	817	1007*	91	45	35	0.64*	0.42	23	2.1	4
407.209	702	821	1010*	79	45	37	0.54	0.47	26	2.3	3
407.217	706	820	1006*	61	52*	55*	0.45	0.39*	26	1.9	4
407.275	701	817	1011	102*	52	49*	0.56*	0.38	24	1.6	3
407.277	622	818	1009	144*	67	52*	0.59	0.46	25	1.9	4
407.278	623	817	1010	103*	49	40	0.56	0.47	27	2.2	4

Table 4

Seed composition data for USDA wild soybean germplasm in
maturity groups 000 to IV, PI 65.549 to PI 483.464

Entry	Matur-									
		ity group	Protein (%)	Oil (%)	Palmitic (%)	Stearic (%)	Oleic (%)	Lino-leic (%)	Lino-lenic (%)	Other (%)
65.549	II	51.4	8.7	14.1	2.6	8.9	55.3	19.2	0.0	
81.762	II	51.9	8.0	14.3	2.6	9.0	55.5	18.6	0.0	
101.404A	II	50.9	8.3	14.2	2.8	9.7	55.3	18.1	0.0	
101.404B	II	50.9	8.3	14.0	2.7	10.0	55.1	18.2	0.1	
135.624	II	51.1	8.7	12.9	2.8	9.5	54.8	20.0	0.1	
326.581	II	49.7	8.2	12.3	3.2	13.5	53.2	17.9	0.1	
326.582A	II	48.6	10.2	12.9	2.8	15.4	53.8	15.3	0.0	
326.582B	I	48.7	9.8	13.5	2.8	11.2	55.8	16.8	0.1	
339.732	IV	50.5	8.1	14.5	3.0	11.5	54.3	16.8	0.1	
339.735A	IV	49.4	9.3	14.1	3.2	11.6	54.6	16.7	0.0	
339.735B	IV	47.5	9.2	13.7	3.0	12.7	54.7	16.0	0.0	
342.618A	II	47.8	10.6	13.7	3.0	11.1	54.7	17.6	0.1	
342.618B	I	49.0	9.9	13.4	2.7	11.0	55.7	17.2	0.0	
342.619A	0	43.4	16.5	11.9	3.7	13.3	59.5	11.6	0.1	
342.619B	00	45.1	16.0	12.1	3.7	13.7	59.6	11.0	0.1	
342.620A	I	49.3	9.1	13.2	2.7	9.0	52.5	22.7	0.0	
342.620B	II	51.6	8.1	13.4	3.2	11.8	54.9	16.7	0.1	
342.621A	00	46.3	9.9	12.7	2.6	15.0	56.0	13.8	0.1	
342.621B	00	46.7	11.5	12.5	2.6	11.8	57.5	15.5	0.1	
342.621C	00	48.1	11.4	12.5	2.8	12.9	57.0	14.9	0.1	
342.622A	I	49.0	8.3	13.8	3.5	11.6	52.3	18.8	0.1	
342.622B	I	46.3	8.8	13.2	2.8	11.3	56.7	16.1	0.0	
366.120	IV	51.2	8.2	14.3	2.8	10.5	54.3	19.1	0.0	
366.121	IV	52.5	6.5	13.4	2.6	8.3	52.6	23.2	0.1	
366.122	IV	52.2	6.8	14.1	3.0	8.8	52.2	22.2	0.1	
366.123	IV	51.7	8.0	13.6	3.2	10.0	53.9	19.5	0.0	
378.692	IV	52.8	7.0	13.4	3.2	9.9	53.6	20.0	0.1	
378.702	IV	51.2	8.3	13.0	2.7	10.7	54.9	18.7	0.1	
391.587	II	48.4	9.4	13.3	3.2	11.9	54.9	16.8	0.1	
406.684	III	51.1	8.7	12.9	3.0	11.8	55.7	16.6	0.0	
407.029	IV	50.4	8.5	13.8	3.2	10.9	54.1	18.2	0.1	
407.162	IV	50.8	9.4	13.8	3.0	12.8	55.6	14.8	0.1	
407.166	IV	50.1	9.7	13.9	3.0	11.8	55.7	15.9	0.0	
407.171	IV	48.2	8.7	15.4	2.7	10.4	54.1	17.5	0.1	
407.175	IV	49.3	9.7	12.6	3.0	12.5	57.8	14.2	0.0	
407.176	IV	48.9	9.2	14.0	2.9	11.5	54.9	16.7	0.1	
407.182	IV	49.1	9.2	13.7	2.7	11.5	53.5	18.7	0.0	
407.184	IV	50.1	8.3	13.5	2.6	11.8	56.6	15.5	0.0	
407.195	IV	50.9	8.7	14.3	2.9	11.0	55.7	16.2	0.1	
407.196	IV	50.2	9.4	13.8	3.1	10.8	56.4	16.0	0.1	
407.200	IV	49.4	10.1	12.9	2.4	12.1	57.1	15.7	0.1	
407.205	IV	50.3	9.5	12.3	2.7	11.6	57.4	16.1	0.1	
407.209	IV	49.0	10.9	13.5	2.4	13.1	55.7	15.5	0.0	
407.217	IV	49.3	8.9	14.0	2.5	11.4	54.7	17.6	0.0	
407.275	IV	49.1	9.2	13.3	2.5	11.2	54.6	18.6	0.0	
407.277	IV	50.1	8.5	14.1	2.7	11.8	53.3	18.2	0.0	
407.278	IV	48.0	10.6	12.2	2.7	13.9	57.0	14.2	0.0	

Table 1

Identification and origin information for USDA wild soybean germplasm
in maturity groups 000 to IV, PI 65.549 to PI 483.464

PI No.	Accession name	Foreign or collector No.	Country of acquisition	Country of origin	Year introduced	Matur- ity group
407.288		W1	China	China	1976	II
407.289		W1	China	China	1976	II
407.290		W2	China	China	1976	II
407.291		W2	China	China	1976	II
407.292		W3	China	China	1976	II
407.293		W3	China	China	1976	II
407.294		W3	China	China	1976	II
407.295		W3	China	China	1976	II
407.296		W4	China	China	1976	II
407.297		W4	China	China	1976	II
407.298		W4	China	China	1976	II
407.299		W4	China	China	1976	II
407.320			South Korea	South Korea	1976	IV
407.321			South Korea	South Korea	1976	IV
423.988	J-077630	F7	Soviet Union	Soviet Union	1978	00
423.989A	J-077631	F8	Soviet Union	Soviet Union	1978	00
423.989B	J-077631	F8	Soviet Union	Soviet Union	1978	0
423.990A	J-077632	F9	Soviet Union	Soviet Union	1978	0
423.990B	J-077632	F9	Soviet Union	Soviet Union	1978	0
423.991	J-077633	F10	Soviet Union	Soviet Union	1978	0
423.992	J-077634	F11	Soviet Union	Soviet Union	1978	00
423.993	J-077635	F12	Soviet Union	Soviet Union	1978	00
423.994	J-077636	F13	Soviet Union	Soviet Union	1978	0
423.995	J-077637	F14	Soviet Union	Soviet Union	1978	0
423.996	J-077638	F15	Soviet Union	Soviet Union	1978	00
423.997	J-077639	F16	Soviet Union	Soviet Union	1978	00
423.998	J-077640	F17	Soviet Union	Soviet Union	1978	00
423.999A	J-077641	F18	Soviet Union	Soviet Union	1978	00
423.999B	J-077641	F18	Soviet Union	Soviet Union	1978	00
424.000	J-077642	F19	Soviet Union	Soviet Union	1978	00
424.001	J-077643	F20	Soviet Union	Soviet Union	1978	00
424.002	J-077644	F21	Soviet Union	Soviet Union	1978	00
424.003	J-077645	F22	Soviet Union	Soviet Union	1978	00
424.004A		74001	South Korea	South Korea	1978	II
424.004B		74001	South Korea	South Korea	1978	II
424.005		74002	South Korea	South Korea	1978	IV
424.006A		74004	South Korea	South Korea	1978	IV
424.006B		74004	South Korea	South Korea	1978	IV
424.032		74035	South Korea	South Korea	1978	IV
424.033		74036	South Korea	South Korea	1978	IV
424.034		74037	South Korea	South Korea	1978	IV
424.036		74039	South Korea	South Korea	1978	IV
424.038A		74053	South Korea	South Korea	1978	IV
424.051A		74066	South Korea	South Korea	1978	IV
424.051B		74066	South Korea	South Korea	1978	IV
424.063		74043	South Korea	South Korea	1978	IV
424.067		74048	South Korea	South Korea	1978	IV

Table 2

Descriptive data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Matur-		Flower <u>Pubescence</u>			Pod color	Seed coat		Hilum	Seed	Other traits		
	ity group	color	Color	Form	Density		Luster	Color	color	shape	Seed	Leaf	Plant
407.288	II	P	T	Va	N	B1	B	B1	B1	Ov			
407.289	II	P	T	Va	N	B1	B	B1	B1	Ov	Fleck		
407.290	II	P	T	Sa	N	B1	B	B1	B1	Ob	Fleck		
407.291	II	P	T	Sa	N	B1	Db	B1	B1	Ob	Fleck		
407.292	II	P	T	Va	N	B1	B	B1	B1	Ov			
407.293	II	P	T	Va	N	B1	Db	B1	B1	Ov			
407.294	II	P	T	Sa	N	B1	Db	B1	B1	Ov	Fleck		
407.295	II	P	T	Va	N	B1	B	B1	B1	Ov			
407.296	II	P	T	Va	N	B1	B	B1	B1	Ov	Fleck		
407.297	II	P	T	Va	N	B1	B	B1	B1	Ov	Fleck		
407.298	II	P	T	Va	N	B1	B	B1	B1	Ov	Fleck		
407.299	II	P	T	Va	N	B1	Db	B1	B1	Ov	Fleck		
407.320	IV	P	T	A	N	B1	B	B1	B1	Ob	Fleck		
407.321	IV	P	T	A	N	B1	Lb	B1	B1	Ob	Fleck		
423.988	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.989A	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.989B	0	P	T	Sa	N	B1	I	B1	B1	Ob	Fleck		
423.990A	0	P	T	A	N	Br	Db	B1	B1	Ob	Fleck		
423.990B	0	P	T	A	N	B1	Db	B1	B1	Ob			
423.991	0	P	T	A	N	B1	Db	B1	B1	Ob			
423.992	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.993	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.994	0	P	T	A	N	B1	Db	B1	B1	Ob			
423.995	0	P	T	A	N	B1	B	B1	B1	Ob			
423.996	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.997	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.998	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.999A	00	P	T	Va	N	B1	Db	B1	B1	Ob			
423.999B	00	P	T	Va	N	B1	B	B1	B1	Ob			
424.000	00	P	T	Va	N	B1	Db	B1	B1	Ob			
424.001	00	P	T	A	N	Dbr	D	B1	B1	Ov	Fleck		
424.002	00	P	T	Va	N	B1	B	B1	B1	Ov		4sd	
424.003	00	P	T	Va	N	B1	Db	B1	B1	Ob			
424.004A	II	P	T	E	N	B1	B	B1	B1	Ob	Fleck		
424.004B	II	P	T	E	N	B1	Lb	B1	B1	Ob	Fleck		
424.005	IV	W	T	A	N	B1	B	Gn	B1	Ob		Sw	
424.006A	IV	P	T	Sa	N	B1	B	B1	Br	Ob	Fleck		
424.006B	IV	P	T	Sa	N	B1	B	B1	Br	Ob	Fleck		
424.032	IV	P	T	A	N	B1	Db	B1	B1	Ob	Fleck		
424.033	IV	P	T	A	N	B1	B	B1	B1	Ob	Fleck		
424.034	IV	P	T	A	N	B1	B	B1	B1	Ob	Fleck		
424.036	IV	P	T	A	N	B1	Lb	B1	B1	Ov	Fleck		
424.038A	IV	P	T	A	N	B1	Lb	B1	B1	Ov	Fleck		
424.051A	IV	P	T	Va	N	B1	B	B1	B1	Ob			
424.051B	IV	P	T	Va	N	B1	Db	B1	B1	Ob			
424.063	IV	P	T	Va	N	B1	Lb	B1	B1	Ob	Fleck		
424.067	IV	P	T	A	N	B1	Lb	B1	B1	Ob	Fleck		

Table 3

Agronomic data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Twining	Flower	Maturity	Leaflet length				Leaflet shape		Pod	Seed
	date (mmdd)	date (mmdd)	date (mmdd)	Height (cm)	lower (mm)	upper (mm)	lower (mm)	upper (mm)	length (mm)	weight (cg/sd)	SMV
407.288	628*	725*	910	54	38	39*	0.43*	0.42	22	1.2	4
407.289	624	724	907	118*	42	46	0.45*	0.27	22	1.2	4
407.290	625	725	910	74*	45	44	0.48*	0.30	25	2.0	3
407.291	626	721	908	63	39	39	0.47*	0.35	22	1.8	2
407.292	628	726*	910	62*	35	38	0.54*	0.33*	23	1.6	4
407.293	626	727*	909*	88*	45	43*	0.35*	0.29	23*	1.7*	4
407.294	619	722	906	70	29	33*	0.56*	0.40	22	1.7	4
407.295	627*	727	909	86	45	35	0.50*	0.38*	21	1.6	3
407.296	625	805	914	91*	48	54*	0.45*	0.28	22	1.3	3
407.297	624	718*	914	80*	49	43*	0.52*	0.41*	21	1.2	3
407.298	702*	724*	917	79*	48	51*	0.46*	0.32	22	1.2	2
407.299	626	727	913	76	54	48*	0.40	0.32	19	1.1	3
407.320	703	823	1009*	86	53	43	0.63*	0.33	26	2.0	4
407.321	705*	822	1012	73	51	44	0.46*	0.34	25	1.5	3
423.988	621	625*	811*	56*	56	59*	0.36*	0.22	24	2.1*	4
423.989A	620	623	812*	69*	47	55*	0.38	0.23	24	2.1*	3
423.989B	627*	701*	824*	60*	43	42*	0.44	0.34	26*	2.0	4
423.990A	629	701*	818	55*	41*	39	0.43	0.21	24	1.7*	4
423.990B	625	702	816	56*	28	36*	0.44	0.28*	23	1.4	4
423.991	625	701	817	38	31	32*	0.41	0.40*	22	1.4	4
423.992	625	626	811*	47	41	49*	0.41	0.28	25	2.0*	4
423.993	628*	627*	814*	38	40	37	0.47	0.22	23	1.5	5
423.994	623	701	816	68*	38	36*	0.43	0.29	24	1.5	4
423.995	619	625	812*	60*	40	50*	0.44	0.17	22	1.9	4
423.996	620	624*	813*	56	42	46*	0.39	0.29	24	1.9	4
423.997	621	625	810*	55*	45	56*	0.43	0.25	22	1.9*	4
423.998	624	626	811*	59*	41	54*	0.43	0.32	24	1.9*	5
423.999A	628	628	812*	42	45	56*	0.45	0.29	24	1.8*	4
423.999B	622	626	813*	55*	43	41*	0.47	0.30*	28	1.8*	5
424.000	626*	628	814*	28	38	43	0.48	0.29	24	1.9*	4
424.001	629*	630*	813*	42	42	49*	0.49	0.32	27*	3.3*	4
424.002	619	629	812	53*	40	44*	0.44*	0.26	23	1.6	4
424.003	620	626	812	47*	45	52*	0.41	0.24	24	1.9*	4
424.004A	625	816	925	127*	60	54*	0.55*	0.43	26	2.1	4
424.004B	625	814	925	120*	59	54*	0.54*	0.39	26	1.8	2
424.005	717*	730*	1006	81	83	62	0.41	0.32	30	3.7*	2
424.006A	702	830	1011*	121	49	42	0.65*	0.57*	26	1.6	4
424.006B	630	831	1011*	121	56	40	0.57	0.46	27	1.5	2
424.032	702	820	1008*	74	47	36	0.61	0.47	24	2.0	4
424.033	704	822	1006*	62	39	31	0.74*	0.42	24	1.8	5
424.034	630	823	1009	91*	52	38	0.68	0.46	25	1.8	4
424.036	630	820	1008*	93	51	37	0.63*	0.41	25	1.9	4
424.038A	707	824	1009	67	45	38	0.71	0.44	25	1.8	4
424.051A	626	819	1009*	84	41	34	0.64*	0.43	26	2.3	4
424.051B	623	817	1006*	82*	43	36	0.63	0.47	25	2.6	4
424.063	702	822	1011	104*	48	35	0.55	0.45	24	1.7	3
424.067	629	818	1011	83	54	44	0.55	0.45	25	1.8	4

Table 4

Seed composition data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464

Entry	Maturity group	Protein (%)	Oil (%)	Palmitic (%)	Stearic (%)	Oleic (%)	Linoleic (%)	Linenic (%)	Other (%)
407.288	II	49.8	8.9	13.1	3.1	11.1	55.3	17.4	0.1
407.289	II	49.6	9.5	13.3	2.9	12.1	54.7	17.0	0.1
407.290	II	49.9	10.0	14.0	2.9	13.3	53.6	16.3	0.1
407.291	II	49.0	10.6	13.4	3.1	15.1	53.6	14.8	0.1
407.292	II	48.8	10.1	12.7	3.0	14.7	54.3	15.3	0.1
407.293	II	49.1	10.2	13.1	2.8	12.1	54.5	17.5	0.1
407.294	II	51.2	8.7	13.5	2.9	12.2	53.8	17.7	0.1
407.295	II	48.4	10.4	13.5	2.9	11.8	55.4	16.5	0.1
407.296	II	47.9	9.0	12.7	2.8	11.2	55.6	17.9	0.0
407.297	II	49.6	8.6	12.9	2.8	10.3	56.1	18.0	0.1
407.298	II	48.3	9.3	12.7	2.9	10.7	56.2	17.5	0.1
407.299	II	48.8	8.9	13.2	3.0	12.6	54.1	17.2	0.1
407.320	IV	49.5	9.8	13.4	2.9	13.9	55.9	13.9	0.2
407.321	IV	49.7	9.4	13.8	3.0	11.2	54.7	17.3	0.1
423.988	00	48.4	11.0	11.5	3.3	16.0	56.6	12.6	0.1
423.989A	00	48.1	10.9	11.5	3.2	15.2	57.7	12.6	0.1
423.989B	0	47.1	11.2	12.1	3.3	12.2	58.3	14.1	0.1
423.990A	0	47.6	10.0	12.8	3.3	12.2	57.0	14.8	0.0
423.990B	0	45.4	8.2	13.0	3.7	11.5	56.7	15.3	0.0
423.991	0	49.0	9.1	13.0	3.4	11.9	56.7	15.3	0.0
423.992	00	47.6	11.3	12.6	2.9	12.8	56.4	15.3	0.1
423.993	00	46.5	11.0	11.8	3.7	18.3	55.3	11.0	0.0
423.994	0	48.5	9.8	13.0	3.4	11.3	56.9	15.5	0.1
423.995	0	47.7	11.2	13.6	3.8	17.4	54.0	11.2	0.1
423.996	00	48.3	11.0	11.7	3.1	14.3	57.8	13.2	0.1
423.997	00	48.0	10.7	12.5	2.7	13.0	56.6	15.2	0.1
423.998	00	46.9	11.0	12.6	2.8	12.6	57.2	15.0	0.1
423.999A	00	47.9	10.2	12.6	2.7	11.8	57.4	15.7	0.1
423.999B	00	47.5	10.2	12.6	2.7	11.4	57.7	15.7	0.1
424.000	00	47.4	9.4	12.8	2.7	13.0	56.7	14.7	0.1
424.001	00	44.4	14.1	13.0	3.1	20.9	51.7	11.4	0.1
424.002	00	49.3	10.4	12.9	3.0	13.5	55.1	15.7	0.0
424.003	00	46.7	11.0	13.4	3.4	17.4	54.7	11.2	0.1
424.004A	II	49.9	9.2	13.9	2.9	13.2	52.9	17.2	0.1
424.004B	II	50.1	9.1	13.7	2.7	11.1	54.4	18.1	0.1
424.005	IV	46.3	12.6	14.4	3.4	12.7	56.5	13.1	0.1
424.006A	IV	50.3	9.0	14.3	3.0	10.1	52.7	20.0	0.0
424.006B	IV	50.9	8.9	13.6	2.5	8.9	52.6	22.5	0.0
424.032	IV	52.0	8.5	15.3	3.1	11.0	53.8	17.0	0.0
424.033	IV	52.7	8.7	15.1	3.2	10.9	53.9	17.0	0.0
424.034	IV	50.6	9.3	14.1	3.1	11.7	54.7	16.5	0.0
424.036	IV	49.5	9.7	14.0	3.0	12.6	54.2	16.2	0.0
424.038A	IV	49.1	10.7	12.5	3.1	11.7	59.9	13.0	0.0
424.051A	IV	49.9	9.3	14.0	3.2	12.9	55.6	14.5	0.0
424.051B	IV	49.8	9.7	13.9	3.2	13.5	55.6	14.0	0.0
424.063	IV	49.8	9.2	14.1	3.1	11.7	54.7	16.6	0.0
424.067	IV	50.8	9.5	14.6	3.1	15.7	50.9	15.7	0.0

Table 1

Identification and origin information for USDA wild soybean germplasm
in maturity groups 000 to IV, PI 65.549 to PI 483.464

PI No.	Accession name	Foreign or collector No.	Country of acquisition	Country of origin	Year introduced	Maturity group
424.068		74049	South Korea	South Korea	1978	IV
424.071		74052	South Korea	South Korea	1978	IV
424.079		74079	South Korea	South Korea	1978	IV
424.088		74088	South Korea	South Korea	1978	IV
424.091A		74091	South Korea	South Korea	1978	IV
424.091B		74091	South Korea	South Korea	1978	IV
424.116		74116	South Korea	South Korea	1978	IV
440.913A			China	China	1980	II
440.913B			China	China	1980	II
447.003A		50204	China	China	1980	0
447.003B		50204	China	China	1980	I
447.004		50368	China	China	1980	III
458.535			China	China	1981	0
458.536			China	China	1981	0
458.537A			China	China	1981	000
458.537B			China	China	1981	000
458.538			China	China	1981	000
458.539A			China	China	1981	000
458.539B			China	China	1981	0
458.540A			China	China	1981	0
458.540B			China	China	1981	0
458.540C			China	China	1981	0
458.540D			China	China	1981	0
464.866A		Long 79-0009	China	China	1982	00
464.866B		Long 79-0009	China	China	1982	00
464.867		Long 79-0304	China	China	1982	000
464.868A		Long 79-0402	China	China	1982	00
464.868B		Long 79-0402	China	China	1982	00
464.869A		Long 79-1809	China	China	1982	0
464.869B		Long 79-1809	China	China	1982	0
464.870		Long 79-3313	China	China	1982	000
464.871A		Long 79-5801	China	China	1982	0
464.871B		Long 79-5801	China	China	1982	0
464.871C		Long 79-5801	China	China	1982	0
464.889A		Gong di 685	China	China	1982	II
464.889B		Gong di 685	China	China	1982	II
464.889C		Gong di 685	China	China	1982	II
464.890A		Gong di 2019	China	China	1982	II
464.890B		Gong di 2019	China	China	1982	I
464.891A		Gong di 2022	China	China	1982	I
464.891B		Gong di 2022	China	China	1982	II
464.891C		Gong di 2022	China	China	1982	II
464.892		Gong di 2024	China	China	1982	I
464.925A		LS-001	China	China	1982	0
464.925B		LS-001	China	China	1982	0
464.925C		LS-001	China	China	1982	I
464.926		LS-004	China	China	1982	000

Table 2

Descriptive data for USDA wild soybean germplasm in maturity groups
000 to IV, PI 65.549 to PI 483.464

Entry	Matur-	Flower				Pubescence		Pod	Seed coat		Hilum	Seed	Other traits		
		group	color	Color	Form	Density	color		Luster	Color	color	shape	Seed	Leaf	Plant
424.068	IV	P	T	A	N		B1	B	B1	B1	Ob		Fleck		
424.071	IV	P	T	A	N		B1	B	B1	B1	Ob				
424.079	IV	P	T	Va	N		B1	B	B1	B1	Ov		Fleck		
424.088	IV	P	T	A	N		B1	Lb	B1	B1	Ob		Fleck		
424.091A	IV	P	T	Va	N		B1	Lb	B1	B1	Ob		Fleck		
424.091B	IV	P	T	A	N		B1	B	B1	B1	Ob		Fleck		
424.116	IV	P	T	Sa	N		B1	Lb	B1	B1	Ob		Fleck		
440.913A	II	P	T	Va	N		B1	Db	B1	B1	Ov		Fleck		
440.913B	II	P	T	Va	N		B1	Db	B1	B1	Ov		Fleck		
447.003A	0	P	T	A	N		B1	B	B1	B1	Ov		Fleck		
447.003B	I	P	T	Va	N		B1	B	B1	B1	Ob		Fleck		
447.004	III	P	T	Va	N		B1	B	B1	B1	Ob				
458.535	0	P	T	A	N		B1	B	B1	B1	Ob				
458.536	0	P	T	E	N		B1	B	B1	B1	Ov				
458.537A	000	P	T	Va	N		B1	Lb	B1	B1	Ob				
458.537B	000	P	T	Va	N		B1	B	B1	B1	Ob				
458.538	000	P	T	Va	N		B1	B	B1	B1	Ob				
458.539A	000	P	T	Va	N		B1	B	B1	B1	Ob				
458.539B	0	P	T	A	N		B1	B	B1	B1	Ob				
458.540A	0	P	T	Sa	N		B1	Db	B1	B1	Ob				
458.540B	0	P	T	Sa	N		B1	Db	B1	B1	Ov				
458.540C	0	P	T	Sa	N		B1	Db	B1	B1	Ov				
458.540D	0	P	T	Sa	N		B1	B	B1	B1	Ov				4sd
464.866A	00	P	T	Va	N		B1	B	B1	B1	Ob				
464.866B	00	P	T	Va	N		B1	B	B1	B1	Ob				
464.867	000	P	T	Va	N		B1	B	B1	B1	Ob				
464.868A	00	P	T	Sa	N		B1	B	B1	B1	Ov				
464.868B	00	P	T	Va	N		B1	B	B1	B1	Ob				
464.869A	0	P	T	E	N		B1	B	B1	B1	Ov				
464.869B	0	P	T	E	N		B1	Db	B1	B1	Ov				
464.870	000	P	T	Va	N		B1	Db	B1	B1	Ob				
464.871A	0	P	T	Sa	N		B1	B	B1	B1	Ov				
464.871B	0	P	T	Sa	N		B1	B	B1	B1	Ov				
464.871C	0	P	T	Sa	N		B1	B	B1	B1	Ov				
464.889A	II	P	T	Va	N		B1	Db	B1	B1	Ov				
464.889B	II	P	T	Va	N		B1	Db	B1	B1	Ov				
464.889C	II	P	T	Va	N		B1	Db	B1	B1	Ov				
464.890A	II	P	T	Sa	N		B1	B	B1	B1	Ob		Fleck		
464.890B	I	P	T	Sa	N		B1	B	B1	B1	Ob		Fleck		
464.891A	I	P	T	Va	N		B1	B	B1	B1	Ov				
464.891B	II	P	T	E	N		B1	B	B1	B1	Ov				4sd
464.891C	II	P	T	E	N		B1	B	B1	B1	Ov				
464.892	I	P	T	Va	N		B1	B	B1	B1	Ob		Fleck		
464.925A	0	P	T	E	N		B1	B	B1	B1	Ov				
464.925B	0	P	T	E	N		B1	Db	B1	B1	Ov				
464.925C	I	P	T	Sa	N		B1	B	B1	B1	Ov				
464.926	000	P	T	Va	N		B1	Db	B1	B1	Ob				

Table 3

Agronomic data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Twining	Flower	Maturity	Leaflet length				Leaflet shape		Pod	Seed
	date (mmdd)	date (mmdd)	date (mmdd)	Height (cm)	lower (mm)	upper (mm)	lower (mm)	upper (mm)	length (mm)	weight (cg/sd)	SMV (score)
424.068	630	817	1008*	100*	43	37	0.70	0.49	24	2.0	4
424.071	624	818	1007*	102*	50	35	0.60*	0.51	27	2.4	4
424.079	629*	817	1007*	104*	49	40	0.63*	0.47	25	1.6	4
424.088	625	819	1010*	70	54	46	0.54	0.34	27	2.0	4
424.091A	630	824	1010*	85	53	40	0.46	0.40	27	2.2	5
424.091B	630	825*	1010*	107*	55	38	0.57	0.43*	27	2.2	4
424.116	702	826	1017	92	57*	44	0.62*	0.45	25	1.8	4
440.913A	620	730	917	93*	41	40*	0.48*	0.35	24	1.4	3
440.913B	623	726	917	82	41	37	0.52*	0.36*	24	1.4	3
447.003A	623	707	825*	63*	37	37*	0.52	0.35*	23	1.2	4
447.003B	623	711	901	79*	31	47*	0.56	0.28	23	1.5	5
447.004	622	731	919	115	49	44*	0.53*	0.38	23	1.5	4
458.535	625	630	817	45	37	33	0.48	0.30	24	1.7*	5
458.536	623	717*	830*	68*	29	32*	0.67	0.31	23	1.4	2
458.537A	620	623	808*	61	45	49*	0.49	0.31	26*	1.8*	3
458.537B	621	624	808	55*	49	48*	0.39	0.27	25	2.0*	3
458.538	623	627	802	38	50	43	0.41	0.34	26	1.9*	4
458.539A	623	626	809*	54	47	44	0.39	0.33	24	1.9*	3
458.539B	625	628	816	51	36	34*	0.53	0.38	25	1.7	4
458.540A	620	709	825	87*	38*	37*	0.44	0.41	22	1.9	3
458.540B	618	711	826	72*	35	34*	0.60*	0.37	21*	1.8	4
458.540C	620	708	825	91*	42	40*	0.50	0.39	24	1.7	4
458.540D	621	708	823	84*	33	38*	0.54	0.36	22	1.5	4
464.866A	627	629	815	52*	39	44*	0.53	0.40*	24	1.6	5
464.866B	704	707*	817	15	23	24	0.57	0.49*	20	1.5	5
464.867	625	627	809*	43	44	42*	0.41	0.37*	25*	1.7*	4
464.868A	625	712	821	80*	38	36*	0.54	0.28	22	1.4	5
464.868B	625	629	815	55*	35	38*	0.52*	0.39	24	1.6	4
464.869A	623	718	830	93*	39	37*	0.51	0.36	22	1.2	2
464.869B	621	718	901	80*	39	42*	0.60	0.28	23	1.4	3
464.870	625	629	810*	50*	52*	55*	0.34	0.28	26	1.7*	4
464.871A	621	715	830	87*	36*	38*	0.53	0.45	22	1.6	4
464.871B	623	715	829	66	41	43*	0.47*	0.33	22	1.6	4
464.871C	620	706*	826	87*	37*	41*	0.50*	0.39	23	1.7	4
464.889A	623	728	913	93*	36	39*	0.55*	0.42	22	1.5	3
464.889B	624	730*	920*	88*	40	36	0.53*	0.45*	23	1.5	3
464.889C	619	728	915	106*	47	42*	0.51*	0.42*	23	1.6	2
464.890A	625	725	913	90*	50	49*	0.49*	0.37*	22	1.7	2
464.890B	625	720	907	100*	41	46*	0.46*	0.34	25	1.9	2
464.891A	624	717	905	74*	39*	35*	0.42	0.40	22*	1.5	3
464.891B	620	726	913	105*	52	38*	0.55*	0.44	23	1.6	2
464.891C	621	725	912	92*	44	40	0.53	0.40*	24	1.6	2
464.892	626	719	906	85*	45*	63*	0.38*	0.31	23	1.3	2
464.925A	621	718	830*	100*	40	51*	0.53	0.33	23	1.4	3
464.925B	621	717	901	121*	40	52*	0.54	0.32	22	1.5	2
464.925C	622	720*	904	94*	44	39*	0.46	0.24	24	1.9	4
464.926	624	626	811*	34	44*	40	0.48	0.29	23	2.1	4

Table 4

Seed composition data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464

Entry	Maturity group	Protein (%)	Oil (%)	Palmitic (%)	Stearic (%)	Oleic (%)	Lino-leic (%)	Lino-lenic (%)	Other (%)
424.068	IV	50.8	9.4	14.4	3.1	16.3	51.2	15.2	0.1
424.071	IV	49.7	9.4	13.7	3.0	12.8	55.4	15.2	0.0
424.079	IV	49.6	9.0	14.3	3.1	11.9	56.1	14.8	0.0
424.088	IV	51.8	8.4	13.9	2.9	11.8	55.1	16.3	0.1
424.091A	IV	50.3	9.5	13.5	3.3	12.4	56.3	14.5	0.0
424.091B	IV	49.9	9.6	13.4	3.2	12.5	56.3	14.8	0.0
424.116	IV	48.2	10.0	13.3	3.2	12.8	53.9	17.0	0.0
440.913A	II	47.8	9.1	13.0	2.6	10.0	56.0	18.6	0.0
440.913B	II	47.9	10.2	12.8	2.6	10.2	56.2	18.2	0.1
447.003A	0	48.9	9.6	14.1	2.9	11.0	55.2	16.8	0.0
447.003B	I	47.7	11.4	13.3	3.0	12.0	55.0	16.9	0.0
447.004	III	46.9	10.1	13.4	3.1	12.2	54.3	17.1	0.0
458.535	0	48.3	9.7	13.5	3.5	16.9	53.6	12.8	0.0
458.536	0	51.4	9.1	13.7	3.2	11.3	53.4	18.4	0.0
458.537A	000	50.1	10.3	12.3	2.6	12.3	57.1	15.8	0.1
458.537B	000	47.9	11.4	12.4	2.6	12.2	57.1	15.8	0.1
458.538	000	47.8	10.5	12.4	2.6	12.9	56.6	15.6	0.1
458.539A	000	48.2	11.4	12.4	2.7	11.6	57.6	15.8	0.0
458.539B	0	48.4	10.2	13.4	3.2	12.6	55.5	15.4	0.0
458.540A	0	49.4	9.2	13.9	3.2	11.9	55.3	15.8	0.0
458.540B	0	49.1	9.4	13.9	3.2	12.1	55.5	15.3	0.0
458.540C	0	49.0	9.3	13.6	3.2	12.3	55.3	15.6	0.1
458.540D	0	48.8	9.3	13.9	3.2	11.1	55.5	16.5	0.0
464.866A	00	48.0	8.9	13.3	3.0	16.5	54.0	13.2	0.1
464.866B	00	47.3	9.0	13.4	3.3	15.0	55.2	13.3	0.0
464.867	000	47.1	10.0	12.6	2.7	10.9	57.6	16.3	0.1
464.868A	00	46.0	8.5	13.7	3.1	10.8	56.4	16.2	0.1
464.868B	00	47.7	10.1	12.5	2.9	11.5	56.8	16.4	0.0
464.869A	0	50.4	9.6	13.6	3.0	11.1	54.0	18.5	0.0
464.869B	0	51.1	9.4	13.1	2.8	10.5	54.5	19.3	0.0
464.870	000	47.2	9.9	12.8	2.6	11.2	57.4	16.1	0.1
464.871A	0	48.5	9.1	13.8	3.2	12.0	54.5	16.7	0.0
464.871B	0	48.7	9.1	13.9	3.2	11.4	54.8	16.8	0.1
464.871C	0	48.1	10.6	13.7	3.2	11.6	55.6	15.9	0.0
464.889A	II	48.6	9.2	12.3	2.7	11.0	57.2	16.9	0.0
464.889B	II	47.8	9.8	12.4	2.7	11.2	56.8	17.0	0.0
464.889C	II	48.4	9.5	12.9	2.8	10.8	54.9	18.6	0.0
464.890A	II	51.0	8.8	13.6	2.8	10.2	52.6	20.8	0.1
464.890B	I	48.7	10.8	13.3	3.0	11.9	54.1	18.0	0.1
464.891A	I	50.3	9.1	13.7	3.1	12.7	53.5	17.1	0.0
464.891B	II	49.8	8.7	13.6	3.0	11.4	53.2	18.9	0.0
464.891C	II	51.4	8.9	13.0	2.8	11.6	55.0	17.8	0.0
464.892	I	47.5	10.3	13.4	3.1	13.6	53.9	16.1	0.0
464.925A	0	49.8	9.9	13.2	2.9	11.3	54.0	18.9	0.0
464.925B	0	50.2	9.3	13.3	2.9	11.4	54.0	18.4	0.0
464.925C	I	50.6	10.5	13.0	3.4	13.3	55.1	15.2	0.1
464.926	000	47.1	12.2	12.7	2.7	14.6	56.3	13.7	0.0

Table 1

Identification and origin information for USDA wild soybean germplasm
in maturity groups 000 to IV, PI 65.549 to PI 483.464

PI No.	Accession name	Foreign or collector No.	Country of acquisition	Country of origin	Year introduced	Maturity group
464.927A		LS-005	China	China	1982	000
464.927B		LS-005	China	China	1982	000
464.927C		LS-005	China	China	1982	000
464.928		LS-008	China	China	1982	00
464.929A		LS-009	China	China	1982	0
464.929B		LS-009	China	China	1982	0
468.396A			China	China	1982	IV
468.396B			China	China	1982	IV
468.397A			China	China	1982	IV
468.397B			China	China	1982	IV
468.398A			China	China	1982	IV
468.398B			China	China	1982	IV
468.398C			China	China	1982	IV
468.399A			China	China	1982	IV
468.399B			China	China	1982	IV
468.399C			China	China	1982	IV
468.400A			China	China	1982	IV
468.400B			China	China	1982	IV
468.904			China	China	1982	00
468.905			China	China	1982	0
468.906			China	China	1982	0
468.907			China	China	1982	I
468.908			China	China	1982	000
468.909			China	China	1982	00
468.910			China	China	1982	00
468.911			China	China	1982	00
468.912			China	China	1982	00
468.913			China	China	1982	00
468.916			China	China	1982	III
468.917			China	China	1982	III
468.918			China	China	1982	III
468.919			China	China	1982	III
479.744	Gong di 50003-2	China	China	1983	I	
479.745	Gong di 50029-2	China	China	1983	I	
479.746A	Gong di 50062	China	China	1983	II	
479.746B	Gong di 50062	China	China	1983	II	
479.747	Gong di 50093-1	China	China	1983	III	
479.748	Gong di 50107-3	China	China	1983	II	
479.749	Gong di 50326-2	China	China	1983	III	
479.750	Gong di 50344	China	China	1983	I	
479.751	Gong di 50351-1	China	China	1983	III	
479.752	Gong di 50388-2	China	China	1983	I	
479.753A	Gong di 50396-1	China	China	1983	II	
479.753B	Gong di 50396-1	China	China	1983	II	
479.767	Long 79-1809	China	China	1983	I	
479.768	Long 79-3313-1	China	China	1983	0	
479.769	Long 79-5801	China	China	1983	0	

Table 2

Descriptive data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Matur-	Flower			Pubescence		Pod Density	Seed coat		Hilum color	Seed shape	Other traits		
		group	color	Color	Form	color		Luster	Color			Seed	Leaf	Plant
464.927A	000	P	T	Va	N	B1	B	B1	B1	Ob				
464.927B	000	P	T	Va	N	B1	Db	B1	B1	Ob				
464.927C	000	P	T	Va	N	B1	B	B1	B1	Ob				
464.928	00	P	T	Va	N	B1	Db	B1	B1	Ob				
464.929A	0	P	T	Sa	N	B1	B	B1	B1	Ov				
464.929B	0	P	T	Sa	N	B1	B	B1	B1	Ov				
468.396A	IV	P	T	Va	N	B1	B	B1	B1	Ov	Fleck			
468.396B	IV	P	T	A	N	B1	B	B1	B1	Ov	Fleck			
468.397A	IV	P	T	Va	N	B1	B	B1	B1	Ob				
468.397B	IV	P	T	Va	N	B1	B	B1	B1	Ob				
468.398A	IV	P	T	A	N	B1	B	B1	B1	Ov	Fleck			
468.398B	IV	P	T	A	N	B1	B	B1	B1	Ov	Fleck			
468.398C	IV	P	T	A	N	B1	B	B1	B1	Ov	Fleck			
468.399A	IV	P	T	A	N	B1	Db	B1	B1	Ov	Fleck			
468.399B	IV	P	T	Va	N	B1	Db	B1	B1	Ov	Fleck			
468.399C	IV	P	T	Va	N	B1	B	B1	B1	Ov	Fleck			
468.400A	IV	P	T	Va	N	B1	B	B1	B1	Ob	Fleck			
468.400B	IV	P	T	Va	N	B1	B	B1	B1	Ov	Fleck			
468.904	00	P	T	Va	N	Br	I	B1	B1	Ov		Na	Sw, 4sd	
468.905	0	P	T	A	N	Br	S	B1	B1	Ov		Na	Sw, 4sd	
468.906	0	P	T	Va	N	Br	S	B1	B1	Ov		Na	Sw	
468.907	I	W	T	Sa	N	Br	B	B1	B1	Nr	Fleck	Na	Sw, 4sd	
468.908	000	W	T	E	N	Br	Db	B1	B1	Nr		Na	Sw	
468.909	00	P	T	Sa	N	Br	I	B1	B1	Ov		Na	Sw	
468.910	00	P	T	E	N	Br	Db	B1	B1	Ov		Na	Sw, 4sd	
468.911	00	P	T	A	N	Br	I	B1	B1	Ov		Na	Sw	
468.912	00	P	T	Va	N	Br	Db	B1	B1	Nr		Na	Sw, 4sd	
468.913	00	P	Ng	A	N	Br	D	Br	Br	Ov			Sw	
468.916	III	P	T	Va	N	B1	B	B1	B1	Ov	Fleck			
468.917	III	P	T	A	N	B1	Db	B1	B1	Ob	Fleck			
468.918	III	P	T	Va	N	B1	B	B1	B1	Ov	Fleck			
468.919	III	P	T	Sa	N	B1	S	B1	B1	Ob	Fleck			
479.744	I	P	T	Sa	N	B1	B	B1	B1	Ov	Fleck			
479.745	I	P	T	Sa	N	B1	Db	B1	B1	Ob	Fleck			
479.746A	II	P	T	A	N	B1	Lb	B1	B1	Ov	Fleck			
479.746B	II	P	T	A	N	B1	B	B1	B1	Ov	Fleck			
479.747	III	P	T	Va	N	B1	B	B1	B1	Ob				
479.748	II	P	T	A	N	B1	B	B1	B1	Ov	Fleck			
479.749	III	P	T	Va	N	B1	Db	B1	B1	Ob	Fleck			
479.750	I	P	T	E	N	B1	Db	B1	B1	Ov	Fleck			
479.751	III	P	T	Va	N	B1	Db	B1	B1	Ob	Fleck			
479.752	I	P	T	Va	N	B1	B	B1	B1	Ov				
479.753A	II	P	T	A	N	B1	B	B1	B1	Ov				
479.753B	II	P	T	Sa	N	B1	Db	B1	B1	Ob				
479.767	I	P	T	Sa	N	B1	B	B1	B1	Ov				
479.768	0	P	T	Va	N	B1	B	B1	B1	Ov	Fleck			
479.769	0	P	T	Va	N	B1	Db	B1	B1	Ov				

Table 3

Agronomic data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Twining	Flower	Maturity	Leaflet Length				Leaflet shape		Pod	Seed
	date (mmdd)	date (mmdd)	date (mmdd)	Height (cm)	lower (mm)	upper (mm)	lower (mm)	upper (mm)	length (mm)	weight (cg/sd)	SMV
464.927A	625	627	812*	46*	42	41*	0.44	0.31	25	1.9*	3
464.927B	623	625	811*	50	55	60*	0.39	0.31	24	1.8*	4
464.927C	623	628	812	61*	48*	43*	0.38	0.33	26*	1.6*	3
464.928	625	628	817	78*	39	39*	0.47*	0.44*	24	1.6	4
464.929A	623	716	830*	70*	41	43*	0.54*	0.42*	22	1.6	4
464.929B	623	715	826	64*	33	36*	0.57	0.38	22	1.7	5
468.396A	702	817	1004	68	40	35	0.47	0.29	23	1.5	4
468.396B	702	817	1004	77	41	38	0.51*	0.27	24	1.6	4
468.397A	628	822*	1004	131*	51	42	0.50*	0.34	27	1.4	3
468.397B	627	822*	1004	93*	40	35	0.45*	0.30	25	1.4*	3
468.398A	Agronomic data not available										
468.398B	Agronomic data not available										
468.398C	Agronomic data not available										
468.399A	703	820	1006	101*	40	36	0.49*	0.35	26	1.7	2
468.399B	705	819	1001	89	40	36	0.50	0.35	25	1.1	1
468.399C	707*	820	1002	96	40	35*	0.58*	0.33	25	1.2	1
468.400A	629	816	1001	101*	36	32	0.58*	0.30	25	1.5	3
468.400B	704	816	1004	128*	46	33	0.52	0.31	24	1.5	1
468.904	617	701	823*	104*	56	66*	0.43	0.38	30*	3.5*	1
468.905	618	703	825*	93*	47*	42*	0.30	0.29	28*	3.1*	2
468.906	620	706*	829*	69	49	48*	0.41	0.36	30	3.7*	4
468.907	628	712	903	110*	98	67*	0.39	0.29	35	4.1	2
468.908	618	622	811*	78	60*	58*	0.41	0.32	31	3.7*	2
468.909	618	704*	823*	105*	49	53*	0.32	0.32	32	3.1	2
468.910	618	626	822*	86	54	65*	0.37	0.30	32	4.3*	1
468.911	618	629	817*	93*	46	60*	0.41	0.28	31*	3.5*	1
468.912	617	623	815*	83*	45	55*	0.42*	0.32	31	3.6*	3
468.913	618	626	819*	87	48	50*	0.46	0.34	33	3.3*	3
468.916	628	806	920	93	39	37	0.55	0.33	20	1.3	3
468.917	624	807	921	95*	50	39	0.55*	0.47	22	2.0	4
468.918	625	803	920	71	45	38	0.45	0.31	20	1.3	3
468.919	626	728	918	152	66	64*	0.46*	0.31	28*	2.9	4
479.744	625	716	831	63	31	19	0.45	0.47	19	1.2	3
479.745	625	716	831	47	26	24	0.42	0.38	19	2.3	4
479.746A	625	720	908	58	39	31	0.36	0.35	24	1.4	3
479.746B	625	720	908	58	39	31	0.36	0.35	24	1.5	3
479.747	627	725	917	59	43	30	0.42	0.40	24	1.5	4
479.748	702	725	908	61	37	31	0.32	0.35	23	1.0	4
479.749	620	725	920	92	48	32	0.35	0.25	25	1.9	4
479.750	620	713	827	42	31	18	0.42	0.39	24	1.3	4
479.751	709	808	925	43	37	34	0.59	0.41	22	1.8	4
479.752	625	713	902	66	53	28	0.32	0.39	23	1.1	3
479.753A	702	725	915	61	46	32	0.33	0.25	23	1.8	4
479.753B	702	725	915	61	46	32	0.33	0.25	23	1.8	3
479.767	620	716	827	46	28	24	0.54	0.42	26	1.5	3
479.768	620	701	815	45	37	23	0.35	0.30	23	1.5	3
479.769	622	716	819	45	31	24	0.45	0.38	21	1.7	3

Table 4

Seed composition data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464

Entry	Maturity group	Protein (%)	Oil (%)	Palmitic (%)	Stearic (%)	Oleic (%)	Linoleic (%)	Linenic (%)	Other (%)
464.927A	000	48.8	10.9	12.8	2.9	12.6	56.8	15.0	0.0
464.927B	000	47.6	10.4	12.6	2.6	11.6	57.1	16.2	0.1
464.927C	000	49.6	10.1	14.4	3.6	14.3	53.4	14.4	0.1
464.928	00	47.9	9.8	13.4	3.2	11.1	58.6	15.4	0.1
464.929A	0	48.1	9.7	13.8	3.1	11.4	55.1	16.7	0.2
464.929B	0	45.1	7.3	14.0	3.0	12.2	55.3	15.5	0.0
468.396A	IV	44.8	12.3	13.6	3.4	13.9	54.9	14.3	0.1
468.396B	IV	45.1	12.1	13.6	3.3	14.1	55.0	14.1	0.0
468.397A	IV	47.9	10.2	13.5	2.8	12.5	55.0	16.3	0.0
468.397B	IV	47.7	10.5	13.5	2.8	12.9	55.1	15.8	0.1
468.398A	IV	Seed composition data not available							
468.398B	IV	Seed composition data not available							
468.398C	IV	Seed composition data not available							
468.399A	IV	48.1	9.3	14.0	2.9	11.0	54.7	17.4	0.0
468.399B	IV	45.5	10.1	13.1	3.3	11.7	53.8	18.3	0.0
468.399C	IV	46.2	9.2	13.1	3.3	11.3	54.2	18.1	0.0
468.400A	IV	46.8	10.6	13.7	3.7	13.4	55.2	14.2	0.1
468.400B	IV	49.1	10.1	13.7	3.0	11.2	55.3	16.8	0.1
468.904	00	48.0	12.0	12.6	2.6	15.3	55.1	14.4	0.1
468.905	0	44.3	12.7	13.6	2.9	14.3	55.8	13.4	0.1
468.906	0	45.0	13.4	13.4	2.8	16.7	54.7	12.5	0.1
468.907	I	46.6	13.9	13.1	3.2	11.9	55.4	16.3	0.1
468.908	000	45.1	13.9	11.8	3.1	15.5	56.7	13.0	0.1
468.909	00	47.4	12.1	12.7	2.7	14.3	55.5	14.8	0.1
468.910	00	46.0	13.6	11.2	3.1	25.7	49.5	10.7	0.1
468.911	00	47.9	11.8	12.7	2.7	16.5	53.6	14.4	0.1
468.912	00	46.8	14.4	12.2	3.2	16.5	57.2	11.0	0.1
468.913	00	45.4	13.2	13.9	2.4	16.9	55.3	11.6	0.1
468.916	III	47.8	9.0	14.0	3.0	12.4	54.6	17.1	0.0
468.917	III	49.4	10.0	13.9	2.9	12.4	54.9	16.0	0.0
468.918	III	47.2	10.0	12.8	2.8	15.9	53.1	15.4	0.1
468.919	III	46.8	13.5	14.6	2.9	14.6	55.7	12.4	0.0
479.744	I	49.1	8.6	12.7	2.7	12.3	54.3	17.9	0.0
479.745	I	50.7	10.6	12.6	3.1	8.9	56.2	19.1	0.0
479.746A	II	49.7	8.8	12.7	3.3	14.5	53.6	15.9	0.0
479.746B	II	50.7	9.3	14.5	3.3	12.2	53.3	16.7	0.0
479.747	III	51.6	8.7	13.6	3.1	11.8	53.7	17.8	0.0
479.748	II	47.3	9.6	13.2	3.1	14.2	52.0	17.5	0.0
479.749	III	49.9	9.8	13.0	2.9	10.9	56.4	16.9	0.0
479.750	I	47.7	9.8	12.6	2.8	8.7	53.3	22.6	0.0
479.751	III	49.0	9.0	14.5	2.7	11.1	56.1	15.5	0.1
479.752	I	49.9	7.5	12.8	2.8	10.5	53.5	20.4	0.0
479.753A	II	50.6	9.8	13.5	3.1	12.4	54.3	16.8	0.0
479.753B	II	50.6	8.5	13.2	3.2	11.7	54.5	17.3	0.0
479.767	I	50.6	8.6	13.2	3.0	15.8	55.6	12.4	0.0
479.768	0	49.5	9.7	12.8	3.2	9.9	55.3	18.7	0.1
479.769	0	47.8	7.2	13.4	3.0	10.6	55.1	17.9	0.0

Table 1

Identification and origin information for USDA wild soybean germplasm
in maturity groups 000 to IV, PI 65.549 to PI 483.464

PI No.	Accession name	Foreign or collector No.	Country of acquisition No.	Country of origin	Year introduced	Matur- ity group
483.071A			China	China	1983	IV
483.071B			China	China	1983	IV
483.459			China	China	1984	I
483.460A			China	China	1984	III
483.460B			China	China	1984	III
483.460C			China	China	1984	IV
483.461			China	China	1984	II
483.462A			China	China	1984	IV
483.462B			China	China	1984	IV
483.463			China	China	1984	III
483.464A			China	China	1984	III
483.464B			China	China	1984	III

Table 2

Descriptive data for USDA wild soybean germplasm in maturity groups
000 to IV, PI 65.549 to PI 483.464

Entry	Matur- ity group	Flower		Pubescence		Pod Density	Seed coat		Hilum color	Seed shape	Other traits		
		color	Color	Form	Sp		Luster	Color			Seed	Leaf	Plant
483.071A	IV	P	T	Sa	N	B1	B	B1	B1	Ob			
483.071B	IV	P	T	Sa	N	B1	B	B1	B1	Ob	Fleck		
483.459	I	P	T	Sa	Sp	Br	I	B1	B1	Ov	Fleck		Sw
483.460A	III	P	T	A	N	B1	B	B1	B1	Ov	Fleck		
483.460B	III	P	T	A	N	B1	B	B1	B1	Nr	Fleck		
483.460C	IV	P	T	A	N	B1	B	B1	B1	Ov	Fleck		
483.461	II	P	T	A	N	B1	B	B1	B1	Ov	Fleck		
483.462A	IV	P	T	A	N	B1	B	B1	B1	Ov			
483.462B	IV	P	T	A	N	B1	B	B1	B1	Ob			
483.463	III	P	T	Va	N	B1	Db	B1	B1	Ov	Fleck		
483.464A	III	P	T	Va	N	B1	B	B1	B1	Ob	Fleck		
483.464B	III	P	T	Va	N	B1	B	B1	B1	Ob	Fleck		

Table 3

Agronomic data for USDA wild soybean germplasm in maturity groups

000 to IV, PI 65.549 to PI 483.464

Entry	Twining	Flower	Maturity	Leaflet length				Pod		Seed
	date (mmdd)	date (mmdd)	date (mmdd)	Height (cm)	lower (mm)	upper (mm)	Leaflet shape	length (mm)	weight (cg/sd)	SMV (score)
483.071A	629	813	1006	76	32	28	0.38	0.29	23	1.3 3
483.071B	629	813	1006	76	32	28	0.38	0.29	23	1.3 3
483.459	623	713	903	58	51	47	0.71	0.47	33	5.9 2
483.460A	618	806	925	103	56	40	0.34	0.28	26	1.3 3
483.460B	618	806	925	103	56	40	0.34	0.28	26	1.0 2
483.460C	618	806	925	103	56	40	0.34	0.28	26	1.3 2
483.461	629	716	905	48	27	23	0.67	0.30	22	1.2 4
483.462A	622	808	927	72	54	37	0.35	0.30	24	1.6 3
483.462B	622	808	927	72	54	37	0.35	0.30	24	1.9 4
483.463	704	803	922	59	31	26	0.39	0.27	20	1.2 4
483.464A	627	725	1002	48	28	26	0.43	0.35	21	1.6 4
483.464B	627	725	1002	48	28	26	0.43	0.35	21	2.0 5

Table 4

Seed composition data for USDA wild soybean germplasm in maturity groups 000 to IV, PI 65.549 to PI 483.464

Entry	Matur- ity group	Protein	Oil	Pal- mitic	Stearic	Oleic	Lino- leic	Lino- lenic	Other
		(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)
483.071A	IV	44.2	12.0	12.9	2.8	14.0	55.5	14.8	0.0
483.071B	IV	42.1	13.3	12.8	3.7	14.9	54.6	13.9	0.1
483.459	I	44.7	14.2	14.6	2.8	9.0	54.8	18.7	0.0
483.460A	III	47.6	10.4	13.1	3.1	11.8	55.8	16.1	0.0
483.460B	III	47.7	7.8	13.5	2.9	10.1	52.5	21.0	0.0
483.460C	IV	47.0	10.6	12.7	3.2	12.7	55.2	16.3	0.0
483.461	II	49.5	9.0	12.5	2.9	12.1	55.0	17.5	0.0
483.462A	IV	44.6	11.8	13.3	3.4	14.8	53.2	15.3	0.1
483.462B	IV	46.6	10.8	13.4	3.4	14.3	53.7	15.2	0.0
483.463	III	46.1	10.5	12.6	2.9	12.2	56.1	16.1	0.0
483.464A	III	47.3	11.6	14.8	3.2	15.5	53.4	13.0	0.0
483.464B	III	48.5	11.0	15.1	3.8	18.3	50.9	11.8	0.0