



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

blandaltman: A command to create variants of Bland–Altman plots

Mark D. Chatfield
Faculty of Medicine
The University of Queensland
Brisbane, Australia
m.chatfield@uq.edu.au

Tim J. Cole
Great Ormond Street Institute of Child Health
University College London
London, U.K.

Henrica C. W. de Vet
Department of Epidemiology and Data Science
Amsterdam Medical Center
Amsterdam, The Netherlands

Louise Marquart-Wilson
School of Public Health
The University of Queensland
Brisbane, Australia

Daniel M. Farewell
School of Medicine
Cardiff University
Cardiff, U.K.

Abstract. Bland–Altman plots can be useful in paired data settings such as measurement-method comparison studies. A Bland–Altman plot has differences, percentage differences, or ratios on the y axis and a mean of the data pairs on the x axis, with 95% limits of agreement indicating the central 95% range of differences, percentage differences, or ratios. This range can vary with the mean. We introduce the community-contributed `blandaltman` command, which uniquely in Stata can 1) create Bland–Altman plots featuring ratios in addition to differences and percentage differences, 2) allow the limits of agreement for ratios and percentage differences to vary as a function of the mean, and 3) add confidence intervals, prediction intervals, and tolerance intervals to the plots.

Keywords: gr0094, `blandaltman`, Bland–Altman plot, limits of agreement, agreement, `baplot`, `batplot`, concord, prediction, tolerance, interval, ratio, percentage difference

1 Introduction

When paired data arise from two different measurement techniques, for example, a new method A and a conventional method B, the data can be plotted as in figure 1a to visualize the correlation between the two methods. However, this plot is not the best for clearly showing the differences between the methods (Bland and Altman 1986). Bland and Altman (1986, 1999) introduced a plot for visualizing agreement that plots the difference between data pairs versus their arithmetic mean (figure 1b). This is known as the Bland–Altman plot, and it can be used in other paired data settings such as measurement repeatability (Bland and Altman 1986, 1999) or longitudinal studies

(Kirkwood and Sterne 2003). Variants of the Bland–Altman plot have ratios or percentage differences on the y axis and may have the geometric mean of data pairs on the x axis (Dewitte et al. 2002). Note that in geometric terms, the Bland–Altman plot rotates figure 1a clockwise by 45° and linearly rescales the axes. Figure 1b shows the data inside the gray box on figure 1a.

According to Bland and Altman (1999), 95% limits of agreement (LOA) provide an interval within which 95% of differences between measurements are expected to lie. If these limits are not too large (this is a contextual consideration in light of the intended use of the measurement method), then the methods can be considered interchangeable. Assuming differences are normally distributed, the LOA can be calculated using the mean and standard deviation (SD) of the paired differences (as $\text{mean} \pm 1.96 \text{ SD}$), and they are routinely added to a Bland–Altman plot as a pair of horizontal lines toward the top and bottom of the data cloud.

However, horizontal LOA are “meaningful only if we can assume the bias [*the mean difference*] and variability [*the SD of the difference*] are uniform throughout the range of measurement, assumptions which can be checked graphically” (Bland and Altman 1999). In figure 1b, the mean difference changes little as the mean of data pairs varies, but the SD of the difference increases steeply with the mean of data pairs, so the data cloud is shaped like a left-pointing arrowhead. Bland and Altman (1999) suggest that this arrowhead pattern is the most common departure from the assumptions underlying horizontal LOA. In this instance, the LOA need to reflect the varying SD. The plot shows LOA calculated assuming that the SD increases linearly with the mean of data pairs, using the regression-based approach of Bland and Altman (1999) to adjust for nonconstant means or SDs of differences.




Figure 1. (a) Plot of two methods for measuring retinol-binding-protein-4 ($\mu\text{mol/L}$) from Brindle et al. (2017), with the line of equality. (b) Bland–Altman plot featuring differences with regression-based estimates of 95% LOA (thin gray solid lines) and mean difference or “bias” (dashed line). The boxes show how rotating (a) clockwise by 45° and rescaling the axes leads to (b).

Log transformation often leads to the mean and SD of differences being constant, which in turn justifies using horizontal LOA. Figure 2a shows the same data as figure 1a but with the axes scaled logarithmically. Figure 2b is the corresponding Bland–Altman plot, where ratios (A/B) are plotted on the y axis and the geometric mean of the data pairs is plotted on the x axis. Both axes are scaled logarithmically. Plotted this way, the rotational symmetry between the plot of raw data and the Bland–Altman plot is preserved.

Bland–Altman plots can also have percentage differences on the y axis, where the percentage difference is defined by dividing a difference by the arithmetic mean of the data pairs and multiplying by 100%. These percentage differences (which can range from -200% to $+200\%$) are often plotted against the arithmetic mean of the data pairs (Dewitte et al. 2002). Other ways of defining percentage differences are possible (Cole and Altman 2017). Dividing a difference by the logarithmic mean of the data pairs was recommended by economists (Törnqvist, Vartia, and Vartia 1985), and multiplying by 100% produces a percentage difference that can be calculated simply as $100(\ln A - \ln B)$ (Cole 2000). These percentage differences (which can range from $-\infty$ to $+\infty$) could be plotted instead of ratios on the y axis of figure 2b, and rotational symmetry could thus be preserved.




Figure 2. (a) Log-log plot of two methods for measuring retinol-binding-protein-4 ($\mu\text{mol/L}$) from Brindle et al. (2017) with the line of equality. (b) Bland–Altman plot featuring ratios with estimates of horizontal 95% LOA (thin gray solid lines) and the geometric mean of ratios (dashed line). The boxes show how rotating (a) clockwise by 45° and rescaling the axes leads to (b).

Stata has no official Bland–Altman plot command, but there are several community-contributed commands. None of them create Bland–Altman plots with ratios, and only the `agree` command (Doménech and Sesma 2021) gives percentage differences. For differences, `batplot` (Mander 2005) and `biasplot` (Taffé et al. 2017) can draw regression-based LOA for datasets with one measurement per method per subject and for those with several measurements for the reference method per subject. The commands `concord` (Steichen and Cox 1998), `baplot` (Seed 2000), `kappaetc` (Klein 2018), `agree` (Doménech and Sesma 2021), and `rmloa` (Linden 2021) present only horizontal LOA.

Bland and Altman (1986, 1999) recommended calculating 95% confidence intervals for LOA. Yet surprisingly, none of the commands cited above displays confidence intervals on the plot.¹ Some authors have recommended prediction and tolerance intervals (TIs) (Ludbrook 2010; Vock 2016; Carkeet and Goh 2018; Francq, Berger, and Boachie 2020).

In this article, we introduce the `blandaltman` command, which uniquely in Stata can 1) create Bland–Altman plots featuring ratios in addition to differences and percentage differences, 2) allow LOA for ratios and percentage differences to vary with the mean of data pairs, and 3) add confidence intervals, prediction intervals (PIs), and TIs to the plots. We show how, by offering variants of Bland–Altman plots, the command can help decide how best to present LOA for measurement-method comparison studies based on how close to horizontal the regression-based LOA are.

1. Confidence intervals are reported in the nongraphical output using `agree`.

The rest of the article is organized as follows: Section 2 presents the `blandaltman` command, syntax, and options; section 3 shows the command in action; and section 4 concludes.

2 The `blandaltman` command

2.1 Description

`blandaltman` produces Bland–Altman plots featuring differences, ratios, or percentage differences on the y axis. By default, regression-based estimates of bias and LOA appear on the plot to show how the distribution of differences, ratios, or percentage differences varies with the mean of data pairs. See the appendix for details. Horizontal lines for bias and LOA can be produced instead.

Summary statistics are provided in the output. The distribution of differences and percentage differences are summarized by mean and SD, and these are used to calculate horizontal LOA. The distribution of ratios is summarized by geometric mean (GMean) and geometric SD (Limpert and Stahel 2011). These can be calculated by antilogging the mean and SD of differences in log-transformed data. For ratios, all calculations are done using log-transformed data before results such as LOA are antilogged.

Confidence intervals for the bias and LOA can also be displayed, as well as a PI and (up to three) TIs (see appendix for details), assuming the distribution of differences, ratios, or percentage differences does not vary with the mean of data pairs.

2.2 Syntax

```
blandaltman varA varB [if] [in], plot(plot_type) [horizontal noregloa
  noregbias hloa hbias level(#) predinterval ticonfidence(#)
  ticonfidence2(#) ticonfidence3(#) cibias ciloa cilevel(#)
  scopts(scatter_options) regloaopts(tw_function_options)
  regbiasopts(tw_function_options) loaopts(tw_function_options)
  biasopts(tw_function_options) piopts(tw_function_options)
  tiopts(tw_function_options) tiopts2(tw_function_options)
  tiopts3(tw_function_options) ciloaopts(tw_pcarrowi_options)
  cibiasopts(tw_pcarrowi_options) addplot(plot ... [|| plot ... [...]])
  twoway_options]
```

2.3 Options

`plot(plot_type)` creates the plot and is required. *plot_type* is any combination of the following:

<i>plot_type</i>	<i>y</i> axis	<i>x</i> axis
difference	$A - B$	$(A + B)/2$
ratio	A/B^*	$\text{GMean}(A, B)^*$
percentlmean	$100(A - B)/\text{LMean}(A, B) = 100(\ln A - \ln B)$	$\text{GMean}(A, B)^*$
percentmean	$100(A - B)/\{(A + B)/2\}$	$(A + B)/2^*$

* Axis has a logarithmic scale.

Multiple plots are created if several plot types are chosen.

$\text{GMean}(A, B) = (A \times B)^{1/2}$ is the geometric mean.

$\text{LMean}(A, B) = (A - B)/(\ln A - \ln B)$ if $A \neq B$, and $\text{LMean}(A, B) = A$ if $A = B$ is the logarithmic mean (Cole 2000).

Only positive-valued data are used with the options **ratio** and **percentlmean**. With the exception of $A = B = 0$, data pairs where $A \geq 0$ and $B \geq 0$ are used with the **percentmean** option.

horizontal displays *horizontal* rather than *regression-based* LOA and bias. This is equivalent to specifying **noregloa** **noregbias** **hloa** **hbias**.

noregloa prevents display of regression-based LOA.

noregbias prevents display of regression-based bias and LOA.

hloa displays horizontal LOA.

hbias displays horizontal bias. This option is assumed whenever horizontal LOA or a PI or a TI is requested.

level(#) specifies the level, as a percentage, for *##%* LOA, *##%* PI, and *##%* TI with a percent confidence as specified in **ticonfidence()**. The default is **level(95)**.

predinterval displays (horizontal) lines for a level percent PI.

ticonfidence(#) displays (horizontal) lines for a level percent TI with *##%* confidence.

ticonfidence2(#) displays (horizontal) lines for a second level percent TI with *##%* confidence.

ticonfidence3(#) displays (horizontal) lines for a third level percent TI with *##%* confidence.

cibias displays a percent confidence interval, as specified in **cilevel()**, for horizontal bias. This option requires that **horizontal** or **hbias** also be specified.

`ciloa` displays (exact) percent confidence intervals, as specified in `cilevel()`, for horizontal LOA. This option requires that `horizontal` or `hloa` also be specified.

`cilevel(%)` specifies the level, as a percentage, for confidence intervals for the bias and LOA. The default is `cilevel(95)`.

`scopts(scatter_options)` alters the display of the scatterplot. *scatter_options* are any of the options allowed with `scatter`; see [G-2] **graph twoway scatter**.

tw_function_options are any of the options allowed with `twoway function`; see [G-2] **graph twoway function**.

`regloaopts(tw_function_options)` alters the display of the regression-based LOA.

`regbiasopts(tw_function_options)` alters the display of the regression-based bias.

`loaopts(tw_function_options)` alters the display of the horizontal LOA.

`biasopts(tw_function_options)` alters the display of the horizontal bias line.

`plopts(tw_function_options)` alters the display of the PI.

`tiopts(tw_function_options)` alters the display of the first TI.

`tiopts2(tw_function_options)` alters the display of the second TI.

`tiopts3(tw_function_options)` alters the display of the third TI.

tw_pcarrowi_options are any of the options allowed with `twoway pcarrowi`; see [G-2] **graph twoway pcarrowi**.

`ciloaopts(tw_pcarrowi_options)` alters the display of the confidence interval for the LOA.

`cibiasopts(tw_pcarrowi_options)` alters the display of the confidence interval for the bias.

`addplot(plot ... [|| plot ... [...]])` adds other plots to the Bland–Altman plot; see [G-3] **addplot_option**.

twoway_options are any of the options documented in [G-3] **twoway_options**.

3 Examples

This section illustrates `blandaltman` in action. The first example shows how the estimated LOA on a Bland–Altman plot vary throughout the range of measurement. The second example shows how to add confidence intervals, a PI, and TIs to a plot.

3.1 Laboratory measurements: Exploring how LOA vary throughout the range of measurement

Brindle et al. (2017) described the simultaneous assessment of seven micronutrient and inflammation status biomarkers via a multiplex immunoassay method in a population of pregnant women. Results from their seven-plex assay were compared with conventional immunoassay results on $N = 206$ plasma samples. We focus on retinol-binding-protein-4 (figure 1a), a surrogate biomarker for vitamin A deficiency, where low levels indicate deficiency. For simplicity, we generate variables named A and B to represent measurements obtained using the new and conventional methods, respectively.

```
. use labmeasures
(Brindle et al. 2017 PLoS ONE 12(10): e0185868; doi: 10.1371/journal.pone.0185868)
. generate A = plexrbp4μmol1
. generate B = nimanurbp4μmol1
. blandaltman A B, plot(difference ratio percentlmean percentmean)
(see output in appendix)
```

The above line of syntax produces the four Bland–Altman plots shown in figure 3. As seen in figure 3a, the estimated LOA for differences are far from horizontal. In contrast, the estimated LOA for ratios (figure 3b) and percentage differences (figures 3c and 3d) are close to horizontal, so horizontal LOA would be justified. Note that figures 3b and 3c are equivalent (they differ only in their y -axis labeling). Figure 3d looks similar to figure 3c but plots a different definition of percentage difference against a different mean.




Figure 3. Bland–Altman plots featuring (a) differences, (b) ratios, (c) percentage differences using the logarithmic mean as denominator, and (d) percentage differences using the arithmetic mean as denominator. Each plot shows regression-based estimates of LOA (gray solid lines) and bias (dashed line). (b) and (c) are equivalent plots, while (d) is different.

The values for horizontal LOA are displayed when the option `horizontal` is specified:

```
. blandaltman A B, plot(ratio percentlmean percentmean) horizontal
```

A: A
B: B

PERCENTAGE DIFFERENCES (using Mean as denominator)...

Calculation	N	Mean	SD	Interval(s)
100*(A-B)/[(A+B)/2]	206	-10.07663	17.10901	
95% limits of agreement: -43.60967 23.45642				

PERCENTAGE DIFFERENCES (using Logarithmic Mean as denominator)...

Calculation	N	Mean	SD	Interval(s)
100*(A-B)/LMean(A,B)	206	-10.16367	17.29902	
95% limits of agreement: -44.06912 23.74177				

RATIOS...

Calculation	N	GMean	GSD	Interval(s)
A/B	206	.9033576	1.188854	
95% limits of agreement: .6435914 1.267971				

LOA from both definitions of percentage difference will often be very similar, as is the case here. Assuming that both percentage differences are approximately normally distributed and ratios are approximately lognormally distributed, there is little to choose between the three LOA above. Convention or personal preference may be the deciding factor in selecting one.

The previous code produces three Bland–Altman plots with horizontal LOA, one of which is shown in figure 4. It features percentage differences (using arithmetic mean as the denominator) on the y axis and arithmetic mean on the x axis, which is a popular choice in bioanalytical method validation studies (Dewitte et al. 2002). By default, the x axis is scaled logarithmically, which helps to space the data out more evenly. However, if users want a linear scale instead, they can specify the option `xscale(nolog)`. Assuming that these percentage differences are approximately normally distributed, LOA are estimated to be -44% and $+23\%$.




Figure 4. Bland–Altman plot featuring percentage differences (using arithmetic mean as denominator) assuming horizontal 95% LOA (gray solid lines) and bias (dashed line)

For control over the labeling of axes, the `xlabel()` and `ylabel()` options can be specified. For example, in our plots featuring ratios (figures 2b and 3b), we specified `ylabel(0.6 (0.2) 1.6)`. See Cox (2018, 2020) for other ways of labeling log-scaled axes.

3.2 Peak expiratory flow rate data: Adding confidence intervals, PI, and TIs

To demonstrate their method, Bland and Altman (1986) measured peak expiratory flow rate in 17 persons using both a Wright flow meter and a mini-Wright flow meter. Like them and others (Ludbrook 2010; Carkeet 2015; Vock 2016), we use the first measurement by each method. Figure 5 shows the data on an (overly busy) Bland–Altman plot. Bland and Altman saw no obvious relation between differences and means and assumed differences were normally distributed. They estimated² 95% LOA to be $-2.1 \pm 2 \times 38.8$ l/min.

Figure 5 illustrates the various intervals that `blandaltman` can produce—see the appendix and the references in this section for the meaning of prediction and TIs. The figure was created with the following syntax:

```
. use pefr, clear
(Bland and Altman (1986) Lancet 327: 307–10.)
. blandaltman Wright Mini, plot(difference) horizontal
> ciloa cibias predinterval ticonfidence(95)
> loaopts(lc(gs11) lp(solid)) cilaopts(mc(gs11) lc(gs11) lp(solid))
> piopts(lc(gs1) lp(dash_dot))
> tiopts(lc(gs1) lp(dot))
> legend(on order(2 "Bias (& 95% CI)"
> 4 "95% limits of agreement (& exact 95% CI)"
> 6 "95% prediction interval"
> 8 "95% tolerance interval with 95% confidence") rowgap(*.7) cols(1))
```

```
A: Wright           Wright peak expiratory flow rate (l/min)
B: Mini             Mini Wright peak expiratory flow rate (l/min)
```

DIFFERENCES...

Calculation	N	Mean	SD	Interval(s)
A-B	17	-2.117647	38.76513	
		95% limits of agreement:	-78.09591	73.86061
		95% prediction interval:	-86.67853	82.44323
		95% tolerance interval with 95% confidence:	-113.4634	109.2281
		95% CI (LLOA):	-124.1608	-53.09493
		95% CI (ULOA):	48.85964	119.9255
		95% CI (Mean diff.):	-22.04884	17.81354

2. Factors 2 and 1.96, respectively, are used in their 1986 and 1999 articles.




Figure 5. Bland–Altman plot of two methods measuring peak expiratory flow rate (l/min), illustrating the variety of intervals that **blandaltman** can produce. No author has suggested using all of these intervals.

We now report which of these intervals different authors have recommended and how they can be implemented with **blandaltman**. Bland and Altman (1999) recommended estimates of 95% LOA and 95% confidence intervals for LOA, and this advice features in reporting standards (Gerke 2020). Royston and Matthews (1991) considered methods to provide a best estimate of an interval containing the central 95% of a distribution. They considered the interval bounded by mean ± 1.96 SD (that is, LOA) to be a good estimate, and they viewed a 95%-expectation TI (equivalent to a 95% PI) to be of questionable value, as they did a 95% TI with $\geq 90\%$ confidence.

A few authors prefer not to present estimates of LOA (Ludbrook 2010; Vock 2016; Carkeet and Goh 2018; Francq, Berger, and Boachie 2020). Some prefer a 95% PI instead (Ludbrook 2010; Francq, Berger, and Boachie 2020), but not Vock (2016), who argued this is rarely appropriate. Others prefer a 95% TI with 50% confidence (Carkeet and Goh 2018).

Vock (2016) encouraged reporting a 95% TI with 95% confidence, as did Ludbrook (2010).³ Francq, Berger, and Boachie (2020) thought this interval may be too large (see table A1 for how intervals depend on sample size) and suggested a 95% TI with 80% or 90% confidence might be presented if needed. Carkeet and Goh (2018) recommended reporting a 95% TI with 2.5% confidence and a 95% TI with 97.5% confidence.

The following syntax creates Bland–Altman plots with intervals recommended by the above-mentioned authors:

```
. blandaltman Wright Mini, plot(difference) name(Bland_Altman, replace)
> horizontal ciloa
> legend(on order(2 "Bias" 4 "95% limits of agreement (& exact 95% CI)")
>       cols(1))
(output omitted)

. blandaltman Wright Mini, plot(difference) name(Ludbrook, replace)
> noreg hbias predinterval ticonfidence(95)
> legend(on order(2 "Bias" 4 "95% prediction interval"
>       6 "95% tolerance interval with 95% confidence") cols(1))
(output omitted)

. blandaltman Wright Mini, plot(difference) name(Francq_et_al, replace)
> noreg hbias predinterval ticonfidence(80)
> legend(on order(2 "Bias" 4 "95% prediction interval"
>       6 "95% tolerance interval with 80% confidence") cols(1))
(output omitted)

. blandaltman Wright Mini, plot(difference) name(Vock, replace)
> noreg hbias ticonfidence(95)
> legend(on order(2 "Bias" 4 "95% tolerance interval with 95% confidence")
>       cols(1))
(output omitted)

. blandaltman Wright Mini, plot(difference) name(Carkeet_Goh, replace)
> noreg hbias ticonfidence(2.5) ticonfidence2(50) ticonfidence3(97.5)
> legend(on order(2 "Bias" 4 "95% tolerance interval with 2.5% confidence"
>       6 "95% tolerance interval with 50% confidence"
>       8 "95% tolerance interval with 97.5% confidence") cols(1))
(output omitted)
```

4 Conclusion

The `blandaltman` command should help Stata users when assessing agreement in measurement-method comparison studies to follow the advice of Bland and Altman (1999) by visually assessing how estimated LOA vary throughout the range of measurement and by reporting corresponding confidence intervals. The command is also flexible enough to allow users to follow recommendations of other authors involving the presentation of a PI or a TI. More generally, in other paired data settings, the command could help users decide whether to summarize differences, ratios, or percentage differences defined in one of two ways.

3. Vock (2016) also considered an interval formed by the outer confidence limits for the LOA as an alternative.

5 Acknowledgment

`blandaltman` uses some syntax from the command `niceloglabels` (Cox 2018, 2020) to provide publication-ready plots when axes are scaled logarithmically. We thank the editor and a referee for help in structuring the article and Professor Annette Dobson for helpful feedback.

6 Programs and supplemental materials

To install the software files as they existed at the time of the publication of this article, type

```
. net sj 23-3
. net install gr0094      (to install program files, if available)
. net get gr0094          (to install ancillary files, if available)
```

To install the latest version of software files, type

```
. ssc install blandaltman (to install program files)
. net get blandaltman     (to install ancillary files)
```

7 References

- Bland, J. M., and D. G. Altman. 1986. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 327: 307–310. [https://doi.org/10.1016/S0140-6736\(86\)90837-8](https://doi.org/10.1016/S0140-6736(86)90837-8).
- . 1999. Measuring agreement in method comparison studies. *Statistical Methods in Medical Research* 8: 135–160. <https://doi.org/10.1177/096228029900800204>.
- Brindle, E., L. Lillis, R. Barney, S. Y. Hess, K. R. Wessells, C. T. Ouédraogo, S. Stinca, et al. 2017. Simultaneous assessment of iodine, iron, vitamin A, malarial antigenemia, and inflammation status biomarkers via a multiplex immunoassay method on a population of pregnant women from Niger. *PLOS ONE* 12: e0185868. <https://doi.org/10.1371/journal.pone.0185868>.
- Carkeet, A. 2015. Exact parametric confidence intervals for Bland–Altman limits of agreement. *Optometry and Vision Science* 92: e71–e80. <https://doi.org/10.1097/OPX.0000000000000513>.
- Carkeet, A., and Y. T. Goh. 2018. Confidence and coverage for Bland–Altman limits of agreement and their approximate confidence intervals. *Statistical Methods in Medical Research* 27: 1559–1574. <https://doi.org/10.1177/0962280216665419>.
- Chatfield, M. 2021. tolerance: Stata module to calculate tolerance intervals (normal distribution). Statistical Software Components S459009, Department of Economics, Boston College. <https://ideas.repec.org/c/boc/bocode/s459009.html>.

- Cole, T. J. 2000. Sympercents: Symmetric percentage differences on the 100 log(e) scale simplify the presentation of log transformed data. *Statistics in Medicine* 19: 3109–3125. [https://doi.org/10.1002/1097-0258\(20001130\)19:22%3C3109::AID-SIM558%3E3.0.CO;2-F](https://doi.org/10.1002/1097-0258(20001130)19:22%3C3109::AID-SIM558%3E3.0.CO;2-F).
- Cole, T. J., and D. G. Altman. 2017. Statistics Notes: What is a percentage difference? *BMJ* 358: j3663. <https://doi.org/10.1136/bmj.j3663>.
- Cox, N. J. 2018. Speaking Stata: Logarithmic binning and labeling. *Stata Journal* 18: 262–286. <https://doi.org/10.1177/1536867X1801800116>.
- . 2020. Software Updates: gr0072_1: Speaking Stata: Logarithmic binning and labeling. *Stata Journal* 20: 1028–1030. <https://doi.org/10.1177/1536867X20976342>.
- Dewitte, K., C. Fierens, D. Stöckl, and L. M. Thienpon. 2002. Application of the Bland–Altman plot for interpretation of method-comparison studies: A critical investigation of its practice. *Clinical Chemistry* 48: 799–801. <https://doi.org/10.1093/clinchem/48.5.799>.
- Doménech, J. M., and R. Sesma. 2021. agree: Passing–Bablok and Bland–Altman methods: User-written command agree for Stata. GitHub. <https://github.com/rsesma/stata/tree/master/agree>.
- Francq, B. G., M. Berger, and C. Boachie. 2020. To tolerate or to agree: A tutorial on tolerance intervals in method comparison studies with BivRegBLS R package. *Statistics in Medicine* 39: 4334–4349. <https://doi.org/10.1002/sim.8709>.
- Gerke, O. 2020. Reporting standards for a Bland–Altman agreement analysis: A review of methodological reviews. *Diagnostics* 10: 334. <https://doi.org/10.3390/diagnostics10050334>.
- Howe, W. G. 1969. Two-sided tolerance limits for normal populations—Some improvements. *Journal of the American Statistical Association* 64: 610–620. <https://doi.org/10.1080/01621459.1969.10500999>.
- Kirkwood, B. R., and J. A. C. Sterne. 2003. *Essential Medical Statistics*. 2nd ed. Oxford: Blackwell.
- Klein, D. 2018. Implementing a general framework for assessing interrater agreement in Stata. *Stata Journal* 18: 871–901. <https://doi.org/10.1177/1536867X1801800408>.
- Limpert, E., and W. A. Stahel. 2011. Problems with using the normal distribution—And ways to improve quality and efficiency of data analysis. *PLOS ONE* 6: e21403. <https://doi.org/10.1371/journal.pone.0021403>.
- Linden, A. 2021. rmloa: Stata module to compute limits of agreement for data with repeated measures. Statistical Software Components S458980, Department of Economics, Boston College. <https://ideas.repec.org/c/boc/bocode/s458980.html>.

- Ludbrook, J. 2010. Confidence in Altman–Bland plots: A critical review of the method of differences. *Clinical and Experimental Pharmacology and Physiology* 37: 143–149. <https://doi.org/10.1111/j.1440-1681.2009.05288.x>.
- Mander, A. 2005. batplot: Stata module to produce Bland–Altman plots accounting for trend. Statistical Software Components S448703, Department of Economics, Boston College. <https://ideas.repec.org/c/boc/bocode/s448703.html>.
- Meeker, W. Q., G. J. Hahn, and L. A. Escobar. 2017. *Statistical Intervals: A Guide for Practitioners and Researchers*. 2nd ed. Hoboken, NJ: Wiley.
- Royston, P., and J. N. S. Matthews. 1991. Estimation of reference ranges from normal samples. *Statistics in Medicine* 10: 691–695. <https://doi.org/10.1002/sim.4780100503>.
- Seed, P. 2000. sbe33: Comparing several methods of measuring the same quantity. *Stata Technical Bulletin* 55: 2–9. Reprinted in *Stata Technical Bulletin Reprints*. Vol. 10, pp. 73–82. College Station, TX: Stata Press.
- Shieh, G. 2018. The appropriateness of Bland–Altman’s approximate confidence intervals for limits of agreement. *BMC Medical Research Methodology* 18: 45. <https://doi.org/10.1186/s12874-018-0505-y>.
- Steichen, T. J., and N. J. Cox. 1998. sg84: Concordance correlation coefficient. *Stata Technical Bulletin* 43: 35–39. Reprinted in *Stata Technical Bulletin Reprints*. Vol. 8, pp. 137–143. College Station, TX: Stata Press.
- Taffé, P., M. Peng, V. Stagg, and T. Williamson. 2017. biasplot: A package to effective plots to assess bias and precision in method comparison studies. *Stata Journal* 17: 208–221. <https://doi.org/10.1177/1536867X1701700111>.
- Törnqvist, L., P. Vartia, and Y. O. Vartia. 1985. How should relative changes be measured? *American Statistician* 39: 43–46. <https://doi.org/10.2307/2683905>.
- Vangel, M. 2005. Tolerance interval. In *Encyclopedia of Biostatistics*, ed. P. Armitage and T. Colton, 8: 5477–5482, 2nd ed. Chichester, U.K.: Wiley. <https://doi.org/10.1002/0470011815.b2a15163>.
- Vardeman, S. B. 1992. What about the other intervals? *American Statistician* 46: 193–197. <https://doi.org/10.1080/00031305.1992.10475882>.
- Vock, M. 2016. Intervals for the assessment of measurement agreement: Similarities, differences, and consequences of incorrect interpretations. *Biometrical Journal* 58: 489–501. <https://doi.org/10.1002/bimj.201400234>.

About the authors

Mark Chatfield is a biostatistician and has enjoyed using Stata for 20 years. He is currently undertaking a PhD, among other things.

Tim Cole is a professor in medical statistics. He has long-standing interests in the statistics of human growth, where logarithms are useful for modeling purposes.

Henrica de Vet is a professor in clinimetrics. A clinical epidemiologist by training, she focuses on methodological research on the development and improvement of measurement instruments.

Louise Marquart-Wilson is a biostatistician with interests in generalized linear mixed models.

Daniel Farewell is a professor in statistics. He is interested in the development of novel statistical theory and methodology, especially for causal inference and missing data.

A Appendix

A.1 Calculation and reporting of regression-based estimates for bias and 95% LOA

The approach described in Bland and Altman (1999, sec 3.2) for differences can be written as follows:

A linear regression is used to estimate how the mean of differences (Mean_Y) varies linearly with the mean of the data pairs. A second linear regression is used to estimate how the SD of differences (SD_Y) varies linearly with the mean of the data pairs. These two relationships are then combined to estimate LOA (95% LOA_Y) that vary linearly with an average of the data pairs.

Equivalently, for a Bland–Altman plot of differences against means, use $Y = A - B$ and $X = (A + B)/2$ in the more general approach we outline below.

Mean_Y (or bias)

Fit a linear regression of Y on X :

```
. regress Y X
```

The resulting regression equation estimates the mean of Y as a linear function of X :

$$\text{Mean}_Y = b_0 + b_1X$$

SD_Y

First, obtain the residuals from the above regression:

```
. predict resid, resid
```

Second, calculate the absolute values of the residual and adjust by multiplying by $\sqrt{\pi/2}$: (Given a value of X , it is assumed that residuals are normally distributed with SD σ , and therefore the mean of absolute residuals is $\sigma\sqrt{2/\pi}$.)

```
. generate adj_abs_resid = abs(resid) * sqrt(_pi/2)
```

Third, fit a linear regression of `adj_abs_resid` on `X`:

```
. regress adj_abs_resid X
```

The resulting regression equation estimates the SD of Y as a linear function of X :

$$SD_Y = b_2 + b_3X$$

Estimated 95% LOA for Y

$$\begin{aligned} 95\% \text{ LOA}_Y &= \text{Mean}_Y \pm 1.96 \text{ SD}_Y \\ &= b_0 + b_1X \pm 1.96(b_2 + b_3X) \end{aligned}$$

that is,

$$\begin{aligned} \text{lower limit of agreement (LLOA)} &= b_0 - 1.96b_2 + (b_1 - 1.96b_3)X \\ \text{upper limit of agreement (ULOA)} &= b_0 + 1.96b_2 + (b_1 + 1.96b_3)X \end{aligned}$$

For a Bland–Altman plot of ratios against geometric means, we apply the approach to log-transformed data but express relationships in terms of ratios and geometric means. We use

$$\begin{aligned} Y &= \ln A - \ln B = \ln(A/B) = \ln(\text{Ratio}) \\ X &= (\ln A + \ln B)/2 = \ln\{\text{GMean}(A, B)\} \end{aligned}$$

For a Bland–Altman plot of percentage differences $100(\ln A - \ln B)\%$ against geometric means, we use

$$\begin{aligned} Y &= 100(\ln A - \ln B) \\ X &= (\ln A + \ln B)/2 = \ln\{\text{GMean}(A, B)\} \end{aligned}$$

For a Bland–Altman plot of percentage differences $100(A - B)/\{(A + B)/2\}\%$ against arithmetic means, we use

$$\begin{aligned} Y &= 100(A - B)/\{(A + B)/2\} \\ X &= \ln\{(A + B)/2\} = \ln\{\text{Mean}(A, B)\} \end{aligned}$$

The following output relates to section 3.1:

```
. blandaltman plexrbp4µmol1 nimanurbp4µmol1,
> plot(difference percentmean percent lmean ratio)
```

A: plexrbp4µmol1 7-Plex RBP4 (µmol/L)
 B: nimanurbp4µmol1 NiMaNu RBP4 (µmol/L)

DIFFERENCES...

Calculation	N	Mean	SD	Interval(s)
A-B	206	-.1022816	.2013955	

```
. regress difference mean
```

Source	SS	df	MS	Number of obs	=	206
Model	.061142301	1	.061142301	F(1, 204)	=	1.51
Residual	8.25368545	204	.040459242	Prob > F	=	0.2204
				R-squared	=	0.0074
				Adj R-squared	=	0.0025
Total	8.31482775	205	.040560135	Root MSE	=	.20114

__000002	Coefficient	Std. err.	t	P> t	[95% conf. interval]
__000003	-.0445447	.0362355	-1.23	0.220	-.1159888 .0268994
_cons	-.0559129	.0402386	-1.39	0.166	-.1352497 .023424

```
-> regression-based bias:        -.0559129 + -.0445447 × Mean(A,B)
```

```
. regress adj_abs_resid mean
```

Source	SS	df	MS	Number of obs	=	206
Model	1.22288054	1	1.22288054	F(1, 204)	=	48.97
Residual	5.09388346	204	.024970017	Prob > F	=	0.0000
				R-squared	=	0.1936
				Adj R-squared	=	0.1896
Total	6.31676401	205	.030813483	Root MSE	=	.15802

__00000C	Coefficient	Std. err.	t	P> t	[95% conf. interval]
__000003	.1992128	.0284665	7.00	0.000	.1430865 .2553392
_cons	-.027725	.0316114	-0.88	0.381	-.0900519 .0346019

```
-> regression-based SD:            -.027725 + .1992128 × Mean(A,B)
-> regression-based 95% LLOA:    -.0015729 + -.4349947 × Mean(A,B)
-> regression-based 95% UL0A:    -.1102529 + .3459053 × Mean(A,B)
```

PERCENTAGE DIFFERENCES (using Mean as denominator)...

Calculation	N	Mean	SD	Interval(s)
100*(A-B)/[(A+B)/2]	206	-10.07663	17.10901	

```
. regress percentmean ln_mean
```

Source	SS	df	MS	Number of obs	=	206
Model	29.4792833	1	29.4792833	F(1, 204)	=	0.10
Residual	59977.7646	204	294.00865	Prob > F	=	0.7518
				R-squared	=	0.0005
				Adj R-squared	=	-0.0044
Total	60007.2439	205	292.718263	Root MSE	=	17.147

__000005	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
__00000A	1.027934	3.246288	0.32	0.752	-5.372644	7.428513
_cons	-10.04945	1.197744	-8.39	0.000	-12.411	-7.687909

```
-> regression-based bias: -10.04945 + 1.027934 * ln(Mean(A,B))
```

```
. regress adj_abs_resid ln_mean
```

Source	SS	df	MS	Number of obs	=	206
Model	239.33826	1	239.33826	F(1, 204)	=	1.33
Residual	36737.0171	204	180.083417	Prob > F	=	0.2503
				R-squared	=	0.0065
				Adj R-squared	=	0.0016
Total	36976.3554	205	180.372465	Root MSE	=	13.42

__00000E	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
__00000A	2.928957	2.540644	1.15	0.250	-2.080331	7.938246
_cons	16.74617	.9373909	17.86	0.000	14.89795	18.59439

```
-> regression-based SD: 16.74617 + 2.928957 * ln(Mean(A,B))
```

```
-> regression-based 95% LLOA: -42.87134 + -4.712716 * ln(Mean(A,B))
```

```
-> regression-based 95% UL0A: 22.77244 + 6.768585 * ln(Mean(A,B))
```

PERCENTAGE DIFFERENCES (using Logarithmic Mean as denominator)...

Calculation	N	Mean	SD	Interval(s)
100*(A-B)/LMean(A,B)	206	-10.16368	17.29902	

```
. regress percentlmean ln_gmean
```

Source	SS	df	MS	Number of obs	=	206
Model	72.0980071	1	72.0980071	F(1, 204)	=	0.24
Residual	61275.3688	204	300.369455	Prob > F	=	0.6247
				R-squared	=	0.0012
				Adj R-squared	=	-0.0037
Total	61347.4668	205	299.255936	Root MSE	=	17.331

__000006	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
__000009	1.609633	3.285436	0.49	0.625	-4.868133	8.087398
_cons	-10.1131	1.211925	-8.34	0.000	-12.5026	-7.723594

```
-> regression-based bias: -10.1131 + 1.609633 * ln(GMean(A,B))
```

```
. regress adj_abs_resid ln_gmean
```

Source	SS	df	MS	Number of obs	=	206
Model	145.256099	1	145.256099	F(1, 204)	=	0.78
Residual	37812.2272	204	185.354055	Prob > F	=	0.3771
				R-squared	=	0.0038
				Adj R-squared	=	-0.0011
Total	37957.4833	205	185.158455	Root MSE	=	13.614

__00000G	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
__000009	2.284716	2.58087	0.89	0.377	-2.803884	7.373317
_cons	16.89376	.9520259	17.75	0.000	15.01669	18.77084

```
-> regression-based SD:      16.89376 + 2.284716 * ln(GMean(A,B))
-> regression-based 95% LLOA: -43.22427 + -2.868329 * ln(GMean(A,B))
-> regression-based 95% UL0A: 22.99807 + 6.087595 * ln(GMean(A,B))
```

```
RATIOS...
```

Calculation	N	GMean	GSD	Interval(s)
A/B	206	.9033576	1.188854	

```
. regress ln_ratio ln_gmean
```

Source	SS	df	MS	Number of obs	=	206
Model	.007209808	1	.007209808	F(1, 204)	=	0.24
Residual	6.12753704	204	.030036946	Prob > F	=	0.6247
				R-squared	=	0.0012
				Adj R-squared	=	-0.0037
Total	6.13474685	205	.029925594	Root MSE	=	.17331

__000008	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
__000009	.0160963	.0328544	0.49	0.625	-.0486813	.080874
_cons	-.101131	.0121192	-8.34	0.000	-.125026	-.0772359

```
-> regression-based GMean Ratio:      .9038146 * GMean(A,B)^ .0160963
```

```
. regress adj_abs_resid ln_gmean
```

Source	SS	df	MS	Number of obs	=	206
Model	.014525613	1	.014525613	F(1, 204)	=	0.78
Residual	3.7812229	204	.018535406	Prob > F	=	0.3771
				R-squared	=	0.0038
				Adj R-squared	=	-0.0011
Total	3.79574851	205	.018515846	Root MSE	=	.13614

__00000I	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
__000009	.0228472	.0258087	0.89	0.377	-.0280388	.0737332
_cons	.1689376	.0095203	17.75	0.000	.1501669	.1877084

```
-> regression-based GSD Ratio:      1.184046 * GMean(A,B)^ .0228472
-> regression-based 95% LLOA Ratio: .6490518 * GMean(A,B)^-.0286833
-> regression-based 95% UL0A Ratio: 1.258576 * GMean(A,B)^ .060876
```

The output for ratios uses the fact that $\exp[\alpha + \beta \ln\{GMean(A, B)\}] = \exp(\alpha) \times GMean(A, B)^\beta$.

A.2 Confidence intervals for LOA, PI, and TIs

Here we detail the calculation of confidence intervals for LOA and the calculation and meaning of prediction and TIs produced by `blandaltman`.

We assume differences (or percentage differences or log ratios) $\mathbf{y} = \{y_1, y_2, \dots, y_n\}$ are randomly sampled from a normal distribution with mean μ , SD σ , and cumulative distribution function F . We denote the sample size n , the sample mean \bar{y} , and the sample SD s .

95% confidence intervals for LOA

Bland and Altman (1999) viewed $\bar{y} \pm 1.96s$ as estimates of

$$\begin{aligned}\text{LLOA} &= \mu - 1.96\sigma \text{ (that is, 2.5th percentile of population)} \\ \text{ULOA} &= \mu + 1.96\sigma \text{ (that is, 97.5th percentile of population)}\end{aligned}$$

They acknowledged that the sampling error affects the estimates of μ , σ , and LOA and proposed calculating a 95% confidence interval for LOA. They described approximate methods assuming the sample size is large. However, there exists an exact method based on the noncentral t distribution (Carkeet 2015; Shieh 2018), which is implemented in `blandaltman`.⁴ The formulas are

$$\begin{aligned}\text{ULOA: } & \bar{y} + k_{\text{inner}} \times s \text{ to } \bar{y} + k_{\text{outer}} \times s \\ \text{LLOA: } & \bar{y} - k_{\text{outer}} \times s \text{ to } \bar{y} - k_{\text{inner}} \times s\end{aligned}$$

where

$$\begin{aligned}k_{\text{inner}} &= t_{n-1, 1.96\sqrt{n}, 0.025} \sqrt{\frac{1}{n}} \\ k_{\text{outer}} &= t_{n-1, 1.96\sqrt{n}, 0.975} \sqrt{\frac{1}{n}}\end{aligned}$$

and the quantities $t_{n-1, 1.96\sqrt{n}, 0.025}$ and $t_{n-1, 1.96\sqrt{n}, 0.975}$ are the 0.025 and 0.975 quantiles of the noncentral t distribution with $n - 1$ degrees of freedom and noncentrality parameter $1.96\sqrt{n}$. In contrast to the approximate confidence intervals, these exact confidence intervals will not appear symmetric about the LOA.

4. While this method is currently not implemented in `centile` (see [R] `centile`), it is now implemented in the recently revised community-contributed command `tolerance` (Chatfield 2021).

PI

A two-sided 95% PI for a single future observation y_{n+1} (Vardeman 1992; Meeker, Hahn, and Escobar 2017) is a random interval $[L(\mathbf{y}), U(\mathbf{y})]$ constructed such that

$$\text{Prob}\{L < y_{n+1} < U\} = 0.95$$

That is, if the process of 1) gathering a sample of size n , 2) constructing a 95% PI, and 3) gathering one additional y_{n+1} is repeated infinitely many times, then 95% of the PIs will contain y_{n+1} .

The 95% PI is calculated as $\bar{y} \pm k_{\text{PI}} \times s$, where

$$k_{\text{PI}} = t_{n-1, 0.975} \sqrt{1 + \frac{1}{n}}$$

and the quantity $t_{n-1, 0.975}$ is the 0.975 quantile of the Student's t distribution with $n-1$ degrees of freedom.

TIs

TIs are statistical intervals that contain at least a specified percentage of a population, either 1) on average or 2) with a stated confidence (Vangel 2005; Vardeman 1992).

1) 95% expectation TI

A two-sided 95% expectation TI is a random interval $[L(\mathbf{y}), U(\mathbf{y})]$ constructed such that

$$E\{F(U) - F(L)\} = 0.95$$

That is, if the process of a) gathering a sample of size n , b) constructing a 95% expectation TI, and c) calculating what percentage of the population is contained by the interval is repeated infinitely many times, then the mean (that is, expected) percentage will be 95%.

Mathematically, it is equivalent to the above-mentioned 95% PI.

2) 95% TI with $C\%$ confidence

A two-sided 95% TI with $C\%$ confidence is a random interval $[L(\mathbf{y}), U(\mathbf{y})]$ constructed such that

$$\text{Pr}\{F(U) - F(L) \geq 0.95\} = C\%$$

That is, if the process of a) gathering a sample of size n , b) constructing a 95% TI with $C\%$ confidence, and c) calculating what percentage of the population is contained by the interval is repeated infinitely many times, then $C\%$ of these intervals will contain at least 95% of the population.

There is no closed-form expression. `blandaltman` calculates an approximate two-sided 95% TI with $C\%$ confidence (Howe 1969): $\bar{y} \pm k_{\text{TI}} \times s$, where

$$k_{\text{TI}} = 1.96 \sqrt{\left(1 + \frac{1}{n}\right) \left(\frac{n-1}{\chi_{n-1, 1-C/100}^2}\right) \left\{1 + \frac{n-3 - \chi_{n-1, 1-C/100}^2}{2(n+1)^2}\right\}}$$

and the quantity $\chi_{n-1, 1-C/100}^2$ is the $(1-C/100)$ quantile of a χ^2 distribution with $n-1$ degrees of freedom.

Table A1. Factors used to calculate i) a 95% confidence interval for LOA, ii) a 95% PI, and iii) approximate 95% TIs with $C\%$ confidence. Intervals are calculated as described in section A.2.

n	(i) $k_{\text{inner}}, k_{\text{outer}}$	(ii) k_{PI}	(iii) k_{TI}	
			$C = 50$	$C = 95$
10	1.16, 3.80	2.37	2.13	3.41
20	1.36, 3.01	2.14	2.04	2.76
50	1.55, 2.53	2.03	1.99	2.38
100	1.66, 2.34	1.99	1.98	2.23
∞	1.96, 1.96	1.96	1.96	1.96