



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

**EFFECT OF VARIED CALCIUM FORMULATIONS AND TIME OF APPLICATION
ON POSTHARVEST QUALITY AND ORGANOLEPTIC ACCEPTABILITY OF
MANGO FRUITS****Bitange NM^{1*}, Chemining'wa GN¹, Ambuko J¹ and WO Owino²****Naphis Mokaya Bitange**

*Corresponding author email: mokayanaphis@gmail.com

¹Department of Plant Science and Crop Protection, University of Nairobi, P.O. Box 29053-00625, Kangemi, Nairobi, Kenya

²Department of Food Science and Technology, Jomo Kenyatta University of Agriculture and Technology, P.O Box 6200-00200, Nairobi, Kenya



ABSTRACT

Mango (*Mangifera indica L.*) is a highly perishable fruit with a short shelf life at ambient conditions, which may lead to post-harvest losses approximated to be 40-45%. This reduces returns to farmers significantly. The problem is compounded by the fact that most farmers do not have access to cold storage facilities. Nutrient management has been shown to affect postharvest characteristics of fruits. Calcium particularly plays a critical role in cell membrane integrity, tissue firmness and delays lipid catabolism. Previous studies have indicated a deficiency of calcium in some mango growing regions in Kenya. A field study was carried out to determine the effect of varied calcium formulations applied at various stages of growth on mango fruits post-harvest quality and organoleptic acceptability. The study was carried out in Embu County Eastern Kenya during seasons 2017/2018 and 2018 /2019 using "Van Dyke" cultivar, aged approximately 10 years. The experiment was set up in a randomized complete block design with a split-split plot arrangement, three trees per replication, replicated thrice. Three calcium formulations: calcium chloride, calcium nitrate and Easygro™ were applied at rates of 0%, 1.0%, 1.5% and 2.0% at fruit set, 30 days after fruit set and 30 days to physiological maturity. The calcium sources formed the main plots, the timing of application formed the subplots while the rates of application formed the sub-sub plots. Total soluble solids (TSS) and percentage titratable acidity (TA) were assessed at harvest and after 12 days of storage under ambient conditions ($25\pm2^{\circ}\text{C}$, $70\pm5\%$ relative humidity) using standard procedures. Selected fruits' sensory attributes were also evaluated after storage using a hedonic scale. Analysis of data was done using the 14th Edition of the Genstat software. The differences among the means of the treatments were compared using Fisher's Protected LSD test at 5% probability level. Fruits sprayed with calcium chloride, 2.0% at fruit set had higher TSS (6.8 ° brix and 6.3° brix) (10.47 ° brix and 9.10 ° brix), TA (1.29% and 1.27%), (0.77% and 0.675%) than other treatments at maturity and after storage in both seasons, respectively. Calcium chloride at 2.0% level of application led to a superior peel color appearance contrary to calcium nitrate and Easygro™ also applied at 2.0%, which led to an inferior peel color appearance and an inferior taste of fruits. Therefore, calcium nitrate and easy gro should be sprayed at concentration of 1.5% for good taste and peel colour appearance.

Key words: Mango, total soluble sugars, total titratable acidity, organoleptic, shelf-life



INTRODUCTION

Mango is an important fruit crop though faced with short shelf life challenges in ambient conditions, which may lead to high post harvest losses. Various physiological processes occur in the mango fruit during ripening that lead to disintegration and softening of the fruit cell wall and eventual senescence. Calcium is a primary component of the pectin that strengthens the cell wall, hence plays a key role in fruit firmness [1, 2] and influences other physiological processes [3].

Calcium compounds have been shown to enhance post harvest shelf life of mango [4, 5, 6] peach [7], guava [8], and avocado [9] fruits. However, excessive application of calcium may lead to inferior eating quality [10]. Calcium in the form of calcium chloride has been reported to delay fruit ripening rate thus increasing the shelf life of guava [11], apple [12], mango [2] and pear [13]. Calcium nitrate on the other hand has been reported to be more effective than calcium chloride in quality preservation of mango fruits [14]. The effectiveness of any calcium formulation depends on the time, frequency and concentration of applications [2, 4, 5]. Therefore, the purpose of this study was to compare the effectiveness of varied formulations of calcium applied at different times and rates in delaying the ripening rate of mango fruits while maintaining fruit quality.

MATERIALS AND METHODS

The experiment was set up in Embu County, Kenya, with an elevation of 1174m ASL and coordinates of 0° 32' S 37° 41'E in seasons 2017/2018 and 2018/2019. "Van Dyke" mango cultivar trees aged approximately 10 years were used. Calcium chloride, calcium nitrate and easy gro were applied at 0%, 1.0%, 1.5% and 2.0 % at three varied stages of fruit development (fruit set, 30 days after fruit set and 30 days to physiological maturity). Randomized complete block design was used with a split-split plot arrangement, three trees per replication, replicated three times. The calcium sources formed the main plots, the timing of application formed the subplots while the rates of application formed the sub-sub plots. Easygro is a foliar based fertilizer with a composition of: 14% nitrogen, 2.5% magnesium, 2% potassium, and 13% calcium. Fruits were harvested at physiological maturity and transported to the post harvest laboratory at Jomo Kenyatta University of Agriculture and Technology in plastic crates which were lined with dampened newspapers to prevent injury and minimize heat load during transit.



Data collection and analysis

The fruits' total soluble solids (TSS) and titratable acidity (TTA) were assessed at harvest and after 12 days of storage under ambient conditions ($25\pm2^{\circ}\text{C}$, $70\pm5\%$, relative humidity). Selected fruit sensory attributes were also determined after 12 days of storage at ambient conditions. Analysis of data was done using the 14th Edition of the Genstat software [15]. The differences among the means of the treatments were compared using Fisher's Protected LSD test at 5% probability level.

Total soluble solids and titratable acidity determination

Three fruits, from each treatment, were selected and a fruit juice obtained from each batch. Three ml of the extracted juice was placed on a hand refractometer (Model 500, Atago, Tokyo, Japan). Total soluble solids level was recorded in °Brix. Five ml of the extracted juice was diluted with 25 ml of distilled water. Ten ml of the diluted solution was then titrated with 0.1 N NaOH using phenolphthalein indicator [16]. The TTA was then expressed as a percentage of citric acid using the equation below:

Citric acid equivalent (%) =

$$\left(\frac{\text{Sample reading (ml)} \times \frac{\text{Dilution factor}}{\text{Sample weight}} \times 0.0064(\text{citric acid factor})}{100} \right) 100$$

Organoleptic attributes determination

Fifteen fruits were taken from each replication and sliced into approximately equal sized slices, which were then anonymously coded and placed on a white paper. A panel of 15 untrained, college researchers was guided on how to score selected sensory attributes (peel color, pulp taste and general acceptability) using a 7-point hedonic scale [17]: 1-Dislike extremely, 2-Dislike very much, 3-Dislike moderately, 4-Neither like nor dislike, 5-Like moderately, 6-Like very much and 7-Like extremely. The general acceptability of the fruits was determined by adding the scores of the evaluated attributes. The average of each of the panelists was then analysed and presented in graphs.

RESULTS AND DISCUSSION

Main and interactive effects of source, time and rate of calcium application on fruit TSS at harvest and after storage in ambient conditions

Source, rate and time of calcium application significantly ($p\leq0.05$) affected the fruit TSS at harvest and after 12 days of storage under ambient conditions in both seasons (Table 1). Interactions between source and time of calcium application



significantly ($P \leq 0.05$) affected the fruit TSS at harvest and after 12 days of storage at ambient conditions in both seasons (Table 2).

Application of calcium significantly ($P \leq 0.05$) reduced the accumulation of soluble solids at harvest and after storage in both seasons. The low TSS among fruits sprayed with calcium implies that they could be stored longer than those with higher TSS because they had not ripened fully hence have an enhanced shelf life. The lower levels of TSS at the end of the storage period in calcium sprayed fruits could be because of the role of calcium in delaying metabolic activities of fruits [6,18]. There was an increase in fruit TSS as the storage period advanced irrespective of the treatment. The increase in TSS with storage is because of the conversion of carbohydrates to sugars through enzymatic activities [2, 19]. Fruits sprayed at fruit set had significantly lower TSS than those sprayed after 30 days, which were in turn significantly lower than those sprayed 30 days to physiological maturity at harvest and after storage in both seasons. This indicates that calcium is more effective when applied in the early periods of fruit development [2].

The interaction between rate and time of calcium application had significant ($P \leq 0.05$) effects on the fruit TSS at harvest and after 12 days of storage in both seasons (Table 3). Fruits sprayed with 2.0% calcium had significantly lower TSS than those sprayed with 1.5%, which in turn had significantly lower values than those sprayed with 1.0% at harvest and after storage during both seasons and in most cases irrespective of the time of application.

The interactions between source, rate and time of calcium application significantly ($P \leq 0.05$) affected the TSS of the fruits at harvest in both seasons (Table 4).

Main and interactive effects of source, time and rate of calcium application on fruit TTA at harvest and after storage in ambient conditions

Source, rate and time of calcium application had significant ($P \leq 0.05$) effects on fruit TTA at harvest and after 12 days of storage in both seasons (Table 5). Interactions between source and time (Table 6) and between rate and time (Table 7) of calcium application significantly ($P \leq 0.05$) affected the fruit TTA at harvest and after storage in both seasons. Fruits sprayed with calcium chloride at fruit set and 30 days later had significantly higher TTA than those sprayed with calcium nitrate, which were in turn not significantly different from those sprayed with easy gro. Most of the fruits sprayed at 30 days to maturity did not have significantly different TTA from each other and were not significantly different from unsprayed ones. Calcium treated fruits had significantly higher TTA content than untreated fruits due to delayed fruit ripening associated with calcium [20, 21]. The high acidity in



calcium treated fruits has been previously reported [6, 22]. Fruits sprayed with calcium chloride had significantly higher TTA than those sprayed with easy gro at harvest and after storage in both seasons. Fruits sprayed at fruit set outperformed those sprayed 30 days after fruit set, which in turn outperformed those sprayed 30 days to physiological maturity at harvest and after storage regardless of the source of calcium. This is because calcium is more available at the early stages of fruit development as alluded in other studies [2,9]. Fruits sprayed at 2.0% had significantly higher TTA than those sprayed with calcium at 1.0% at harvest and after storage in both seasons.

Interactions between source, rate and time of calcium application significantly ($P \leq 0.05$) affected the fruit TTA at harvest in both seasons (Table 8) and in season 1 only after storage.

There was a general decrease in fruit acidity after storage. During storage under ambient conditions, fruits start to ripen as they approach climacteric period, which triggers an increase in malic enzyme and pyruvate decarboxylation reaction [23] hence a decrease in acidity with storage. Ripening leads to the accumulation of TSS that is a result of breakdown of starch to sugars. The ripening process was probably slower in calcium-treated fruits than control, thus a high acidity comparatively even after storage. Similar results showing decline in TTA after storage has been reported [2,15, 24].

Effect of source, rate and time of calcium application on sensory quality of fruits after storage in ambient conditions

All the calcium sources invariably affected fruit peel color in both seasons (Fig 1). Application of calcium chloride led to higher peel color scores than untreated fruits. Peel color was ranked with superior scores as the rate of calcium chloride increased. Calcium nitrate on the other hand led to high scores as evaluated by the panelists for concentration 1.0% and 1.5%. However, as the concentration of calcium increased from 1.5% to 2.0%, peel color was given inferior scores irrespective of the time of application. Similarly, application of easygro at higher rates (1.5% and 2.0%) registered lower scores than the control in both seasons. The enhanced peel color appearance due to calcium chloride application has been reported previously [4]. The same authors reported a deteriorated skin colour due to application of calcium ammonium nitrate. The deterioration in color could be due to nitrogen in easy gro and calcium nitrate that affected color formation. Fruit color decreases as nitrogen content increases [25,26] because nitrogen inhibits anthocyanin synthesis and accumulation [27].






Figure 1: Effect of different sources of calcium on the peel color of ripe mango fruits

Bars represent standard errors of the means at $p \leq 0.05$

S₁-Calcium chloride, S₂-Calcium nitrate, S₃-Easygro, X₀-Control, R₁-1.0%, R₂-1.5%,

R₃-2.0%, T₁-Fruit set, T₂-30 days after fruit set, T₃-30 days to physiological maturity

All the calcium sources affected the taste of fruits negatively in both seasons (Fig. 2). As the rate of calcium concentration increased the taste of the fruits deteriorated regardless of the source and time of application. Application of calcium at 30 days to maturity had superior taste scores than application at fruit set and 30 days later. This could be because earlier application led to more availability of calcium than later application [2, 9]. Calcium application led to deteriorated mango taste probably due to reduced accumulation of soluble solids with the application of calcium as previously reported [1, 14, 20], which led to a poor taste comparatively. This has also been reported in peach [28], apricot [29] and jujube [30]. Additionally, calcium may have imparted bitterness and saltiness, which results from residual calcium on the fruit mesocarp hence the unfavorable taste scores [31, 32, 33].




Figure 2: Effect of different sources of calcium on the taste of ripe mango fruits

Bars represent standard errors of the means at $p \leq 0.05$

S₁-Calcium chloride, S₂-Calcium nitrate, S₃-Easygro, X₀-Control, R₁-1.0%, R₂-1.5%,

R₃-2.0%, T₁-Fruit set, T₂-30 days after fruit set, T₃-30 days to physiological maturity

The general acceptability of the fruits was significantly ($p \leq 0.05$) affected by the source, rate and timing of application in both seasons (Fig.3). Fruits sprayed with calcium nitrate (1.5%) at 30 days to fruit maturity had significantly the highest scores. Application of calcium nitrate (2.0%) had the lowest acceptability rating irrespective of the timing of application in both seasons.




Figure 3: Effect of different sources of calcium on the general acceptability of ripe mango fruits

Bars represent standard errors of the means at $p \leq 0.05$

S₁-Calcium chloride, S₂-Calcium nitrate, S₃-Easygro, X₀-Control, R₁-1.0%, R₂-1.5%, R₃-2.0%, T₁-Fruit set, T₂-30 days after fruit set, T₃-30 days to physiological maturity

CONCLUSION

Calcium chloride (2.0%), applied at fruit set was more effective in maintenance of post harvest quality than other treatments. The taste of fruits was, however, negatively affected when calcium concentration increased to 2.0%, irrespective of the calcium source. Therefore, availability of calcium to the fruit should be optimised by either lowering the concentration to below 2% or spraying at 30 days after fruit set to achieve optimum levels. Similarly, calcium nitrate and easygro led to inferior peel colour when sprayed at 2.0%. Therefore, calcium nitrate and easy gro should only be sprayed at concentration of below 1.5% to achieve good peel colour.

ACKNOWLEDGEMENTS

This Research was supported by the Rockefeller Foundation (Yield Wise Initiative), Grant Number 2016 YWS 328.

Conflict of interest

Authors have declared that no competing interests exist.

Table 1: Main effects of source, time and rate of calcium application on fruit total soluble solids (°Brix) at harvest and after storage in ambient conditions

	Season 1		Season 2	
	At harvest	After storage	At harvest	After storage
Source				
Easy gro	11.73 ^b	15.43 ^b	10.43 ^b	14.37 ^b
Calcium nitrate	11.32 ^{bc}	15.10 ^b	10.26 ^b	14.16 ^b
Calcium chloride	10.19 ^c	14.50 ^b	9.81 ^b	13.27 ^b
Control	16.44 ^a	18.28 ^a	16.19 ^a	18.21 ^a
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05)	1.13	2.08	1.91	2.12
Cv%	18	14.5	18.9	15.7
Time				
T ₁	9.20 ^c	12.67 ^c	9.10 ^c	11.78 ^c
T ₂	12.24 ^b	16.12 ^b	10.52 ^b	15.09 ^b
T ₃	13.41 ^a	17.22 ^a	12.69 ^a	16.21 ^a
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05)	1.04	0.73	1.18	0.93
Cv%	17.50	9.30	21.40	12.60
Rate				
1.0%	11.82 ^b	15.59 ^b	11.6 ^b	14.84 ^b

1.5%	11.16 ^{bc}	15.07 ^{bc}	10.05 ^c	13.87 ^{bc}
2.0%	10.26 ^c	14.37 ^c	8.86 ^d	13.09 ^c
0%	16.44 ^a	18.28 ^a	16.19 ^a	18.21 ^a
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05)	1.13	1.19	0.94	1.68
Cv%	18.00	14.30	16.10	15.3

T₁=Fruit set; T₂=30 days after fruit set; T₃=30 days to physiological maturity; LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters in the same column are significantly different according to LSD at p ≤0.05



Table 2: Interactive effects of source and time of calcium application on fruit total soluble solids (°Brix) at harvest and after storage in ambient conditions

	Season 1		Season 2	
	At harvest	After storage	At harvest	After storage
Ctrl T ₁	16.67 ^a	18.20 ^a	16.17 ^a	18.30 ^a
Ctrl T ₃	16.50 ^a	18.17 ^a	16.13 ^a	18.03 ^a
Ctrl T ₂	16.17 ^a	18.22 ^a	16.27 ^a	18.30 ^a
S ₃ T ₃	13.96 ^b	17.58 ^{ab}	12.89 ^b	16.13 ^{bc}
S ₂ T ₃	13.19 ^c	17.01 ^{bc}	12.31 ^b	16.36 ^b
S ₃ T ₂	12.30 ^d	16.06 ^d	9.87 ^{cd}	15.37 ^{cd}
S ₂ T ₂	12.17 ^d	16.02 ^d	9.79 ^{cd}	14.91 ^d
S ₁ T ₃	12.06 ^d	16.77 ^c	11.72 ^b	15.53 ^{bcd}
S ₁ T ₂	10.96 ^e	15.50 ^d	9.99 ^c	13.92 ^e
S ₃ T ₁	8.94 ^f	12.66 ^e	8.54 ^e	11.60 ^{fg}
S ₂ T ₁	8.61 ^f	12.28 ^e	8.68 ^{de}	11.21 ^{fg}
S ₁ T ₁	7.57 ^g	11.23 ^f	7.72 ^e	10.34 ^g
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05) S*T	0.74	0.64	1.20	0.90
Cv%	6.80	4.40	11.80	6.70

S₁=Calcium chloride; S₂=Calcium nitrate; S₃=Easygro; R₁=1.0%; R₂=1.5%; R₃=2.0%; Ctrl=Control; T₁=Fruit set; T₂=30 days after fruit set; T₃=30 days to physiological maturity; S=Source; T=Time; R=Rate; LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters in the same column are significantly different according to LSD at p ≤0.05



Table 3: Interactive effects of rate and time of calcium application on fruit total soluble solids (°Brix) at harvest and after storage in ambient conditions

	Season 1		Season 2	
	At harvest	After storage	At harvest	After storage
Ctrl T ₁	16.67 ^a	18.20 ^a	16.17 ^a	18.30 ^a
Ctrl T ₃	16.50 ^a	18.17 ^a	16.13 ^a	18.03 ^a
Ctrl T ₂	16.17 ^a	18.47 ^a	16.27 ^a	18.30 ^a
R ₁ T ₃	13.90 ^b	17.57 ^{bc}	13.62 ^b	16.98 ^b
R ₂ T ₃	13.08 ^c	17.09 ^{cd}	12.36 ^c	15.87 ^c
R ₁ T ₂	12.32 ^d	16.30 ^{ef}	11.31 ^d	15.52 ^c
R ₃ T ₃	12.22 ^d	16.70 ^{de}	10.94 ^d	15.18 ^{cd}
R ₂ T ₂	11.88 ^{de}	15.83 ^{fg}	9.73 ^e	14.67 ^{de}
R ₃ T ₂	11.22 ^e	15.44 ^g	8.60 ^f	14.01 ^e
R ₁ T ₁	9.24 ^f	12.90 ^h	9.86 ^e	12.02 ^f
R ₂ T ₁	8.53 ^f	12.29 ⁱ	8.07 ^f	11.07 ^g
R ₃ T ₁	7.34 ^g	10.98 ⁱ	7.02 ^g	10.07 ^h
P-value	<.001	<.001	<.001	<.001
Lsd ($P \leq 0.05$) R*T	0.75	0.54	0.61	0.75
Cv%	6.8	3.8	6.0	5.5

R₁=1.0%; R₂=1.5%; R₃=2.0%; Ctrl=Control; T₁=Fruit set; T₂=30 days after fruit set; T₃=30 days to physiological maturity; T=Time; R=Rate; LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters in the same column are significantly different according to LSD at $p \leq 0.05$



Table 4: Interactive effects of source, rate and time of calcium application on fruit total soluble solids (°Brix) at harvest

Source	Rate	Season 1			Season 2		
		T₁	T₂	T₃	T₁	T₂	T₃
Calcium chloride	1.0%	8.30 ^{pq}	11.53 ^{ij}	13.40 ^{cd}	9.17 ^{jk}	11.43 ^{efg}	12.93 ^{cd}
	1.5%	7.60 ^r	11.00 ^{jk}	12.10 ^{ghi}	7.67 ^{no}	9.83 ^{ij}	11.77 ^{ef}
	2.0%	6.80 ^s	10.33 ^{lm}	10.67 ^{kl}	6.33 ^p	8.70 ^{klm}	10.47 ^{hi}
Calcium nitrate	1.0%	9.47 ^{no}	12.60 ^{efg}	13.80 ^c	10.80 ^{gh}	11.03 ^{fgh}	13.47 ^c
	1.5%	8.67 ^p	12.20 ^{gh}	13.17 ^{de}	8.17 ^{mn}	9.87 ^{ij}	12.30 ^{de}
	2.0%	7.70 ^{qr}	11.70 ^{hi}	12.60 ^{efg}	7.07 ^{op}	8.47 ^{lmn}	11.17 ^{fgh}
Easy gro	1.0%	9.97 ^{mn}	12.83 ^{def}	14.50 ^b	9.60 ^{ij}	11.47 ^{efg}	14.47 ^b
	1.5%	9.33 ^o	12.43 ^{fg}	13.97 ^{bc}	8.37 ^{lmn}	9.50 ^{jk}	13.00 ^{cd}
	2.0%	7.53 ^r	11.63 ^{hi}	13.40 ^{cd}	7.67 ^{no}	8.63 ^{klm}	11.20 ^{fgh}
Control	0%	16.67 ^a	16.17 ^a	16.50 ^a	16.17 ^a	16.27 ^a	16.13 ^a
P-value		0.008			0.038		
Lsd (P≤0.05)	SxTxR	0.63			0.87		
Cv (%)		3.30			4.90		

T1=Fruit set; T2=30 days after fruit set; T3=30 days to physiological maturity; S=Source; T=Time; R=Rate; LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters are significantly different according to LSD at p ≤0.05



Table 5: Main effects of source, time and rate of calcium application on fruit total titratable acidity (%) at harvest and after storage in ambient conditions

Source	Season 1		Season 2	
	At harvest	After storage	At harvest	After storage
Easy gro	0.41 ^b	0.23 ^b	0.49 ^b	0.26 ^b
Calcium nitrate	0.42 ^b	0.26 ^{ab}	0.54 ^{ab}	0.30 ^{ab}
Calcium				
chloride	0.59 ^a	0.33 ^a	0.67 ^a	0.35 ^a
Control	0.13 ^c	0.05 ^c	0.18 ^c	0.15 ^c
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05)	0.14	0.09	0.13	0.07
Cv %	39.1	26.5	34.5	23.5
Time				
T ₁	0.72 ^a	0.41 ^a	0.74 ^a	0.41 ^a
T ₂	0.37 ^b	0.24 ^b	0.55 ^b	0.27 ^b
T ₃	0.24 ^c	0.10 ^c	0.28 ^c	0.19 ^c
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05)	0.10	0.07	0.10	0.05
Cv %	40	34.8	26.5	26.4

Rate

1.0%	0.35 ^b	0.19 ^b	0.45 ^b	0.23 ^{bc}
1.5%	0.46 ^b	0.25 ^b	0.56 ^{ab}	0.29 ^b
2.0%	0.62 ^a	0.38 ^a	0.68 ^a	0.40 ^a
0%	0.13 ^c	0.05 ^c	0.18 ^c	0.15 ^c
<hr/>				
p-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05)	0.14	0.09	0.13	0.06
Cv%	37.5	63.7	43.8	39.1

T1=Fruit set; T2=30 days after fruit set; T3=30 days to physiological maturity;
 LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters in
 the same column are significantly different according to LSD at $p \leq 0.05$



Table 6: Interactive effect of source and time of calcium application on fruit total titratable acidity (%) at harvest and after storage in ambient conditions

Treatment	Season 1		Season 2	
	At Harvest	After storage	At harvest	After storage
S ₁ T ₁	0.93 ^a	0.50 ^a	0.99 ^a	0.52
S ₂ T ₁	0.74 ^{bc}	0.48 ^{abc}	0.73 ^{bc}	0.40
S ₃ T ₁	0.69 ^{bc}	0.39 ^{abc}	0.69 ^{bcd}	0.38
S ₁ T ₂	0.57 ^c	0.36 ^{bc}	0.64 ^{bcd}	0.30
S ₃ T ₂	0.32 ^d	0.20 ^d	0.55 ^d	0.27
S ₂ T ₂	0.29 ^{de}	0.22 ^d	0.60 ^{cd}	0.29
S ₁ T ₃	0.29 ^{de}	0.13 ^{de}	0.36 ^e	0.23
S ₃ T ₃	0.23 ^{de}	0.09 ^e	0.24 ^f	0.15
S ₂ T ₃	0.23 ^{de}	0.10 ^e	0.29 ^{ef}	0.21
S ₄ T ₁	0.14 ^{de}	0.04 ^e	0.20 ^f	0.18
S ₄ T ₂	0.14 ^{de}	0.08 ^e	0.19 ^f	0.16
S ₄ T ₃	0.11 ^e	0.02 ^e	0.14 ^f	0.09
P-value	<.001	<.001	<.001	0.09
LSD _(P≤0.05)	S*T	0.18	0.14	0.16
CV%	31.20	41.40	22.30	21.20

S₁=Calcium chloride; S₂=Calcium nitrate; S₃=Easygro; T₁=Fruit set; T₂=30 days after fruit set; T₃=30 days to physiological maturity; S=Source; T=Time; LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters in the same column are significantly different according to LSD at p ≤0.05



Table 7: Interactive effects of rate and time of calcium application on fruit total titratable acidity (%) at harvest and after 12 days of storage in ambient conditions

Treatment	Season 1		Season 2	
	At harvest	After storage	At harvest	After storage
R ₃ T ₁	0.99 ^a	0.65 ^a	0.96 ^a	0.53 ^a
R ₂ T ₁	0.76 ^b	0.41 ^b	0.78 ^{bc}	0.41 ^b
R ₁ T ₁	0.61 ^c	0.31 ^c	0.67 ^{cd}	0.35 ^c
R ₃ T ₂	0.53 ^c	0.34 ^c	0.71 ^{bcd}	0.42 ^b
R ₂ T ₂	0.36 ^{de}	0.24 ^d	0.59 ^{de}	0.25 ^d
R ₃ T ₃	0.33 ^{de}	0.17 ^e	0.36 ^f	0.24 ^{de}
R ₁ T ₂	0.28 ^{de}	0.20 ^{de}	0.48 ^e	0.18 ^f
R ₂ T ₃	0.25 ^{ef}	0.09 ^f	0.31 ^{fg}	0.19 ^{ef}
R ₁ T ₃	0.17 ^{fg}	0.06 ^g	0.22 ^{gh}	0.16 ^f
Ctrl T ₁	0.14 ^{fg}	0.04 ^g	0.20 ^{gh}	0.18 ^f
Ctrl T ₂	0.14 ^{fg}	0.08 ^g	0.19 ^{gh}	0.16 ^f
Ctrl T ₃	0.11 ^g	0.02 ^g	0.14 ^h	0.09 ^g
P-value	<.001	<.001	<.001	<.001
Lsd _(P≤0.05) R*T	0.11	0.06	0.15	0.05
Cv%	27	27.1	20.7	19.3

S₁=Calcium chloride; S₂=Calcium nitrate; S₃=Easygro; T₁=Fruit set; T₂=30 days after fruit set; T₃=30 days to physiological maturity; S=Source; T=Time; LSD=Least significant difference; CV=Coefficient of variation; Ctrl=Control. Treatments with different letters in the same column are significantly different according to LSD at p ≤0.05



Table 8: Interactive effects source, rate and time of calcium application on fruit total titrable acidity (%) at harvest

Source	Rate	Season 1			Season 2		
		T ₁	T ₂	T ₃	T ₁	T ₂	T ₃
Calcium chloride	1.0%	0.66 ^{cd}	0.46 ^f	0.21 ^{jklm}	0.80 ^{cd}	0.52 ^{jk}	0.24 ^p
	1.5%	0.83 ^b	0.55 ^e	0.30 ^{ghi}	0.91 ^b	0.64 ^{ghi}	0.37 ^{mn}
	2.0%	1.29 ^a	0.69 ^c	0.35 ^g	1.27 ^a	0.76 ^{cdef}	0.48 ^{kl}
Calcium nitrate	1.0%	0.59 ^{de}	0.20 ^{jklm}	0.15 ^{klmn}	0.64 ^{ghi}	0.50 ^{kl}	0.19 ^{pq}
	1.5%	0.78 ^b	0.30 ^{ghi}	0.23 ^{ijk}	0.71 ^{efg}	0.60 ^{hij}	0.33 ^{no}
	2.0%	0.85 ^b	0.37 ^g	0.32 ^{gh}	0.83 ^{bc}	0.70 ^{fg}	0.36 ^{mn}
Easy gro	1.0%	0.57 ^e	0.18 ^{jklmn}	0.15 ^{klmn}	0.56 ^{ijk}	0.42 ^{lm}	0.22 ^{pq}
	1.5%	0.67 ^{cd}	0.24 ^{hij}	0.22 ^{jk}	0.72 ^{defg}	0.55 ^{ik}	0.24 ^p
	2.0%	0.82 ^b	0.54 ^{ef}	0.33 ^g	0.79 ^{cde}	0.67 ^{gh}	0.26 ^{op}
Control	0%	0.14 ^{mn}	0.14 ^{mn}	0.11 ⁿ	0.20 ^{pq}	0.19 ^{pq}	0.14 ^q
p-value		<.001			<.001		
Lsd ($P \leq 0.05$)	SxTxR	0.08			0.09		
Cv (%)		11			10		

T₁=Fruit set; T₂=30 days after fruit set; T₃=30 days to physiological maturity; S=Source; T=Time; R=Rate; LSD=Least significant difference; CV=Coefficient of variation. Treatments with different letters are significantly different according to LSD at $p \leq 0.05$



REFERENCES

1. **Daundasekera WAM, Liyanage GLG, Wijerathne RY and R Pieris** Preharvest calcium chloride application improves postharvest keeping quality of tomato (*Lycopersicon esculentum* Mill.). *Ceylon Journal of Science*, 2015; **44**(1):55-60.
2. **Karemera UNJ and S Habimana** Performance of calcium chloride sprays on ripening, shelf life and physical chemical properties of mango fruits (*Mangifera indica* L.) CV. Totapuri. *International Invention Journal of Agricultural and Soil Science*, 2014; **2**(3):33-38.
3. **Njoroge CK, Kerbel EL and DP Briskin** Effect of calcium and calmodulin antagonists on ethylene biosynthesis in tomato fruits. *Journal of the Science of Food and Agriculture*, 1998; **76**:209-214.
4. **Anjum MA and H Ali** Effect of various calcium salts on ripening of mango fruits. *Journal of Research (Science)*, 2004; **15**(1):45-52.
5. **Poojapant and CP Singh** Response of pre harvest spray of chemicals on shelf life and quality of mango (*Mangifera indica* L.) Cv. Dashehari. *International Journal of Basic and Applied Agricultural Research*, 2014; **12**(3):374-378.
6. **Singh V, Pandey G, Sarolia DK, Kaushik RA and JS Gora** Influence of pre harvest application of calcium on shelf life and fruit quality of mango (*Mangifera indica* L.) cultivars. *International Journal of Current Microbiology and Applied Sciences*, 2017; **6**(4):1366-1372.
7. **EI-Alakmy HA** Effect of calcium application and wrapping treatments on fruit quality of Earlgrande Peach Trees (*Prunus persica* L.). *Journal of Applied Sciences Research*, 2012; **8**(7):3845-3849.
8. **Penter MG and PJC Stassen** The effect of pre-and post- harvest calcium application on the post-harvest quality of Pinkerton avocados. *South Africa Avocado Growers' Association Yearbook*, 2000; **23**:1-7.
9. **Njuguna J, Ambuko J, Hutchinson M and W Owino** Incidence of jelly seed disorder in 'Tommy Atkins' and 'Van Dyke' Mangoes as affected by agro-ecological conditions in Kenya. *International Journal of Plant and Soil Science*, 2016; **11**(5):1-9.



10. **Gangle A, Kirar SK and CS Pandey** Effect of post-harvest treatments on shelf life and quality of guava (*Psidium guajava L.*) cv. Allahabad Safeda. *International Journal of Current Microbiology and Applied Sciences*, 2019; **8(10)**:2104-2114.
11. **Hussain PR, Meena RS, Dar MA and AM Wani** Effect of post-harvest calcium chloride dip treatment and gamma irradiation on storage quality and shelf-life extension of Red delicious apple. *Journal of Food Science and Technology*, 2012; **49**:415-426.
12. **Sajid M, Mukhtiar M, Rab A, Shah S T and I Jan** Influence of Calcium Chloride on fruit quality of pear (*Pyrus communis*) cv. le conte during storage. *Pakistan Journal of Agricultural Sciences*, 2014; **51**:113-121.
13. **Mounika T, Reddy NN, Jyothi L N and V Joshi** Studies on the effect of post harvest treatments on shelf life and quality of mango [*Mangifera indica* L.]cv. Amrapali. *Journal of Applied and Natural Science*, 2017; **9(4)**:2055-2061.
14. **Ngamchuachit P, Sivertsen H K, Mitcham E J and DM Barrett** Effectiveness of calcium chloride and calcium lactate on maintenance of textural and sensory qualities of fresh-cut mangos. *Journal of Food Science*, 2014; **0(0)**:1-9.
15. **Payne R W, Murray D A, Harding S A, Baird D B and D M Soutar** An introduction to Genstat for Windows (14th Edition). VSN International, Hemel Hempstead, UK 2011.
16. **Rangana S** Manual of analysis of fruit and vegetable Products. Tata McGraw Hill Pub. Co. Ltd., New Delhi, 1979; 634.
17. **Galan SV, Fernandez GD and R Calvo** Incidence of softnose on mangoes in the Canary Islands. *Proceedings of the Fla State Horticulture Society*, 1984; **97**:358-360.
18. **Conway W S, Sams C E and KD Hickey** Pre- and postharvest calcium treatment of apple fruit and its effect on quality. *Acta Horticulturae*, 2002; **594**:413-419.
19. **Wahdan M T, Habib S E, Bassal M A and EM Qaoud** Effect of some chemicals on growth, fruiting, yield and fruit quality of "Succary Abiad" mango cv. *Journal of Horticulture Science*, 2011; **7(2)**:651-658.

20. **Madani B, Mirshekari A, Sofo A and MTM Mohamed** Preharvest calcium applications improve post harvest quality of papaya fruits. *Carica papaya L. Cv. Eksotika II. Journal of Plant Nutrition*, 2016; **39**(10):1483-1492.
21. **Mahmud S, Shibly A Z, Hossain M M, Islam S M and R Islam** The effect of calcium carbide and different calcium salts on mango fruits ripening in Bangladesh. *Journal of Pharmacology and Phytochemistry*, 2015; **4**(1):210-215.
22. **Dhillon B S and K Sukhjit** Effect of post harvest application of calcium chloride on storage life of mango var. Dashehari fruits. *HortFlora Research Spectrum*, 2013; **2**(3):265-267.
23. **Rhodes MJC, Woodtorton LSC, Gallard T and AC Hulme** Metabolic changes in excised fruit tissue iv. Factors affecting the development of a malate decarboxylation system during the ageing of disc of pre-climacteric apples. *Phytochem*, 1968; **7**:271–276.
24. **Islam KMD, Khan M ZH, Sarkar MAR, Absar N and SK Sarkar** Changes in acidity, TSS and sugar content at different storage periods of the post harvest mango (*Mangifera indica L.*) influenced by Bavistin DF. *International Journal of Food Science*, 2013; 1-2.
25. **Week KD, Southwick FW and M Drake** Relationship of leaf nitrogen, potassium content and color development of apple fruit. *Proceedings of American Society for Horticultural Science*, 1952; **60**:11-15.
26. **Fallahi E, Colt WM, Baird CR, Fallahi B and I J Chun** Influence of nitrogen and bagging on fruit quality and mineral concentrations of 'BC-2 Fuji' apple. *Hort Technology*, 2001; **11**:462-466.
27. **Cheng L and H Wang** Nitrogen fertilization has differential effects on red color development and flesh starch breakdown of 'Gala' Apple. New York. *Fruit Quarterly*, 2011; **19**(3):11-15.
28. **Gofure A, Shafique MZ, Helali M, Ibrahim M, Rahman MM and MS Alam** Studies on extension of post-harvest storage life of mango (*Mangifera indica L.*). *Bangladesh Journal of Scientific and Industrial Research*, 1997; **32**:148-152.

29. **Hamzehzad K, Rabiei V, Naseri L and S Hemati** Effect of UV-C irradiation and calcium chloride treatment on the quality and storage life of peach fruit, cv. Zafarany. *Iranian Journal of Horticultural Sciences*, 2010; **40**(4):53– 59.
30. **Antunes MDC, Correia MP, Miguel MG, Martins MA and MA Neves** The effect of calcium chloride postharvest application on fruit storage ability and quality of 'Beliana' and 'Lindo' apricot (*Prunus armeniaca* L.) cultivars. *Acta Horticulturae*, 2003; **604**:721–726.
31. **Moradinezhad F, Ghesmati M and M Khayyat** Postharvest calcium salt treatment of fresh jujube fruit and its effects on biochemical characteristics and quality after cold storage. *Journal of Horticultural Research*, 2019; **27**(2):39–46.
32. **Lovera N, Ramallo L and V Salvadori** Effect of processing conditions on calcium content, firmness, and color of papaya in syrup. *Journal of Food Processing*, 2014; 1-8.
33. **Luna-Guzmán and DM Barrett** "Comparison of calcium chloride and calcium lactate effectiveness in maintaining shelf stability and quality of fresh-cut cantaloupes". *Postharvest Biology and Technology*, 2000; **19**(1):61–70.