



**AgEcon** SEARCH  
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

## CLIMATE CHANGE AND VARIABILITY AFFECT RANGELAND QUALITY AND PRODUCTIVITY- HOW?

Timpong-Jones E C<sup>1</sup>, Owusu-Bremang R<sup>2</sup>, Mopipi K<sup>3</sup> and FO Sarkwa<sup>1\*</sup>



Felix O Sarkwa

\*Corresponding author email: [ofsarkwa@ug.edu.gh](mailto:ofsarkwa@ug.edu.gh) or [sarkwafelix@yahoo.com](mailto:sarkwafelix@yahoo.com)

<sup>1</sup>Livestock and Poultry Research Centre, School of Agriculture, College of Basic and Applied Sciences, University of Ghana, Legon, Ghana

<sup>2</sup>Department of Animal Science, School of Agriculture, College of Basic and Applied Sciences, University of Ghana, Legon, Ghana

<sup>3</sup>Department of Range and Forestry Resources, Faculty of Natural Resources, Botswana University of Agriculture and Natural Resource, Gaborone, Botswana



## ABSTRACT

Climate change is a principal threat to the sustainability of our planet which consists partly of the world's rangelands. According to numerous studies and reports, the trend at which the global climate is changing is a clear indication that the long term sustainability of global rangelands is uncertain. There is abundant evidence of climate change across the globe. The global average surface temperature is predicted to increase between 1.8°C and 4.0°C by 2100. This will no doubt have adverse and tremendous impact on the quality and productivity of rangelands, which in turn is expected to affect ruminant livestock production since about 70 % of forages fed on by animals are produced from these rangelands. The current climate change is basically as a result of anthropogenic activities leading to a build up of greenhouse gases in the atmosphere. These gases are trapped in the atmosphere and in return increase global surface temperature and alter known climatic patterns globally. This review, therefore, highlights the effect of climate change on rangeland quality and productivity considering how the drivers of climate change impact on the length of the growing period, plant species composition, nutritional value of forages, drought stress in plants, lignification, vegetation flammability and mortality/extinction of range plant and animal species. The review revealed that, climate change will result into the extinction of over one million terrestrial species in the next five decades. The elevation of CO<sub>2</sub> levels will increase non-structural carbohydrate content of grasses by 25 % and reduce their nitrogen content by about 8 %. It is also reported that, for every 1 % rise in temperature, there will be 0.4 % rise in Neutral Detergent Fibre decreasing forage intake and digestibility. Rising temperature also results in faster decomposition of soil organic matter. However, minerals will be lost through leaching and erosion if mineral uptake by rangeland plant species is slower than the rate of mineral release into the soil. It is clear that changing climate is having a toll on rangeland quality and productivity, the impact is mixed and difficult to quantify. It is recommended that, much research efforts be put into quantification of climate change impacts on rangelands through modelling and projections simplified for the comprehension of the resource poor livestock farmers particularly in developing countries. Vulnerable rural based livestock farmers should be assisted with adaptation strategies to remain in production.

**Key words:** Composition of plant species, drought stress, nutritional value, plant mortality



## INTRODUCTION

Climate change is presently regarded as a major and predominant threat to the sustainability of the planet Earth, which includes rangelands [1]. Naturally, the earth has experienced some alterations in its climate through changes in the orbital variation and also from volcanic eruptions, causing changes in the amount of solar radiations reaching the earth. However, human activities are substantially contributing to climate change [2, 3]. The agricultural sector is therefore vulnerable by the exposure and responsiveness to these climate conditions and the ability to adapt or cope with the changing climate [1]. There is therefore the need for farmers to have several strategies to employ to cope with the anticipated climate variability, which include adjusting their farming practices as required (that is applying adaptive range management and climate adaptation strategies). The total earth's terrestrial surface is covered by about 50 % of rangeland ecosystems [4]. Millions of people around the world depend on the grazing resources of rangelands as their source of livelihood.

Livestock contributes about 30 % of agricultural gross domestic product in developing countries, with a projected increase to about 40 % by the year 2030 [5] and is becoming the fastest growing sub-sector in agriculture. However, report indicates that there is an uncertainty in the long term sustainability of rangelands [6], since the effect of climate change will have a major impact on these rangelands as well as farmers. Economically developing countries and regions of the world are the ones whose rangeland resources are likely to suffer and are the most vulnerable since they are not capable of adapting to the unexpected change in climate and other extreme environmental conditions. Many of these economically developing countries are characterized by high population growth and as such are forced to use areas of very marginal productivity as grazing lands, and this tends to put stress on already stressed landscape due to climate change [7].

The agriculture sector is expected to be affected by climate change due to direct and indirect effect of climate change on crop and rangeland production. Rangelands provide about 70 % of the forage needs of ruminant livestock across the globe [4], thus, the livestock sub-sector will not be spared. Drivers of climate change will cause changes in rangeland ecosystems. The impact is expected to be severe in sub-Saharan Africa where a vast land area falls under the Arid and Semi-arid regions of the world and also because Agricultural activities in these regions is heavily dependent on climate [8]. Therefore, for a sustainable livestock production, there is the need to adequately understand the effect of climate change on



rangelands. This will be helpful in the formulation of appropriate range management decisions to optimize forage production in a changing climate. Thus, this review highlights the impact of climate change on rangeland quality and productivity emphasizing how drivers of climate change affect the length of growing period, plant species composition, nutritional value of forages, drought stress in plants, vegetation flammability and mortality/extinction of valuable plant and animal species.

## LITERATURE REVIEW

### Climate Change

Climate is often described by the statistical interpretation of rainfall and temperature data recorded over a long period of time for a given region or location [2]. Climate is usually the weather conditions in a given location averaged over a 30-year time period. The Intergovernmental Panel on Climate Change Third Assessment Report (IPCC TAR) defined climate change as any alteration in the climate over a period of time, as a result of anthropogenic or natural factors [9]. It is also defined as a change of climate which is associated to human activity that modifies the composition of the atmosphere and also includes natural climate variability observed over a period of time. Climate variability on the other hand, is defined as “the variation in climate on all temporal and spatial scales beyond that of individual weather events”. The United Nations Framework on Climate Change [10] makes a contrast between ‘climate change’ which is caused by human activities on atmospheric composition of the world and ‘climate variability’ which is attributable to natural causes. Based on these definitions, it can therefore be implied that climate change can occur as a result of natural variability and human activity [11].

### Greenhouse gases and climate change

The atmosphere is composed of a mixture of gases which are essential to life, and as such our lives depend on it [11]. Greenhouse gases (GHGs) which consist of the trace gases in the atmosphere tend to be increasing fast and contributing significantly to climate change. The GHGs include carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), Chlorofluorocarbons (CFCs), and tropospheric ozone (O<sub>3</sub>). Carbon dioxide’s involvement in global warming is massively higher than the other GHGs because of its high emission into the atmosphere [12]. Other GHGs such as CH<sub>4</sub>, CFCs and N<sub>2</sub>O are also fairly increasing. In the past 200 years, CO<sub>2</sub> concentration in the atmosphere has increased by more than 30 %, CH<sub>4</sub> by more than 150 % and N<sub>2</sub>O by 17 % [13]. Scientists are more than 95 % certain that global warming is caused mostly by increasing concentrations of these GHGs



through anthropogenic activities [14]. Some of these activities include burning of fossil fuel, livestock production (during respiration, they release CO<sub>2</sub> and CH<sub>4</sub>, their excreta releases N<sub>2</sub>O and NH<sub>3</sub>), felling of wood indiscriminately which affect the carbon balance, landfill systems and other forms of industrial air-pollution [11]. The aforementioned confirms that profound changes in the mechanisms of the atmosphere is due to human activities [12].

### **Evidence of climate change**

The IPCC [15] provides clear evidence that climate change is a reality, as shown by the increase in global temperatures and changes in rainfall patterns experienced globally. Evidence shows that the relative changes in climate over the past century have already had significant impacts on a wide range of plant species in terms of their distribution, abundance, phenology and physiology. Many instances of alteration in species distributions toward the poles have been documented [16]. Reports from IPCC revealed that there will be more wet areas which will lead to severe floods as well as dry areas which will also lead to severe drought, all in the tropics with low latitudes [2, 14]. Additionally, temperatures are anticipated to rise further by 1 to 4 °C in some decades to come [17]. Report on the African continent indicates that, there has been a rise in the mean annual land surface temperature above the 0.7 °C recorded in the 20th century [18]. The above-mentioned evidence of climate change is the reality the entire earth is faced with today.

### **Rangelands and Climate Change**

Rangelands can be defined as ecosystems which are capable of supporting grazing and browsing animals [19]. Globally, rangelands are estimated to occupy 50 % of the entire earth's terrestrial surface [19]. Their expanse makes rangelands very important to humans with their importance far outweighing their challenges [4]. The composition and productivity of rangelands are basically influenced by rainfall, fire, grazing, changes in temperature and the concentration of atmospheric CO<sub>2</sub> over longer time frames [20].

Climate change puts enormous pressure on rangelands [21] and as such, the phenomenon is now recognized as a dominant issue of concern for their sustainability. Species distribution in rangeland communities are likely to be altered in relation to changes in rainfall patterns and temperature levels. Some rangeland species are likely to migrate as temperature increases similarly to the reported occurrence in the early to mid-Holocene (10,500 to 5,000 years ago) [22]. Climate change is reported to be responsible for the extinction of many plant species, although, the extent and nature of this future occurrence is very difficult to predict.



In most plant species, there is an increase in the rate of photosynthesis when the concentration of atmospheric CO<sub>2</sub> increases and this leads to increase of plant growth, decreased water use and reduced tissue concentrations of protein and nitrogen [23]. The quality of forages on rangelands declines as the concentration of plant tissue nitrogen and protein decreases [24]. A rise in the atmospheric CO<sub>2</sub> also affects the primary productivity of rangelands [25]. In Africa, most grassy rangelands have been taken over by shrubs and trees which results in bush encroachment, a phenomenon that tends to increase plant biomass but limits the productivity of rangelands [26]. A study on 37 selected sites of the Mongolian rangelands to understand the impact of climate change on the productivity of these rangelands, showed that the Mongolian Steppe was already affected by climate change leading to marginal increase in the soil organic carbon level, and resulting in reduction of the productivity of forage plants [27].

### **Effect of climate change and variability on rangeland quality and productivity Length of growing period (LGP)**

Climate change has either lengthened or shortened the length of the growing season in various regions globally. In West Africa, areas around the west Guinean coast are reported to experience about 20 % decrease in the length of the growing period whereas regions around the east are expected to experience an increase of about 5-10 % [27]. The central and eastern Sahel areas of Africa are to experience over 30 % increase in the length of the growing seasons while the western Sahel regions will experience shorter growing seasons [28]. Most parts of the sub-Saharan Africa will experience a decrease in the length of growing period as a result of the combined impact of changes in temperature and rainfall, whereas the decrease will be severe in some cases [20].

The length of the growing season has an impact on the quality and quantity of forages as well as the duration and period when live forages are accessible [27]. The semi-arid rangelands of the Sahel have been affected by climate change (high temperatures and recurrent droughts) which has led to a higher ratio of actual to potential evapotranspiration. This has limited plant growth and reduced the length of the growing season [28]. A reduction in the length of the growing season translates into lower forage yields and shorter periods of availability of lush pasture for ruminant livestock.

In the temperate regions, increasing temperatures will affect the length of growing season through reduced cold stress and increased heat loads during summer [37]. It is projected that by 2050, most parts of the world will possibly have an increase



in temperature during the growing season than currently experienced, and this temperature increase will eventually be associated with more rainfall variability [29].

### **Drought stress in plants**

Rangeland quality and composition can be affected by the interactions existing among temperature, rainfall, atmospheric CO<sub>2</sub> concentration as well as land cover change and grazing. For example, drought stress in plants is increased when there is an increase in temperature [20]. Prolonged drought is associated with a comparable reduction in the net annual productivity of rangelands. The survival of long-lived perennial tree and pasture species is threatened by severe droughts. Browse species have high chance of survival than grasses during drought periods as a result of the effect of climate change and as such, farmers with large ruminants such as cattle are forced to shift from cattle production to the production of small ruminants such as goats and sheep since they require less feed and water [11]. Drought stress leads to physiological and morphological changes in forage plants. This is a natural response to avoid water deficit and heat stress. This response ultimately reduces the nutritional value and digestibility of the forage plants.

### **Changes in plant species composition**

Global average surface temperature is predicted to increase between 1.8°C to 4.0°C by the year 2100 according to IPCC [2]. Approximately 20 to 30 % of some plant and animal species are reportedly at the risk of becoming extinct with an increase of 1.5°C to 2.5°C [30]. This will lead to severe outcomes for food security in developing countries. Plant species composition is influenced by climate change mainly by shifting the amount, seasonal pattern, and vertical placement of soil water which will turn out to affect the outcome of competition by plants [31]. A rise in CO<sub>2</sub> concentration contributes to changes in composition of species by lowering transpiration to prolong soil water availability within a growing season [32]. The current CO<sub>2</sub> level of 350ppm in the atmosphere is projected to rise to 700ppm by 2100 [30]. This projected rise may tend to favour C<sub>3</sub> shrubs and legumes over C<sub>4</sub> grasses, consequently leading to changes in plant species composition. A study on plant species diversity in South Africa revealed that 11 % of the species investigated were at risk of extinction [33]. It is also reported that, climate change will result in the extinction of over one million terrestrial species in the next five decades [34]. Loss in plant species biodiversity is said to result from increase in temperature and prolonged droughts creating the right conditions for intensive rangeland fires. These fires together with changes in hydrologic and life cycles are expected to alter the habitats that support vulnerable plant species, thus leading to





loss of species or changes in species composition in favor of more resilient species [35].

### Impact on plant nutritional value

Forage quality is affected by rise in CO<sub>2</sub> and temperature as well as variations in precipitation, which complicates the quality of forages [27]. Forage quality can be increased by concentrating nitrogen moderately into plant tissues especially during the dry conditions and delaying maturity [36]. An increase in CO<sub>2</sub> tends to increase total shoot nonstructural carbohydrates and this adds up to improve forage quality [37]. This is because increase in CO<sub>2</sub> increases the rate of photosynthesis which converts the CO<sub>2</sub> into sugars and therefore, elevated CO<sub>2</sub> will stimulate the production of nonstructural carbohydrates. However, in dry and nitrogen limited rangelands, a rise in CO<sub>2</sub> reduces forage quality by increasing the ratio of carbon to nitrogen, lowering crude protein content, and finally increasing the fiber content of forage which will eventually decrease digestibility [38]. Studies have shown that nonstructural carbohydrate content of grasses increased by 25 % under elevated CO<sub>2</sub> levels but under the same condition, nitrogen content decreased by 8 % [39]. This reduction in the concentration of plant tissue nitrogen and protein will undoubtedly lead to a decline in forage quality in pastures and rangelands [24]. However, CO<sub>2</sub> elevation in the atmosphere is also expected to increase the population of C<sub>3</sub> legume species in rangelands, thus, compensating for the reduction in the concentration of plant tissue nitrogen and protein in elevated CO<sub>2</sub> levels [40]. C<sub>3</sub> plants respond to elevated CO<sub>2</sub> levels by increasing photosynthesis and reducing conductance in the stomata. This makes it possible for legumes to increase N<sub>2</sub> fixation, thus, reducing the impact of drought on N<sub>2</sub> fixation [41]. Other studies also suggest that, the response of legumes to elevated CO<sub>2</sub> is only in managed systems and not in natural systems [41]. This is because nutrient deficiency can limit the response of legumes to elevated CO<sub>2</sub> [42]. Phosphorous limited the response of legumes to elevated CO<sub>2</sub> in a N-poor calcareous grassland which subsequently limited N transfer to non-leguminous species in the grassland ecosystem [43].

Ozone (O<sub>3</sub>) acts as a shield to prevent harmful shortwave ultraviolet solar radiations from reaching the surface of the earth. However, the presence of O<sub>3</sub> on the earth's surface has added to the impact of GHGs to plants. Plants exposed to high concentrations of ground-level O<sub>3</sub> suffer from reduced photosynthetic activities, experience accelerated rate of senescence, reduced rate of plant growth and lower yields [44].



Lack of adequate soil water coupled with high atmospheric temperatures increase evapotranspiration, which results in increased lignification of forage species thereby reducing intake and digestibility by livestock that feed on them [45]. It has been reported that increase in temperature does not only increase drought stress in plants but also increases lignification of their tissues which affects both digestibility of plants as well as their rate of decomposition [20]. Lignin and cell wall contents in plants also increase as a result of a rise in temperature and this lowers digestibility. Neutral Detergent Fibre (NDF) is also said to increase with rise in temperature. It is reported that, NDF increases by 0.4 +/- 0.06 % for every 1°C rise in temperature [46], and the higher the NDF content, the lower the feed intake by ruminants. According to US Global Change Research Programme [24], temperatures are expected to increase by 2.0–3.08 °C by 2050 and 2.2– 5.58 °C by 2100. There is, therefore, no doubt that the grasses in rangelands are going to be less nutritious, so livestock that feed on them are going to gain less nutrients leading to their poor condition and a subsequent reduction in productivity.

### **Increase in vegetation flammability**

Increased temperatures coupled with lowered rainfall increases vegetation flammability [20]. Some rangeland grasses promote fire and are also by themselves promoted by fire, whereas the woody plant populations are sensitive to fire in their seedling stage but may be competitively superior once established. This makes rangelands dominated with woody plants less prone to fire than grasslands [47]. An increase in fuel load is expected under high levels of CO<sub>2</sub> since plant growth would be high. Temperature increases driven by climate change will cause areas with low rainfall to experience an increase in fuel dryness and a reduction in relative humidity [48]. With high amounts of fuel load and warmer environments, rangelands can experience wildfires caused by natural means (lightning) or by anthropogenic activities (hunting, grazing and burning of croplands).

In northern Australia's open woodlands, it is perceived that woodland thickening is as a result of reduction of pasture burning [49]. Continuous fire of annual grasses can cause a shift from grassland to shrublands and woodlands, whereas on a large scale change from shrublands to homogenous grasslands can affect evapotranspiration, moisture transfer and rainfall. The net effect could be an increase in aridity of the region [22].

Wildfires (driven by climate change) in rangelands may lead to the loss of forage biomass essential for livestock feeding. However, these wildfires can also clear



close woody canopies and enhance sunlight penetration into rangelands. This will enhance growth of herbaceous species and encourage plant diversity.

### **Mortality of plants**

When rangelands are exposed long enough to high temperatures, they undergo stress which eventually leads to a large-scale mortality of species in these rangelands. Some climatic events can lead to the death of some specific rangeland species [50]. A severe and prolonged drought can result in the mortality of perennial plants and cause a shift to an annual-dominated vegetation [51]. There has been considerable death of trees across the globe which is attributable to severe drought [52]. For example, in Africa there were tree mortalities of *Cedrus atlantica* in Belezma Park, Algeria and quiver tree (*Aloe dichotoma*) in Tirasberg mountains in Namibia [52]. Loss of species also affects the interactions with other species, thus, having an impact on the overall function of the ecosystem [35]. There is also the extinction of animal species like the checkerspot butterfly. This is as a result of an increase in the variability of precipitation leading to a reduction of temporal overlap between the larvae of the butterfly and its host, the dwarf plantain [53]. An increase in air temperature above the physiological limits has also led to the extinction of 48 lizards. The presence of all these organisms contribute to rangeland ecosystem function [54].

### **Effect of Climate Change on soil**

With the advent of climate change and variability, CO<sub>2</sub> content in the atmosphere is reported to have increased by some 30 % in the last fifteen decades [55]. Atmospheric CO<sub>2</sub> concentration is projected to rise to 800ppm by the end of the 21<sup>st</sup> Century [56]. With respect to the Carbon pool in the soil, it is also reported that the soil alone contains 75 % of the total carbon pool in the ground [57]. The soil serves as a means where plants and some microorganisms directly or indirectly secure their food, thus serving as carbon sink. The rise in CO<sub>2</sub> level in the atmosphere coupled with fluctuations in temperature and precipitation lead to alterations in important soil processes. This exposes the soil to physio-chemical degradation, salinization, erosion, reduction in water availability and storage of soil nutrients [57].

Rise in temperatures caused by climate change induces rapid decomposition of organic matter in the soil. This leads to loss of CO<sub>2</sub> from the soil and a reduction in soil organic carbon stock [57]. Generally, a rise in temperature will result in faster rate of soil organic matter decomposition, faster rate of mineralization and nitrification of Nitrogen [58]. The high rate of soil organic matter decomposition will be due to the fact that warmer temperatures stimulate microbial activities. This high



rate of decomposition will lead to loss of soil nutrients through leaching and erosion, thus reducing nutrient availability to forage plants [59]. A rise in temperature of about 0.3 to 6 °C increases the rate of Phosphorus mineralization in litter by 48 % [60]. Additionally, when the rate of mineralization is faster than rate of plant mineral uptake, the result is leaching. Extremely high temperatures and droughts result in evaporation of soil moisture, which in turn reduces the amount of dissolved nutrients for plants uptake from the soil [11].

Precipitation plays a vital role in the growth and development of plants. Precipitation which is expected to increase in some areas of the world by reason of climate change will make more soil moisture available to plants. Soil moisture will in turn make minerals available to plants. However, extreme precipitation depletes the nutrients in the soil through leaching and erosion through runoff water. It is estimated that an increase in precipitation by 1 % in a given area can lead to an increase in erosion by 1.5 – 2 % [56]. Soil moisture and fertility among others are the factors to consider during the production of rangelands. Therefore, any environmental alterations which affects the process by which soil water is attained will have a severe influence on the stability and productivity of forage plants [4].

Changes in the carbon to nitrogen ratio (C: N) of vegetation which comes as a result of climate change affect soil nutrients indirectly and causes changes in some characteristics of soil as well [61]. Response to these climate change drivers such as increasing CO<sub>2</sub>, temperature and rainfall tend to vary with different locations because of the difference in soil characteristics such as the water-holding capacity [62]. In addition, structures and functions of some aspects of the ecosystem will be affected due to the overall effect of these climate change drivers on the availability of soil water [31]. However, rangelands can be managed to help alleviate anthropogenic causes of climate change through a process known as carbon sequestration. This is the storage of carbon from the atmosphere into the soil and vegetation through photosynthesis, humification and aggregation [4] and store the carbon in organic forms, including plant biomass and soil organic matter. The process enhances soil fertility due to the increasing amounts of soil organic matter.

## CONCLUSION

Climate change no doubt has been and is still a topical issue across the globe. The Earth's climate has changed through natural events and now in an era where anthropogenic activities are immensely contributing to this phenomenon. Gases such as CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub> and CFC's collectively known as GHGs are responsible for the change in climate. Climate change is adversely affecting rangeland's quality



and productivity. High temperatures with little or no precipitation affect the length of growing periods of plants, nutritional value of forages, species composition, vegetation flammability, drought stress and mortality of range plant and animal species. Climate change also affects soil chemical and physical properties, leading to changes in the availability of soil water to plants. For sustainable livestock production, there is the need to fully understand the effect of climate change on rangelands. This will be helpful in the formulation of appropriate range management decisions to optimize forage production and utilization by livestock.

It is recommended that:

- Novel range management policies be enacted and implemented for sustainable rangeland use in the face of climate change.
- Scientists be properly resourced to effectively quantify the impact of climate change on rangeland resources and results communicated to the understanding of all including the resource poor livestock farmers.
- Vulnerable rural based livestock farmers should be assisted with adaptation strategies to remain in production.



## REFERENCES

1. **Muller C and SE Shackleton** Perceptions of climate change and barriers to adaptation amongst commonage and commercial livestock farmers in the semi-arid Eastern Cape Karoo. *African Journal of Range & Forage Science*, 2014; **31(1)**: 1-12.
2. **IPCC**. Intergovernmental Panel on Climate Change. Climate Change: Impacts, Adaptation and Vulnerability. Summary for policy makers, 2007.
3. **Trenberth KE** Climate change caused by human activities is happening and it already has major consequences, *Journal of Energy & Natural Resources Law*, 2018; **36(4)**: 463-481. <https://doi.org/10.1080/02646811.2018.1450895>
4. **Brown JR and J Thorpe** Climate Change and Rangelands: Responding Rationally to Uncertainty? *Rangelands*, 2008; **30(3)**: 3-6. <https://doi.org/10.1046/j.1365-2486.2002.00515.x>
5. **Food and Agriculture Organization of the United Nations**. The second report on the state of the world's plant genetic resources, 2010. <http://www.fao.org/agriculture/crops/core-themes/theme/seeds-pgr/sow/sow2/en/> Accessed on February 15<sup>th</sup>, 2013.
6. **Sietz D, Ludeke MKB and C Walther** Categorisation of typical vulnerability patterns in global drylands. *Global Environmental Change: Human and Policy Dimensions*, 2011; **21**:431–440.
7. **Parry L, Carter TR and NT Konijn** The Impact of Climatic Variations on Agriculture Volume 2: Assessments in Semi-Arid Regions. Kluwer, Dordrecht, The Netherlands, 1988.
8. **Olila DO and VW Oliver** Climate change, Savanna grassland, autoregressive model, time series data. 5<sup>th</sup> International conference of the African Association of Agricultural Economists. Sept 23-26, Addis Ababa, Ethiopia, 2016.
9. **IPCC TAR**. Climate Change: Impacts, Adaptation and Vulnerability. IPCC Third Assessment Report, Cambridge University Press, 2001.
10. **UNFCCC**. Climate change: Impacts, vulnerabilities and adaptation in developing countries, 2007.



11. **Timpong-Jones EC, Adogla-Bessa T, Naazie A and LK Adjorlolo** Climate change: implications for the resource poor livestock farmers in Africa. In: Yangyouru M, Naazie A, MacCarthy DS, Nkansah GO and SK Torkpo (eds.). *Agricultural Research for Food Security and Sustainable Livelihood: University of Ghana Readers Series*. (Ch. 14, pp. 231-248, ISBN: 978-9988-2-0787-8). Tema, Ghana: Digibooks Ghana Ltd, 2014.
12. **FAO**. Livestock's long shadow- environmental issues and options FAO 2009. *The state of food and agriculture. Livestock in the balance*. Rome, Italy: FAO, 2006; 168 p.
13. **IPCC (Intergovernmental Panel on Climate Change)**. Climate change 2001: technical summary: climate change 2001: impacts, adaptation, and vulnerability. A report of Working Group II of the Intergovernmental Panel on Climate Change. Geneva: IPCC, 2001.
14. **IPCC**. Climate Change 2014: Synthesis Report. In: Core Writing Team, Pachauri, R.K. and L A Meyer (Eds), Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland, 2014.
15. **IPCC**. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.
16. **Hughes L** Biological consequences of global warming: Is the signal already apparent. *Trends Ecological Evolution*, 2000; 15: 56–61.
17. **Wiens JJ** Climate-Related Local Extinctions Are Already Widespread among Plant and Animal Species. *PLOS Biology*, 2016; **14(12)**: e2001104  
<https://doi.org/10.1371/journal.pbio.2001104>
18. **UNEP (United Nations Environmental Programme) and GRID-Arendal**. Vital Climate Graphics Africa, 2011.
19. **Holechek J, Pieper RD and CH Herbel** Range management: principles and practices. 5th ed. Upper Saddle River, NJ, USA: Prentice Hall, 2004; 587 p.
20. **Hoffman T and C Vogel** Climate Change Impacts on African Rangelands. *Rangelands*, 2008; **30(3)**:12-17.

21. **Thornton PK, Jones PG, Alagarswamy G and J Andresen** Spatial variation of crop field response to climate change in East Africa. *Global Environmental Change*, 2009; **19**:54–65.
22. **Chambers JC and P Pellant** Climate Change Impacts on Northwestern and Intermountain United States Rangelands. *Rangelands*, 2008; **30**: 29-33.
23. **Taub D** Effects of Rising Atmospheric Concentrations of Carbon Dioxide on Plants. *Nature Education Knowledge*, 2010; **3(10)**:21.
24. **US Global Change Research Program.** Global climate change impacts in the United States. Karl T R, Melillo J M and T C Peterson [EDS.]. Cambridge, UK: Cambridge University Press, 2009. Available at: <http://www.globalchange.gov/usimpacts> Accessed 5 August 2013.
25. **Bolortsetseg B** Climate change Vulnerability and Adaptation in the Livestock Sector of Mongolia. Assessments of Impacts and Adaptations to Climate Change (AIACC) final report. Ulaanbaatar, 2006; 44 pp.
26. **Cook KH and EK Vizy** Impact of climate change on mid-twenty-first century growing seasons in Africa. *Clim. Dyn.*, 2012; **39**:2937-2955.
27. **Polley HW, Briske DD, Morgan JA, Wolter K, Bailey DW and JR Brown** Climate Change and North American Rangelands: Trends, Projections, and Implications. *Rangeland Ecology and Management*, 2013; **66**:493–511 | September 2013. <https://doi.org/10.2111/REM-D-12-00068.1>
28. **Le Houérou HN, Bingham RL and W Skerbek** Relationship between the variability of primary production and the variability of annual precipitation in world arid lands. *Journal of Arid Environments*, 1988; **15**:1-18.
29. **Battisti D and RL Naylor** Historical warnings of future food insecurity with unprecedented seasonal heat. *Science*, 2009; **323**:240–244.
30. **FAO.** Wildlife in a changing climate. FAO Forestry Paper, Rome, Italy, 2012; 118 p.
31. **Knapp AK, Beier C, Briske DD, Classen AT, Luo Y, Reichstein M, Smith M D, Smith SD, Bell JE, Fay PA, Heisler JL, Leavitt SW, Smith RB and E Weng** Consequences of more extreme precipitation regimes for terrestrial ecosystems. *BioScience*, 2008; **58**:811–821.



32. **Polley HW, Jin VL and PA Fay** CO<sub>2</sub>-caused change in plant species composition rivals the shift in vegetation between mid-grass and tallgrass prairies. *Global Change Biology*, 2012; **18**:700–710.
33. **Midgley GF, Hannah L, Millar D, Rutherford MC and LW Powrie** Assessing the vulnerability of species richness to anthropogenic climate change in a biodiversity hotspot. *Global Ecology and Biogeography*, 2002; **11**: 445-451.
34. **Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC and BFN Erasmus** Extinction risk from climate change. *Nature*, 2004; **427**:145-148.
35. **Sintayehu DW** Impact of climate change on biodiversity and associated key ecosystem services in Africa: a systematic review. *Ecosystem Health and Sustainability*, 2018; **4(9)**:225-239.
36. **Craine JM, Town EG, Joern A and RG Hamilton** Consequences of climate variability for the performance of bison in tallgrass prairie. *Global Change Biology*, 2009; **15**:772–779.
37. **Lilley JM, Bolger TP, Peoples MB and RM Gifford** Nutritive value and the nitrogen dynamics of *Trifolium subterraneum* and *Phalaris aquatic* under warmer, high CO<sub>2</sub> conditions. *New Phytologist*, 2001; **150**:385–395.
38. **Jin L, Wang Y, Iwasa AD, Xu Z, Schellenberg MP, Zhang YG, Liu XL and TA McAllister** Effect of condensed tannins on ruminal degradability of purple prairie clover (*Dalea purpurea* Vent.) harvested at two growth stages. *Animal Feed Science and Technology*, 2012; **176**: 17-25.
39. **Dumont B, Andueza D, Niderkorn V, Luscher A, Porqueddu C and C Picon-Cochard** A meta-analysis of climatic change effects on forage quality in grasslands: specificities of mountain and Mediterranean areas. *Grass Forage Sci.*, 2015; **70**:239-254.
40. **Uddin ME and E Kebreab** Review: Impact of Food and Climate change on pastoral industries. *Frontiers in Sustainable Food Systems*, 2020; 4: Article 543403.
41. **Rogers A, Ainsworth E and ADB Leakey** Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiol.*, 2009; **151(3)**: 1009-1016.



42. **van Groenigen KJ, Six J, Hungate BA, de Graaff MA, van Breemen N and C van Kessel** Element interactions limit soil carbon storage. *Proc Natl. Acad. Sci. USA*, 2006; **103**:6571–6574.
43. **Niklaus PA and C Korner** Synthesis of a six-year study of calcareous grassland responses to in situ CO<sub>2</sub> enrichment. *Ecol Monogr*, 2004; **74**:491-511.
44. **Gonzalez-Fernandez I, Bass D, Muntifering R, Mills G and J Barnes** Impact of ozone pollution on productivity and forage quality of grass/clover swards. *Atmos. Environ.*, 2008; **42**: 8755-8769.
45. **Ferraretto LF and RD Shaver** Effects of whole-plant corn silage hybrid type on intake, digestion, ruminal fermentation and lactation performance by dairy cows through a meta-analysis. *Journal of Dairy Science*, 2015; 98:2662–2675. <https://doi.org/10.3168/jds.2014-9045>
46. **Lee MA, Davis AP, Chagunda MGG and P Manning** Forage quality declines with rising temperatures, with implications for livestock production and methane emissions. *Biogeosciences*, 2017. **14**:1403–1417. <https://doi.org/10.5194/bg-14-1403-2017>
47. **Westoby M, Walker BH and I Noy-Meir** Opportunistic management for rangelands not at equilibrium. *Journal of Range Management*, 1989; **42**:266-274.
48. **Howden SM, Moore JL, McKeon GM, Reyenga PJ, Carter JO and JC Scanlan** Global Change Impacts on Fire Dynamics in the Mulga Woodlands of South-West Queensland Working Paper Series 99/ 05. CSIRO Wildlife and Ecology, Canberra, 1999.
49. **Burrows WH, Henry BK, Back PV, Hoffman MB, Tait LJ, Anderson ER, Menke N, Danaher T, Carter JO and GM Mckeon** Growth and carbon stock change in eucalpt woodlands in northeast Australia: Ecological and greenhouse sink implications. *Global Change Biology*, 2002; **8(8)**:769-784. <https://doi.org/10.1046/j.1365-2486.2002.00515.x>
50. **Neilson RP** High-resolution climatic analysis and Southwest biogeography. *Science*, 1986; 23227-34.

51. **Hein AL** Climate change in Africa. Cicerone, 2006; (6).  
<http://www.cicero.uio.no/fulltext/index.aspx?id=5249&lang=NO> Accessed 6/05/2020.
52. **Allen CD, Macalady AK, Chenchouni H, Bachelet D, Mcdowell N, Vennetier M, Kitzberger T, Rigling A, Breshears DD, Hogg EH, Gonzalez P, Fensham R, Zhang Z, Castro J, Demidova N, Lim JH, Allard G, Running SW, Semerci A and N Cobb** A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 2010; **259**:660–684.
53. **McLaughlin JF, Hellmann JJ, Boggs CL and PR Ehrlich** Climate change hastens population extinctions. *Proc. Natl Acad. Sci. USA*. 2002; **99**:6070–6074. <https://doi.org/10.1073/pnas.052131199>
54. **Sinervo B, Méndez-de-la-Cruz F, Miles DB, Heulin B, Bastiaans E, Villagrán-Santa CM, Lara-Resendiz R, Martínez-Méndez N, Calderón-Espinosa ML, Meza-Lázaro RN, Gadsden H, Avila LJ, Morando M, De la Riva IJ, Sepulveda PV, Rocha CFD, Ibargüengoytia N, Puntriano CA, Massot M, Lepetz V, Oksanen TA, David G Chapple DG, Bauer AM, Branch WR, Jean Clobert J and JW Sites** Erosion of lizard diversity by climate change and altered thermal niches. *Science*, 2010; **328**:894– 899. <https://doi.org/10.1126/science.1184695>
55. **Shukla SK, Shee S, Maity S, Shukla SK, Shee S, Maity SK, Solomon S, Awashi SK, Gaur A, Pathak AD and VP Jaiswal** Soil carbon sequestration and crop yields in rice-wheat and sugarcane-ratoon-wheat cropping systems through crop residue management and inoculation of *Trichoderma viride* in Subtropical India. *Sugar Tech.* ,2016; **19**:347-358.
56. **Elbasiouny H, El-Ramady H, Elbehiry F, Rajput VD, Minkina T and S Mandzhieva** Plant Nutrition under Climate Change and Soil Carbon Sequestration. *Sustainability*, 2022;**14**(2): 914.
57. **Sarfraz R, Hussain A, Sabir A, Ben Fekih I, Ditta A and S Xing** Role of biochar and plant growth promoting rhizobacteria to enhance soil carbon sequestration—A review. *Environmental Monitoring and Assessment*, 2019;**191**(4):1-13.

58. **Cookson WR, Osman M, Marschner P, Abaye DA, Clark I, Murphy DV, Stockdale EA and CA Watson** Controls on soil nitrogen cycling and microbial community composition across land use and incubation temperature. *Soil Biology and Biochemistry*, 2007; **39(3)**:744-756.
59. **Lotze-Campe H** Climate Change, Population Growth, and Crop Production: An Overview. In *Crop Adaptation to Climate Change*, 1st ed.; Yadav, S.S., Redden, R.J., Hatfield, J.L., Lotze-Campen, H., Hall, A.E., Eds.; John Wiley & Sons, Ltd.: Hoboken, NJ, USA, 2011.
60. **Wrage N, Chapius-Lardy L and J Isselstein** Phosphorus, plant biodiversity and climate change. In: *Sociology, Organic Farming, Climate Change and Soil Science. Sustainable Agriculture Reviews 3*, Springer: Dordrecht, The Netherlands, 2009.
61. **IPCC**. Guidelines for national greenhouse gas inventories, 1996.
62. **Polley HW, Briske DD, Morgan JA, Wolter K, Bailey DW and JR Brown** Climate Change and North American Rangelands: Trends, Projections, and Implications. *Rangeland Ecological Management*, 2013; **66**:493–511.  
<https://doi.org/10.2111/REM-D-12-00068.1>