



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

1142
S080i

UNITED STATES DEPARTMENT OF AGRICULTURE
AGRICULTURAL ADJUSTMENT AGENCY

OIL and MEAL YIELDS PER ACRE from COTTONSEED, PEANUTS & SOYBEANS

A study of farms, counties and areas producing cotton and peanuts or cotton and soybeans, Southern Region, AAA, 1942

I. W. DUGGAN, Director
C. D. WALKER, Assistant Director
SOUTHERN DIVISION

WASHINGTON, D.C. DECEMBER, 1943

CONTENTS

| | Page |
|--|------|
| Introduction | 1 |
| Scope of Study and Source of Data | 1 |
| Method of Analysis | 2 |
| Limitations of Study | 4 |
| Peanuts Versus Cotton for Oil Production. | 4 |
| Area Comparisons | 5 |
| Alabama | 7 |
| Arkansas | 9 |
| Florida | 10 |
| Georgia | 11 |
| Louisiana | 12 |
| Mississippi | 13 |
| Oklahoma | 14 |
| South Carolina | 15 |
| Texas | 16 |
| Peanuts Versus Cottonseed for Meal Production | 18 |
| Soybeans Versus Cotton for Oil Production | 19 |
| Soybeans Versus Cottonseed for Meal Production | 20 |

MAP

Areas and Counties Included in Study of Oil and Meal Yields From Cottonseed, Peanuts, and Soybeans Facing 2

TABLES

| No. | | Page |
|-----|--|------|
| 1 | Yield of oil and meal per 100 pounds of seed from peanuts, soybeans, and cottonseed | 21 |
| 2 | Factors applied to peanut, soybean, and cotton lint yields to convert such yields to oil and meal yields per acre | 22 |
| 3 | Comparative data on oil yields per acre from cottonseed and peanuts, by areas, Southern Region, 1942 | 23 |
| 4 | Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas, Southern Region, 1942 | 24 |
| 5 | Comparative data on oil yields per acre from cottonseed and peanuts, by States, Southern Region, 1942 | 26 |
| 6 | Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by States, Southern Region, 1942 | 27 |
| 7 | Rank of Southern Region States by various factors related to oil yields per acre from cottonseed and peanuts, 1942. | 28 |
| 8 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Alabama Counties, 1942 | 30 |
| 9 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Alabama, 1942. | 31 |
| 10 | Relative advantage of peanuts over cottonseed in per acre oil production, by areas in Alabama, 1942 | 32 |
| 11 | Frequency distribution of farms by ratio of peanut oil yield per acre to cottonseed oil yield per acre, by areas in Alabama, 1942. | 33 |
| 12 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Arkansas Counties, 1942 | 34 |
| 13 | Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas in Arkansas, 1942 | 35 |
| 14 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Florida Counties, 1942 | 36 |
| 15 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, selected Florida Counties, 1942 | 37 |
| 16 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Georgia Counties, 1942 | 38 |
| 17 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Georgia, 1942. | 39 |
| 18 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Louisiana Parishes, 1942. | 40 |
| 19 | Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas in Louisiana, 1942. | 41 |
| 20 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Mississippi Counties, 1942 | 42 |
| 21 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Mississippi, 1942 | 43 |
| 22 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Oklahoma Counties, 1942 | 44 |
| 23 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Oklahoma, 1942 | 45 |
| 24 | Comparative data on oil yields per acre from cottonseed and peanuts, selected South Carolina Counties, 1942 | 46 |
| 25 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, selected South Carolina Counties, 1942 | 47 |
| 26 | Comparative data on oil yields per acre from cottonseed and peanuts, selected Texas Counties, 1942 | 48 |
| 27 | Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Texas, 1942 | 49 |
| 28 | Relative advantage of peanuts over cottonseed in per acre oil production, by areas in Texas, 1942 | 50 |
| 29 | Frequency distribution of farms by ratio of peanut oil yield per acre to cottonseed oil yield per acre, by areas in Texas, 1942 | 51 |
| 30 | Comparative data on meal yields per acre from cottonseed and peanuts, by areas, Southern Region, 1942 | 53 |
| 31 | Comparative data on meal yields per acre from cottonseed and peanuts, by States, Southern Region, 1942. | 54 |
| 32 | Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by States, Southern Region, 1942. | 55 |
| 33 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Alabama Counties, 1942 | 56 |
| 34 | Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, by areas in Alabama, 1942 | 57 |
| 35 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Arkansas Counties, 1942. | 58 |
| 36 | Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, by areas in Arkansas, 1942 | 59 |
| 37 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Florida Counties, 1942 | 60 |
| 38 | Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, selected Florida Counties, 1942 | 61 |
| 39 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Georgia Counties, 1942 | 62 |
| 40 | Frequency distribution of meal yields per acre from cottonseed and peanuts, by areas in Georgia, 1942 | 63 |
| 41 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Louisiana Parishes, 1942 | 64 |
| 42 | Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Louisiana, 1942 | 65 |
| 43 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Mississippi Counties, 1942. | 66 |
| 44 | Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Mississippi, 1942. | 67 |
| 45 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Oklahoma Counties, 1942. | 68 |
| 46 | Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Oklahoma, 1942. | 69 |
| 47 | Comparative data on meal yields per acre from cottonseed and peanuts, selected South Carolina Counties, 1942. | 70 |
| 48 | Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, by areas in South Carolina, 1942 | 71 |
| 49 | Comparative data on meal yields per acre from cottonseed and peanuts, selected Texas Counties, 1942. | 72 |
| 50 | Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Texas, 1942. | 73 |
| 51 | Comparative data on oil yields per acre from cottonseed and soybeans, State and area summary, Southern Region, 1942 | 75 |
| 52 | Relative advantage of soybeans over cottonseed in per acre oil production, by areas in Arkansas and Mississippi, 1942. | 76 |
| 53 | Frequency distribution of farms by ratio of soybean oil yield to cottonseed oil yield per acre, by areas in Ark. and Miss., 1942 | 77 |
| 54 | Comparative data on oil yields per acre from cottonseed and soybeans, selected Arkansas Counties, 1942. | 78 |
| 55 | Frequency distribution of farms by oil yield per acre from cottonseed and soybeans by sample counties, Arkansas Delta, 1942. | 79 |
| 56 | Comparative data on oil yields per acre from cottonseed and soybeans, selected Louisiana Parishes, 1942 | 80 |
| 57 | Frequency distribution of farms by oil yields per acre from cottonseed and soybeans, by areas in Louisiana, 1942 | 81 |
| 58 | Comparative data on oil yields per acre from cottonseed and soybeans, selected Mississippi Counties, 1942. | 82 |
| 59 | Frequency distribution of farms by oil yield per acre from cottonseed and soybeans, selected Mississippi Counties, 1942 | 83 |
| 60 | Comparative data on meal yields per acre from cottonseed and soybeans, State and area summary, Southern Region, 1942 | 85 |
| 61 | Comparative data on meal yields per acre from cottonseed and soybeans, selected Arkansas Counties, 1942 | 86 |
| 62 | Frequency distribution of farms by meal yield per acre from cottonseed and soybeans, by sample counties, Arkansas Delta, 1942 | 87 |
| 63 | Comparative data on meal yields per acre from cottonseed and soybeans, selected Louisiana Parishes, 1942 | 88 |
| 64 | Frequency distribution of farms by meal yields per acre from cottonseed and soybeans, by areas in Louisiana, 1942 | 89 |
| 65 | Comparative data on meal yields per acre from cottonseed and soybeans, selected Mississippi Counties, 1942 | 90 |
| 66 | Frequency distribution of farms by meal yields per acre from cottonseed and soybeans, selected Mississippi Counties, 1942 | 91 |

NOTE: Pages 29, 52, 74, and 84 have been left blank so that related tables face each other in the report.

OIL AND MEAL YIELDS PER ACRE FROM COTTONSEED, PEANUTS, AND SOYBEANS 1/

INTRODUCTION

Military events that followed the attack on Pearl Harbor made it necessary for United States farmers to expand tremendously their acreages of peanuts and soybeans. Farmers set all-time records with these crops in 1942, expanded them further in 1943, and are expected to plant still greater acreages in 1944. Farmers were encouraged to expand their production of peanuts and soybeans primarily to obtain much needed oil that we were no longer able to import. Goals - State, county, and individual farm - were established throughout the Southern States in 1942, not only in areas where peanuts and soybeans are commonly grown, but in areas and on farms where these two crops had not been grown extensively in recent years. Proof that farmers as a whole did an excellent job in 1942 is revealed in the production records.

Farmers have done exceedingly well; they have planted what their Government asked them to plant. They have planted peanuts for oil; they have planted soybeans for oil; and they have planted cotton, which also produces oil. Each of the three crops produces high protein meals also. In addition there is the cotton lint, hulls, and linters from the cotton crop and hay from the peanut crop.

Every Southern farmer cannot grow peanuts or soybeans. Not all of them can grow cotton. Many can grow cotton and one of the other two, but few can produce successfully all three of these oil-bearing crops.

This study was made to show for specific areas the comparative advantage of producing cottonseed or peanuts and cottonseed or soybeans for oil and meal. Cotton lint, the most important product of the cotton plant, has been omitted from the present analysis, as well as cottonseed hulls and linters; also peanut hay from the peanut crop.

Scope of Study and Source of Data

The study includes 149 sample counties from 9 cotton-producing States, i.e., Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Oklahoma, South Carolina, and Texas (see map facing page 2). These 9 States also grow peanuts and soybeans, but only three, Arkansas, Louisiana, and Mississippi, produce a significant acreage of soybeans.

From each major production adjustment area producing cotton, and either peanuts or soybeans, one or more representative counties was selected for study. Data were then taken from AAA records for all farms, but not

1/ This study was made by John E. Mason under the direction of F. H. Whitaker, Chief, Economic and Statistical Section, Southern Division, AAA. Ocie Coston assisted in planning the study. The statistical sections of the nine State AAA offices in the Southern Region and the employees of the 149 county AAA offices in which farm yield data were tabulated are due special acknowledgement for their assistance.

exceeding a total of 300 farms per county in most States, growing cotton and either peanuts or soybeans. Per farm yield data were tabulated for cotton, peanuts, and soybeans. The information is confined to the 1942 crop year because this is the only year for which representative data are available throughout the Southern States for all three crops (1943 data will be available shortly).

Oil and meal yields per 100 pounds of cottonseed are based on the total quantity of seed crushed and the amount of oil and meal produced, August 1942 through July 1943, as reported by the Bureau of the Census. Oil and meal yields per 100 pounds of seed from peanuts are based on Table 5 of the March 1943 issue of the Fats and Oils Situation, Bureau of Agricultural Economics, U. S. Department of Agriculture. Oil and meal yields per bushel of seed from soybeans are based on information furnished the Southern Division, AAA by mills that crushed the 1942 crop of soybeans produced in the Southern States (tables 1 and 2).


Method of Analysis

From appropriate AAA records showing acreage and production on individual farms for cotton, peanuts, and soybeans, the county AAA offices listed average 1942 yields for each of the three crops for a sample number of farms growing cotton and one or both of the other two oil-bearing crops. The county AAA offices in a majority of the States were instructed to list the data for 300 farms or for all farms growing cotton and one of the other oil-bearing crops, whichever number was the smaller. The county AAA offices were further instructed to place a check mark (\checkmark) by any yield figure considered to be unreliable.

Next, the county tabulations were reviewed and edited in the State AAA offices by the AAA statistician and, in 6 of the 9 States, by a member of the Washington staff. All zero yields and obviously incorrect yields were deleted from the study through the editing process. Only those farms with a yield figure for cotton and one or both of the other crops were retained in the study.

After the editing had been completed, a per acre oil and meal outturn was computed for each crop on each farm by applying the appropriate factors given in table 2. From here on the statistical analysis is apparent from the tables presented herein, with the exception of the area and State totals. In the case of peanut and cottonseed comparisons, in order to give proper weight to the counties having more than 300 farms, but which were limited to 300 farms in taking the sample, area yields for cotton lint and peanuts were computed by weighting the county yields by the harvested peanut acreage in the counties included in the sample. The State figure was arrived at by weighting the area averages thus obtained by the respective harvested peanut acreages for the entire area, including counties in and out of the sample. Practically all farms growing peanuts in the areas included in the study also produce cotton but all farms growing cotton do not produce peanuts, therefore,

AREAS AND COUNTIES INCLUDED IN STUDY OF OIL AND MEAL
YIELDS FROM COTTONSEED, PEANUTS, AND SOYBEANS
(AREA BOUNDARIES ADJUSTED TO COUNTY LINES)



AREA NAMES

- | | |
|-----------------------------|--------------------------------------|
| 1-HIGH PLAINS | 12-COAST PRAIRIE |
| 2-TRANS-PECOS | 13-COASTAL PLAIN |
| 3-RIO GRANDE PLAIN | 14-OZARK-OUACHITA HIGHLANDS |
| 4-EDWARDS PLATEAU | 15-MISSISSIPPI RIVER ALLUVIAL LAND |
| 5-ROLLING PLAINS | 16-COAST MARSH LAND |
| 6-OKLAHOMA CENTRAL PRAIRIES | 17-BROWN LOAMS |
| 7-OSAGE HILLS | 18-SAND-CLAY HILLS |
| 8-EAST OKLAHOMA PRAIRIES | 19-BLACK BELT |
| 9-CROSS TIMBERS | 20-UPPER COASTAL PLAIN |
| 10-GRAND PRAIRIE | 21-LIMESTONE BASIN |
| 11-TEXAS BLACKLANDS | 22-SAND MOUNTAIN |
| | 23-APPALACHIAN HIGHLANDS |
| | 24-BLUE RIDGE MOUNTAINS |
| | 25-PIEDMONT PLATEAU |
| | 26-FALL LINE SAND HILLS |
| | 27-COASTAL PLAIN-RED BELT |
| | 28-MIDDLE COASTAL PLAIN |
| | 29-LOWER COASTAL PLAIN |
| | 30-GULF COAST FLATWOODS |
| | 31-ROLLING SANDY LANDS AND FLATWOODS |
| | 32-HIGH SANDS AND FLATWOODS |
| | 33-EVERGLADES |
| | 34-BIG CYPRESS AREA |

in order to obtain area averages for farms producing both crops, harvested peanut acreages were used for weights when comparing cotton and peanuts. For soybean and cottonseed comparisons it made little difference which weight was used because of the more uniform distribution of the two crops throughout the areas studied; therefore, cotton yields were weighted by cotton acreages harvested and soybean yields by soybeans for beans acreages. 2/

Limitations of Study

It is important to recognize certain limitations in the data contained in this report. First, the yield information is for one crop-year only. It was not possible to obtain yield data for all three crops, or any two of them, farm by farm for a longer period.

Second, peanuts were grown in new areas in 1942, and the yields are not necessarily indicative of the adaptability of the soil or of the farmers' ability to grow the crop satisfactorily. In some cases high yields were obtained on small acreages; in many other cases low yields prevailed because the farmers did not know the best cultural practices. For these reasons, one should be careful in concluding that a new area is or is not adapted to peanut production.

Third, the factors used to convert individual farm yields of cotton lint, peanuts, and soybeans to oil and meal yields per acre are based on State or area averages. The counties, and most certainly individual farms, could not be expected to produce oil-bearing crops with a uniform oil content in all parts of a State. Nevertheless, in computing oil and meal outturn it was necessary to apply uniform factors to the per acre yields by areas or States.

Fourth, the area and State totals are not necessarily comparable with other totals for the same areas or States. The data presented herein apply to farms on which cotton and either peanuts or soybeans were grown - not to all cotton farms, all peanut farms, or all soybean farms in a State. The State figures, therefore, must be expected to vary from published figures on cotton, peanut, or soybean yields by States.

I. PEANUTS VERSUS COTTON FOR OIL PRODUCTION

In all but 3 of the 136 counties in which peanuts and cottonseed are compared in this study, peanuts exceeded cotton in the quantity of oil produced per acre in 1942. One of these counties was Hale, in the Black Belt of Alabama; the other two, Holmes and Simpson, are located in Mississippi. On a county basis the ratio of peanut oil yield per acre to cottonseed oil yield per acre ranged from 88 percent in Hale County, Alabama to 1,129 percent in Atascosa County, Texas. For the entire sample of 23,707 farms, peanuts produced 3-1/2 times as much oil per acre as cotton.

The ratio of peanut oil to cottonseed oil yield per acre is a relative comparison and does not indicate the absolute quantity produced. The pounds 2/ Arnold J. King, Statistical Laboratory, Bureau of Agricultural Economics, gave helpful advice in choosing the method of weighting.

produced per acre on farms growing both cotton and peanuts are shown in this report by counties, areas, and States. On a State basis, Mississippi ranks first in oil outturn per acre from cottonseed but eighth from peanuts (table 7). Georgia heads the list of States in oil outturn per acre from peanuts, but Texas is at the top in the ratio of peanut oil yield to cottonseed oil yield per acre.

In all States and in nearly all counties a certain percentage of the farms produced more oil per acre from cottonseed than from peanuts, ranging from 4 percent in Florida to 45 percent in Mississippi and averaging 17 for the Southern Region.

Table 7 also ranks the 9 States of the Southern Region according to the man-labor requirements per acre from cotton and peanuts. With the exception of Arkansas, Louisiana, and Mississippi, State labor requirement data were used. In order to eliminate the effect of the high man-labor requirements in the Mississippi River Delta areas of these 3 States, where peanuts are not produced commercially, man-labor requirements for cotton and peanuts were taken from special State studies where the two crops are grown in competition (see footnotes to table 7).

Of the 9 States, South Carolina has the highest per acre man-labor requirements for cotton, and the lowest ratio of peanut labor to cotton labor. Peanuts require only 52 percent as much labor per acre in South Carolina as cotton, but in Oklahoma peanuts require 86 percent as much labor as cotton. The other States fall between these two percentages.

AREA COMPARISONS

The 9 States of the Southern Region of the AAA comprise two post-war planning regions of the U. S. Department of Agriculture. The regional post-war planning committee in each of these regions has prepared a map to show areas reasonably homogenous as to physical resources and character of problems arising from the use of those resources. The 9 States contain a total of 34 such areas. From 23 of these areas one or more representative counties was selected for this study. The other areas were omitted because either cotton or peanuts was not produced at all or in such small quantities that it was not feasible to include them. Some of the 23 areas fall wholly within a single State, insofar as this study is concerned, while others include parts of two or more States. Four areas fall wholly within Texas, two are entirely in Oklahoma, three in Alabama, one in Georgia, and one is confined to Florida. Twelve areas cut across two or more States.

Pearnuts and cotton compete throughout the 23 areas, but in point of farms producing the two crops, area 5 (Rolling Plains), area 9 (Cross Timbers), area 13 (Coastal Plain - South Central States), and area 28 (Middle Coastal Plain) are the most important areas.

The Sand Mountain area of Alabama led all areas in the yield of cotton lint per acre and also in the yield of peanuts. This is a relatively small area but the adjacent Limestone Basin area ranked second with cotton and with peanuts. For the Sand Mountain area, the computed oil outturn per acre from cottonseed is 126 pounds, compared with 254 pounds per acre from peanuts on the same farms. Comparable figures are 80 and 213 for the Limestone Basin.

The oil outturn from cottonseed ranges from 14 pounds per acre in the Rio Grande Plain to 126 pounds in the Sand Mountain area. The average for the Southern Region is 50 pounds per acre, based on this study of farms growing both cotton and peanuts. Using the 1942 average cottonseed yield for all farms the oil outturn would amount to about 80 pounds per acre. Large contiguous areas in the Southwest averaged less than 40 pounds of oil per acre from cottonseed. The Rolling Plains of Oklahoma and Texas, the Ozark-Cuachita Highlands of Arkansas and Oklahoma, and the Rolling Sandy Lands of Florida produced 40 to 49 pounds of oil per acre from cottonseed. The High Plains of Texas, the Black Belt, the Fall Line Sand Hills, and the Middle Coastal Plains of the Southeast produced from 50 to 59 pounds of oil per acre. The highest cottonseed oil yields came from nearly all parts of Mississippi and the northern parts of Alabama, Georgia, and South Carolina, with averages of 70 pounds or more per acre.

The computed oil outturn from peanuts ranged from 71 pounds per acre in the Black Belt to 254 pounds in the Sand Mountain area, averaging 175 pounds for the Southern Region. When mapped by broad geographical areas, the lowest oil yields from peanuts, under 100 pounds per acre, occur in the Brown Loams of Mississippi and Louisiana, the Black Belt of Alabama and Mississippi and the Appalachian Highlands of Alabama and Georgia. The Piedmont Plateau, the Sand-Clay Hills and Upper Coastal Plain of Alabama and Mississippi, and the Ozark-Cuachita Highlands produced from 100 to 124 pounds of oil per acre from peanuts. Some scattered areas averaged between 125 and 149 pounds of oil per acre. Large areas of Oklahoma and Texas produced from 150 to 174 pounds per acre. When averaged in with Oklahoma and Texas, parts of Arkansas and Louisiana are also covered by this yield range. The highest yields, above 175 pounds per acre, were in the Middle Coastal Plain of the Southeast, the northern part of Alabama, and the High Plains of Texas.

In none of the 23 areas did cottonseed average as much oil per acre as peanuts. Cottonseed came closest to peanuts in per acre oil outturn in the Brown Loam area, where peanuts exceeded cotton by only 18 percent. Three other areas, the Sand-Clay Hills, the Black Belt, and the Appalachian Highlands produced less than 1-1/2 times as much oil per acre from peanuts as from cottonseed. The Rio Grande Plain produced more than 10 times as much oil per acre from peanuts as from cottonseed. The Edwards Plateau, Rolling Plains, Grand Prairie, Cross Timbers, and Coastal Plain (South Central States) areas each produced more than 4 times as much oil per acre from peanuts as from cottonseed. The High Plains, Oklahoma Central Prairies, East Oklahoma Prairies, and the Middle Coastal Plain areas of the Southeast each produced between

3-1/2 and 4 times as much oil per acre from peanuts as from cottonseed. On farms producing both crops in 1942, this study indicates that the average for the Southern Region is 3-1/2 times as much oil per acre from peanuts as from cottonseed.

Peanuts excelled cottonseed in per acre oil yields in each of the 23 areas. Significantly, cottonseed excelled on a certain percentage of farms in every area; it was as low as 2 percent and as high as 48 percent by areas, for an average of 17 percent for the Southern Region (table 3).

In most areas it is possible to determine the typical oil yield per acre from cottonseed, but not so easy for peanuts. When the farms are set up in frequency distributions by oil yields per acre, there is nearly always a distinct modal group for cottonseed, but peanut oil yields per acre range from very low to very high, with a tendency to an even distribution among all class intervals (tables 4 and 6). With such a wide range and even distribution, an average yield of oil per acre from peanuts must be used with care because it typifies only a small percentage of all farms.

For all farms included in the study, 42 percent produced under 40 pounds of oil per acre from cottonseed, 64 percent under 60 pounds, 79 percent under 80 pounds, and 89 percent under 100 pounds. Fifty percent of the farms produced between 20 and 60 pounds of oil per acre from cottonseed (table 6).

For peanuts, 12 percent of the farms produced under 40 pounds of oil per acre, 20 percent under 60 pounds, 30 percent under 80 pounds, 39 percent under 100 pounds, 36 percent from 100 to 199 pounds, 18 percent from 200 to 299 pounds, 6 percent from 300 to 399 pounds, and 1 percent 400 pounds or more.

ALABAMA

The sample for Alabama included 3,875 farms in 16 counties selected to represent 8 major areas of the State. The 1942 peanut yield on these farms, when weighted out by harvested peanut acreage in the respective areas, indicates a yield of 593 pounds (table 8). This is 57 pounds below the State yield for all peanut farms. The cotton yield on the farms growing peanuts was also considerably lower than the State average for all cotton farms.

This study indicates that Alabama farmers growing both crops produced 48 pounds of oil per acre from cottonseed and 362 percent as much, or 174 pounds from peanuts. These are the weighted averages for the State, but it is significant that 22 percent of the farms surveyed produced more oil per acre from cottonseed than from peanuts.

Although Alabama ranks second among the 9 States in per acre oil outturn from peanuts, there is considerable variation from farm to farm, county to county, and area to area. On a county basis, the computed oil outturn per acre from peanuts in Hale County and in Lee County is only 57 pounds per acre, compared with 307 pounds in Cullman County. On an area basis, the Black Belt makes the poorest showing, with an average of 68 pounds of oil per acre. The Sand Mountain area produces more than 3-1/2 times as much per acre as does the Black Belt, or an average of 254 pounds. The old peanut area, the Middle Coastal Plain, is somewhat above the State average, with a computed oil outturn per acre of 198 pounds from peanuts.

Oil production from cottonseed ranges from 32 pounds per acre in Lee County to 128 pounds in Cullman County; it varies from 39 pounds in the Sand-Clay Hills to 126 pounds in the Sand Mountain area. The Sand Mountain area has higher yields for both crops than any other major area of the State.

Coffee County, in the Middle Coastal Plain, produces more than 5 times as much oil per acre from peanuts as from cottonseed; the area as a whole produces about 4-3/4 times as much from peanuts as from cottonseed. The next best area for peanuts, compared with cottonseed, is the Limestone Basin where both cotton and peanuts produce well but where peanuts produce 2.7 times as much oil per acre as cottonseed. In the Black Belt, the Piedmont Plateau, the Appalachian Highlands, and the Upper Coastal Plain, peanuts turn out only about 1.4 to 1.6 times as much oil per acre as can be expected from cottonseed, based on this study of 1942 yields. In these areas, 31 to 48 percent of the farms actually produced more oil per acre from cottonseed than from peanuts. However, in the Middle Coastal Plain and in the Limestone Basin only 2 percent of the farms had yields indicating that more oil per acre was produced from cottonseed than from peanuts.

Table 9 gives a frequency distribution of the 3,875 farms by the per acre oil yields from cottonseed and peanuts for each of the major areas. One of the striking features of this distribution is the narrow range of yields from cottonseed compared with the wide range from peanuts. A significant percentage of both crops yield less than 20 pounds of oil per acre. The outturn of oil from cottonseed rarely exceeds 200 pounds per acre, but a large percentage of the farms exceed this amount with peanuts. In some areas, it appears that oil yields per acre from peanuts are more or less evenly distributed from below 20 pounds to 300 pounds or more.

For the State as a whole 31 percent of the farms produced less than 40 pounds of oil per acre from cottonseed, compared with 16 percent for peanuts; 50 percent of the farms produced less than 60 pounds per acre from cottonseed, compared with 26 percent for peanuts; 79 percent produced less than 100 pounds from cottonseed, compared with 43 percent for peanuts. None of the farms exceeded 239 pounds of oil per acre from cottonseed, but 17 percent of them produced 240 pounds or more per acre from peanuts.

A special analysis was made of the per acre oil yields from peanuts and cottonseed in Alabama, to show by areas the percent of farms producing one, two, three, four, five, or six times as much oil per acre from peanuts as from cottonseed. In the first place, 22 percent produced more oil per acre from cottonseed than from peanuts. However, 11 percent produced 6 times as much from peanuts as from cottonseed, 15 percent 5 times or more, 21 percent 4 times or more, 31 percent 3 times or more, 49 percent 2 times or more, and 78 percent equal or better. These percentages varied by areas, as shown in table 10. In the Middle Coastal Plain, 36 percent of the farms produced 6 times or more as much oil per acre from peanuts as from cottonseed, 75 percent of the farms in this area did 3 times as well with peanuts compared with cottonseed. For further details see tables 10 and 11.

ARKANSAS

In Arkansas, 1,150 farms from 14 counties were included in the sample. For purposes of analysis the counties have been grouped into 2 major areas. The farms in the Ozark-Ouachita Highland area had a cotton lint yield in 1942 of 184 pounds and a peanut yield of 427 pounds. The Coastal Plains area had a cotton yield of 191 pounds and a peanut yield of 367 pounds. The weighted average cotton lint yield for the combined areas was 188 pounds and peanuts 396 pounds (table 12).

The computed oil outturn per acre from cottonseed ranged from a low of 38 pounds in both Montgomery County and Searcy County to a high of 86 pounds in Sharp County. The State average was 56 pounds per acre for farms growing both cotton and peanuts.

The computed oil outturn per acre from peanuts was only 92 pounds in Little River County, but averaged 152 pounds in Montgomery County, the highest county average. The Ozark-Ouachita Highland area had a per acre oil yield from peanuts of 128 pounds; the Coastal Plains 110 pounds; and the weighted average from the combined areas was 119 pounds.

Every county in both areas produced more oil per acre from peanuts than from cottonseed, ranging from about 1-1/2 times as much in Izard County to 4 times as much in Montgomery County. The weighted average for the areas of the State growing both crops was 2.1 times as much oil per acre from peanuts as from cottonseed.

In Arkansas, as in the other States, a certain percentage of the farms produced more oil per acre from cottonseed than from peanuts. For the competing areas in the State as a whole, 17 percent of the farms excelled with cottonseed, ranging from 1 percent in Faulkner County to 46 percent in Sharp County (table 12).

In the Ozark-Ouachita Highland area a large percentage of the farms produced 40 to 59 pounds of oil per acre from cottonseed (table 13). Similar results were obtained in the Coastal Plains area, and of course, the State figures would show about the same results. Peanut yields are such in both areas that approximately 10 to 12 percent of the farms fell in each of the following oil yield per acre groups: 40-59, 60-79, 80-99, 100-119, 120-139, and 140-159. A few farms yielded below 40 pounds of oil per acre from peanuts and the remainder were distributed all the way from 160 pounds to more than 400 pounds.

For all farms included in the sample, 32 percent produced less than 40 pounds of oil per acre from cottonseed, compared with 11 percent for peanuts; 63 percent and 22 percent, respectively, produced under 60 pounds of oil per acre from cottonseed and peanuts; 95 percent and 45 percent, respectively, produced less than 100 pounds (table 13).

FLORIDA

Four counties were included in the study from Florida and these have been grouped into 2 areas. Data were tabulated for 876 farms. The cotton lint yield averaged 151 pounds per acre and peanuts 535 pounds. Santa Rosa County had the highest average yields for both, 185 pounds per acre for cotton and 803 pounds for peanuts. Leon County had the lowest yields, 91 pounds per acre for cotton and 321 pounds for peanuts (table 14).

The computed oil outturn from cottonseed ranged from 25 pounds per acre in Leon County to 51 pounds in Santa Rosa County, averaging 42 pounds for those areas of the State growing the two crops.

The computed oil outturn per acre from peanuts amounted to 93 pounds in Leon County and 241 pounds in Santa Rosa County, averaging 159 pounds for all areas growing the two competing crops.

On a relative basis peanuts did best in Santa Rosa County where $4\frac{3}{4}$ times as much oil per acre was produced from peanuts as from cottonseed. Peanuts made the poorest relative showing in Suwannee County, but even here 3 times as much oil per acre came from peanuts as from cottonseed. The ratio of peanut oil yield per acre to cottonseed oil yield per acre for all areas included in the study was 379 percent.

Four percent of the farms produced more oil per acre from cottonseed than from peanuts, ranging from 1 percent in both Jackson and Santa Rosa Counties to 13 percent in Suwannee County.

For cottonseed a large percentage of the farms are concentrated around the oil yield group of 20 to 39 pounds per acre. Peanuts show only a slight tendency to fall around any particular yield, and range all the way

from below 20 pounds of oil per acre to over 400 pounds. Only 3 percent of the farms produced 100 pounds or more of oil per acre from cottonseed, but 77 percent of the farms exceeded this amount from peanuts; 36 percent produced 200 pounds or more of oil per acre from peanuts (table 15).

GEORGIA

Georgia counties have been grouped into 7 areas by the Southeast Regional Post-War Planning Committee, but only 5 of these are important in the production of peanuts. For this study 18 representative counties growing both peanuts and cotton were selected for analysis, from which data were tabulated for 4,054 farms. The weighted average cotton yield on these farms in 1942 was 203 pounds; the peanut yield 637 pounds (table 16). The 1942 State yields for all farms were: cotton, 240 pounds; peanuts, 610 pounds. Of the 5 areas growing both crops, the Piedmont Plateau had the highest cotton yields (245 pounds per acre), but the lowest peanut yields (347 pounds per acre). The area with the highest peanut yields, the Middle Coastal Plain, had relatively low cotton yields. By counties, cotton lint yields ranged from a low of 149 pounds per acre in Talbot County to 352 pounds per acre in Morgan County. Peanut yields averaged as low as 283 pounds per acre in Talbot County to as high as 900 pounds in Bulloch County.

Converted to oil, cottonseed on all farms producing the two crops would turn out about 55 pounds of oil per acre, compared with 191 pounds from peanuts. By areas, cottonseed shows up best in the Piedmont Plateau, while peanuts are outstanding in the Middle Coastal Plain. Peanuts produce almost 4 times as much oil per acre in the Middle Coastal Plain as cottonseed. The ratio is 3-1/2 times as much from peanuts as from cottonseed for all areas in the sample; about 1-1/2 times for the Piedmont Plateau. In Bulloch County and in Toombs County, the computed oil outturn for 1942 is about 5-1/2 times as much per acre from peanuts as from cottonseed.

In the Middle Coastal Plain 97 percent of the farms produced more oil per acre from peanuts than from cottonseed; in the Piedmont Plateau only 66 percent of the farms excelled with peanuts; the State average is 91 percent (table 16).

The frequency distribution (table 17) shows the largest percentage of farms in all but one area in the 40-59 pounds of oil per acre group for cottonseed. Peanut yields do not show nearly so strong a tendency to cluster around any particular yield. In the Middle Coastal Plain, the principal peanut area of Georgia, oil yields range from below 20 pounds to over 400 pounds per acre, without any pronounced tendency to concentrate around any yield figure in between. In other areas some slight concentration is noted. For example, in the Piedmont Plateau 67 percent of the farms had oil yields per acre from peanuts of 20 to 119 pounds, but a fraction of 1 percent in

this area exceeded 400 pounds. In the Fall Line Sand Hills 39 percent of the farms had oil yields of 100 to 159 pounds per acre.

For the State as a whole, 31 percent of the farms had oil yields per acre from cottonseed less than 40 pounds; 62 percent less than 60 pounds; 83 percent less than 80 pounds; 93 percent less than 100 pounds; and 7 percent 100 pounds or more. With peanuts, only 5 percent of the farms had oil yields of less than 40 pounds per acre; 10 percent less than 60 pounds; 16 percent less than 80 pounds; 23 percent less than 100 pounds; 41 percent from 100 to 199 pounds; 27 percent from 200 to 299 pounds; 8 percent from 300 to 399 pounds; and 1 percent 400 pounds or more (table 17).

LOUISIANA

In Louisiana, 1,302 farms from 9 counties were included in the study. The weighted average cotton lint yield for the areas growing cotton and peanuts was 171 pounds; the peanut yield 306 pounds.

The computed oil outturn per acre from cottonseed ranged from 32 pounds in Caddo Parish to 75 pounds in Rapides Parish, on farms producing both cotton and peanuts. The average for all Louisiana areas growing the two crops was 48 pounds per acre (table 18).

The computed oil outturn per acre from peanuts ranged from 58 pounds in Caddo Parish to 177 pounds in Allen Parish, averaging 92 pounds for all areas growing the two crops.

On a relative basis, peanuts produced nearly twice as much oil per acre, on the average, as cottonseed, ranging as low as 1.4 times in Webster and Washington Parishes to 2.8 times in Allen Parish.

Twenty-two percent of the farms studied produced more oil per acre from cottonseed than from peanuts. In Washington Parish, 40 percent of the farms excelled with cottonseed; in Allen Parish not any of the 28 farms did better with cottonseed; but a substantial percentage of the farms in other parishes produced more oil per acre from cottonseed than from peanuts.

Thirty-five percent of the farms produced less than 40 pounds of oil per acre from cottonseed; 60 percent less than 60 pounds; 82 percent less than 80 pounds; and 93 percent less than 100 pounds (table 19).

From peanuts, 17 percent produced less than 40 pounds of oil per acre; 27 percent less than 60 pounds; 48 percent less than 80 pounds; 61 percent less than 100 pounds; and 39 percent 100 pounds or more (table 19).

MISSISSIPPI

From 5 major areas of Mississippi, 2,750 farms were selected in 13 counties for a comparison of oil and meal yields from cottonseed and peanuts. Mississippi ranks first among the 9 States on cotton lint yields but eighth on peanut yields for those areas of the State growing the two crops. By counties, Lowndes County was low with a cotton lint yield of 203 pounds per acre; Simpson County was high with 357 pounds; and the average for all areas growing both crops was 279 pounds (table 20).

Peanut yields averaged 347 pounds per acre, ranging from 233 pounds in Holmes County to 450 pounds in Itawamba County.

The computed oil outturn from cottonseed averaged 81 pounds per acre for all the farms included in the sample. All areas except the Black Belt averaged very close to this quantity. Only 59 farms were included in the Black Belt area, and the computed cottonseed oil outturn per acre from these amounted to only 58 pounds.

The outturn of oil from peanuts was only 104 pounds per acre for all farms included, ranging from 70 pounds in Holmes County to 135 pounds in Itawamba County.

On a farm to farm basis in Mississippi, peanuts yield only a very little more oil per acre than cottonseed. The ratio of peanut oil yield per acre to cottonseed oil yield per acre for the 2,750 farms was 128 percent. In two of the 13 counties cotton excelled peanuts. In only 2 of the 13 counties did peanuts yield as much as 1-1/2 times the oil per acre as came from cottonseed. Forty-five percent of the farms included in the study produced more oil per acre from cottonseed than from peanuts; the lowest county average was 25 percent of the farms in favor of cottonseed.

From the frequency distribution (table 21) of farms by oil yield per acre is revealed the fact that approximately half of the farms produced between 60 and 99 pounds of oil per acre from cottonseed. Peanut oil yields are distributed all the way from below 20 pounds to more than 400 pounds, with more than half of them below 100 pounds per acre.

Only 8 percent of the farms produced less than 40 pounds of oil per acre from cottonseed, compared with 21 percent for peanuts; 24 percent produced less than 60 pounds from cottonseed, compared with 31 percent for peanuts; 47 percent produced less than 80 pounds from cottonseed, compared with 48 percent for peanuts; 71 percent produced less than 100 pounds from cottonseed, compared with 58 percent for peanuts; and 29 percent produced 100 pounds or more from cottonseed, compared with 42 percent for peanuts (table 21).

OKLAHOMA

From 20 counties, representing 6 major areas of Oklahoma, 3,762 farms were studied to compare per acre oil outturn from cottonseed and peanuts, farm by farm. These farms had a weighted average cotton lint yield of 147 pounds and a peanut yield of 528 pounds per acre. By counties the cotton lint yield ranged from 97 pounds to 235 pounds. The Rolling Plains area averaged 192 pounds of lint per acre, compared with 133 pounds in the Coastal Plains. Peanuts averaged 640 pounds in the Coastal Plains, and almost as high with 622 pounds in the Rolling Plains. The lowest county average for peanuts among the 20 counties was 327 pounds in Latimer County; the highest was 807 pounds in Caddo County (table 22).

The computed oil outturn from cottonseed ranged from 25 pounds per acre to 57 pounds, by counties; by areas, from 34 to 47 pounds. The weighted average for all farms included in the study was 37 pounds per acre.

The computed oil outturn from peanuts was only 98 pounds per acre in Latimer County but went up to 242 pounds in Caddo County, averaging 158 pounds for all farms in the study. The Coastal Plains has the highest area average (192 pounds) followed in order by Rolling Plains (187 pounds), Cross Timbers (166 pounds), Central Prairies (153 pounds), Eastern Prairies (143 pounds), and the Ozark-Cuachita Highlands (106 pounds).

On a comparative basis, peanuts produce 4-1/4 times as much oil per acre as cottonseed, ranging by counties from 2.6 times to 5.6 times as much. The Coastal Plains area produces more than 5-1/2 times as much oil per acre from peanuts as from cottonseed; the Cross Timbers, 4-1/2 times; the Rolling Plains, the Central Prairies, and the Eastern Prairies, nearly 4 times; and the Ozark-Cuachita Highlands, about 2-3/4 times as much from peanuts as from cottonseed.

Although peanuts produce considerably more oil per acre, on the average, cottonseed excelled on 7 percent of the farms studied, ranging from 3 to 10 percent by counties (table 22).

In all areas except the Rolling Plains, about 60 percent of the farms produced less than 40 pounds of oil per acre from cottonseed. By areas, up to 12 percent of the farms produced less than 40 pounds of oil per acre from peanuts. Peanut oil yields range all the way from near-failure to over 400 pounds per acre, with only slight tendency to group around any particular yield.

Twenty percent of the 3,762 Oklahoma farms included in the study produced less than 20 pounds of oil per acre from cottonseed, compared with 2 percent for peanuts; 59 percent produced less than 40 pounds from cottonseed, compared with 8 percent for peanuts; 84 percent produced less than 60 pounds per acre from cottonseed, compared with 15 percent for peanuts;

94 percent produced less than 80 pounds from cottonseed, compared with 24 percent for peanuts; 98 percent produced less than 100 pounds from cottonseed, compared with 33 percent for peanuts; and only 2 percent produced 100 pounds or more from cottonseed, compared with 67 percent for peanuts. Thirty-seven percent of the farms produced from 100 to 199 pounds of oil per acre from peanuts; 20 percent from 200 to 299 pounds; 7 percent from 300 to 399 pounds; and 3 percent 400 pounds or more (table 23).

SOUTH CAROLINA

In South Carolina, 1,742 farms from 13 representative counties from 4 major areas were included in the study.

The weighted average cotton lint yield for the South Carolina areas growing both cotton and peanuts was 215 pounds per acre; peanuts averaged 437 pounds (table 24).

The highest county average cotton lint yield was 448 pounds in Marion County; the lowest, 158 pounds in Barnwell County. The weighted average by areas gives the Piedmont Plateau 310 pounds; the Fall Line Sand Hills 219 pounds; the Lower Coastal Plain 218 pounds; and the Middle Coastal Plain 202 pounds. With these yields the computed oil outturn from cottonseed averages 60 pounds per acre for all areas. By counties, it ranges from 43 pounds per acre in Barnwell County to 122 pounds in Marion County. By areas, the Piedmont Plateau leads with 90 pounds of oil per acre, followed in order by the Fall Line Sand Hills with 63 pounds, the Lower Coastal Plain with 59 pounds, and the Middle Coastal Plain with 55 pounds.

Peanut yields averaged 910 pounds for the 69 farms included from Horry County but only 300 pounds for the 267 farms in Allendale County. By areas, the Fall Line Sand Hills was highest, with an average of 512 pounds, followed by the Piedmont Plateau where a few farms from Anderson County brought the average yield for the area up considerably. The Middle Coastal Plain had an average peanut yield of 412 pounds and the Lower Coastal Plain 367 pounds. These yields resulted in a computed oil outturn of 130 pounds per acre for all areas included in the sample. By counties, the computed oil outturn per acre ranged from 90 pounds in Allendale and Dorchester to 273 pounds in Horry. The Fall Line Sand Hills averaged 152 pounds of oil per acre; the Piedmont Plateau and the Middle Coastal Plain 128 pounds each; and the Lower Coastal Plain 106 pounds.

The ratio of peanut oil yield per acre to cottonseed oil yield per acre shows Horry County producing 3 times as much from peanuts, but Edgefield County only 1.2 times. For all areas peanuts produce 2.2 times as much oil per acre as cottonseed. The Fall Line Sand Hills show an advantage for peanuts of 2.4 times; the Middle Coastal Plain 2.3 times; the Lower Coastal Plain 1.8 times; and the Piedmont Plateau only 1.4 times (table 24).

Twenty-one percent of the farms produced more oil per acre from peanuts than from cottonseed. In Dorchester County 58 percent did better with cottonseed, but only 31 farms were included in the sample. In the Middle Coastal Plain, with more than a thousand farms in the sample, 23 percent produced more oil per acre from cottonseed than from peanuts.

The frequency distribution (table 25) shows that 27 percent of the farms produced less than 40 pounds of oil per acre from cottonseed, 51 percent less than 60 pounds, 68 percent less than 80 pounds, 82 percent less than 100 pounds, and 18 percent 100 pounds or more.

For peanuts, 11 percent of the farms produced less than 40 pounds of oil per acre, 22 percent less than 60 pounds, 35 percent less than 80 pounds, 45 percent less than 100 pounds, 34 percent from 100 to 199 pounds, 14 percent from 200 to 299 pounds, 5 percent from 300 to 399 pounds, and 2 percent 400 pounds or more.

TEXAS

In Texas, 4,196 farms from 29 representative counties from 7 major areas growing both cotton and peanuts were included in the study.

The weighted average cotton yield for these farms was only 105 pounds compared with the 1942 yield for the State of 182 pounds. Peanuts averaged 469 pounds per acre, which is only 11 pounds less than the 1942 State average of 480 pounds (table 26).

Cotton lint yields were as low as 63 pounds in the Rio Grande Plain and as high as 222 pounds in the High Plain. The average for the Rolling Plains was 161 pounds; Coastal Plain 125 pounds; Edwards Plateau 103 pounds; Grand Prairie 102 pounds; and Cross Timbers 92 pounds. The computed oil outturn from these yields indicates that the High Plains would yield about 50 pounds of oil per acre from cottonseed, 36 pounds in the Rolling Plains, 29 pounds in the Coastal Plain, 23 pounds in the Edwards Plateau, 21 pounds in the Cross Timbers, and 24 pounds as the average for all areas growing cotton and peanuts.

Peanut yields averaged as low as 157 pounds per acre for 43 farms in Starr County and as high as 860 pounds for 53 farms in Lamb County. By areas, the High Plains ranked first with an average of 602 pounds, followed in order by the Cross Timbers with 510 pounds, Edwards Plateau with 503 pounds; Rio Grande Plain with 479 pounds; Rolling Plains with 454 pounds; Coastal Plain with 423 pounds; and Grand Prairie with 350 pounds. The computed oil outturn from these yields would be as low as 47 pounds per acre in Starr County to as high as 258 pounds in Lamb County. The High Plains would turn out 181 pounds of oil per acre from peanuts, Cross Timbers 153 pounds, Edwards Plateau 151 pounds, Rio Grande Plain 144 pounds,

Rolling Plains 136 pounds, Coastal Plain 127 pounds, Grand Prairie 105 pounds, and the average for all areas growing cotton and peanuts 141 pounds per acre.

The ratio of peanut oil yield per acre to cottonseed oil yield per acre shows that peanuts have a considerable advantage in all counties, ranging from a low of 2.2 times in Wilbarger County to 11.3 times in Atascosa County. In the Rio Grande Plain peanuts produce 10.3 times as much oil per acre as cottonseed, based on a sample of 309 farms; the Cross Timbers produce 7.3 times, based on a sample of 515 farms; the Edwards Plateau 6.6 times, based on 295 sample farms; the Coastal Plain 4.4 times, based on 1,875 sample farms; the Grand Prairie 4.2 times, based on 84 farms; the High Plains 3.6 times, based on 289 farms; and all areas growing cotton and peanuts show peanuts producing 5.9 times as much oil per acre as cottonseed, based on a weighted average for the 4,196 farms included in the study.

The frequency distribution (table 27) shows the range of oil yields per acre by 20-pound class intervals up to 400 pounds of oil per acre. In no area does any of the farms exceed 159 pounds of oil per acre from cottonseed, but in 6 of the 7 areas a small percentage of the farms produced 400 pounds of oil or more per acre from peanuts. For the State as a whole, 35 percent of the farms produced less than 20 pounds of oil per acre from cottonseed, 78 percent less than 40 pounds, 93 percent less than 60 pounds, 97 percent less than 80 pounds, and 99 percent less than 100 pounds.

With peanuts only 4 percent produced less than 20 pounds of oil per acre, 14 percent less than 40 pounds, 24 percent less than 60 pounds, 34 percent less than 80 pounds, 45 percent less than 100 pounds, 36 percent from 100 to 199 pounds, 13 percent from 200 to 299 pounds, 5 percent from 300 to 399 pounds, and 1 percent 400 pounds or more.

As in Alabama to represent the Southeast, a special analysis was made in Texas to represent the Southwest. Tables 28 and 29 give the results of the special study designed to show, by areas, the percentage of farms producing one, two, three, four, five, and six times or more as much oil per acre from peanuts as from cottonseed. For the entire sample, 36 percent of the farms produced more than 6 times as much oil per acre from peanuts as from cottonseed. Fifty-three percent of the farms produced more than 4 times, 79 percent more than twice as much, and 93 percent as much or more from peanuts as from cottonseed. Similar results are shown by areas, with the Rio Grande Plain, Cross Timbers, and Edwards Plateau having higher percentages of the farms producing four, five, and six times as much oil per acre from peanuts as from cottonseed.

II. PEANUTS VERSUS COTTONSEED FOR MEAL PRODUCTION

Computations for the meal outturn from cottonseed and peanuts are presented in tables similar to the presentation of the data on the oil outturn. The data are given for 23,713 farms, by physical resource areas and by States and counties in tables 30 to 50. One series of tables gives the 1942 cotton lint and peanut yields, computed meal outturn from cottonseed and peanuts, ratio of peanut meal yield per acre to cottonseed meal yield per acre, and the percent of farms producing more meal per acre from cottonseed or from peanuts. The second series of tables gives a frequency distribution of the 23,713 farms by the meal yields per acre from cottonseed and peanuts, in 50-pound class intervals, by States and physical resource areas.

For purposes of this report, only a few of the major highlights will be cited here. For State and county details see tables 30 to 50, inclusive.

Obviously, the high and low producing areas for meal will be the same as reported above for oil, but the advantage of peanuts over cottonseed in the production of meal is less pronounced than in the production of oil. For the Southern Region as a whole 17 percent of the farms produced more oil per acre from cottonseed, but 36 percent of the farms produced more meal per acre from cottonseed. Peanuts yielded 3.5 times as much oil per acre as cottonseed but only 1.9 times as much meal. Each of the 23 areas averaged more oil per acre from peanuts than from cottonseed, but in 6 of the 23 areas cottonseed excelled peanuts in the per acre production of meal (table 30). The average for all areas included in the study for Mississippi shows that State producing only 71 percent as much meal per acre from peanuts as from cottonseed. The per acre yields, by areas, show a computed meal outturn of only 44 pounds from cottonseed in the Rio Grande Plain. On the other extreme, Sand Mountain farmers produced 341 pounds of meal per acre from cottonseed. The average for the Southern Region was 141 pounds. By States, the meal yields varied from 74 pounds in Texas to 211 pounds in Mississippi.

The computed meal outturn from peanuts averaged 269 pounds per acre for the Southern Region, ranging from 101 pounds in the Black Belt to 364 pounds in the Sand Mountain area, or from 149 pounds in Mississippi to 274 pounds in Georgia.

Relatively, the Rio Grande Plain of Texas shows up best for peanuts, by producing 5.4 times as much meal per acre from peanuts as from cottonseed; the Brown Loams of Mississippi is relatively best for cotton, as peanuts in that area produced only 0.6 as much meal per acre as cottonseed on farms growing both crops. Large areas of Oklahoma and Texas and the Middle Coastal Plain of the Southeast produced more than twice as much meal per acre from peanuts as from cottonseed.

A certain percentage of the farms in all areas produced more meal per acre from cottonseed than from peanuts, ranging from 7 percent in the Edwards Plateau to 76 percent in the Brown Loams area, and averaging 36 percent for the Southern Region. In 6 of the 23 areas, approximately two-thirds of the farms excelled with cottonseed; in 2 areas approximately one-half did likewise; in 3 areas about one-third; in 5 areas about one-fourth; in 4 areas about one-fifth; and in 3 areas a still smaller percentage did better with cottonseed than peanuts in meal production per acre. By States, 76 percent of the farms in Mississippi produced more meal per acre from cottonseed than peanuts; in Florida, only 15 percent; Texas, 19 percent; Oklahoma, 20 percent; Georgia, 25 percent; Arkansas, 38 percent; Louisiana, 41 percent; Alabama, 47 percent; and South Carolina, 51 percent.

III. SOYBEANS VERSUS COTTON FOR OIL PRODUCTION

Only 3 of the 9 Southern Region States are important in the production of soybeans. From these 3, Arkansas, Louisiana, and Mississippi, plus Texas, 28 representative counties were selected, from which per acre yield data were tabulated for 4,057 farms growing cotton and soybeans. The majority of these farms were in the Mississippi River Delta areas of Arkansas, Louisiana, and Mississippi.

Table 51 gives a State and area summary of the information as it relates to oil production. Weighted average cotton yields for the areas growing soybeans and cotton were: Arkansas, 518 pounds; Louisiana, 386 pounds; Mississippi, 447 pounds; and Texas, 291 pounds. The Mississippi River Delta area averaged 493 pounds; and the Red River Delta area 326 pounds.

Soybeans averaged 17 bushels per acre in Arkansas; 11.5 bushels in Louisiana; 15.8 bushels in Mississippi; and 8.7 bushels in the Texas areas growing soybeans and cotton.

The high cotton yields and high oil outturn from the Arkansas Delta cottonseed puts cotton far out ahead of soybeans in the per acre production of oil. The computed oil outturn from cottonseed is 167 pounds per acre, compared with 130 pounds for soybeans. The State figure includes a few farms from the Red River Delta where soybeans gave better results per acre than cottonseed. Both Louisiana and Mississippi produced more oil per acre from cottonseed than from soybeans, while soybeans did better than cottonseed on the few farms included in the study from Texas. The weighted average for all areas included in the study shows that soybeans did only 90 percent as well as cottonseed in per acre oil production, ranging from 78 percent in Arkansas to 115 percent in Texas. In Arkansas 76 percent of the farms produced more oil per acre from cottonseed than from soybeans; Louisiana, 64 percent; Mississippi, 64 percent; and Texas, 47 percent.

The Mississippi River Delta, the principal soybean area of the Southern Region, produced 161 pounds of oil per acre from cottonseed and only 81 percent as much or 130 pounds per acre from soybeans. In the Mississippi River Delta areas of Arkansas and Louisiana, soybeans produced about three-fourths as much oil per acre as cottonseed; in the Delta areas of Mississippi, 88 percent as much. For all the Delta areas 73 percent of the farms produced more oil per acre from cottonseed than from soybeans.

A special analysis of the data for Arkansas and Mississippi (tables 52 and 53) shows that 18 percent of the farms produced less than half as much oil per acre from cottonseed as from soybeans and 71 percent produced less oil per acre from soybeans than from cottonseed. Three percent of the farms produced twice as much oil per acre from cottonseed as from soybeans. These contrasts are more striking in the Mississippi River Delta areas than in other parts of these two States.

Tables 54 to 59 give, by counties, frequency distribution of farms by oil yield per acre from cottonseed and soybeans, 1942 yield per acre for cotton and soybeans, computed oil outturn from cottonseed and soybeans, ratio of soybean oil yield per acre to cottonseed oil yield per acre, percent of farms producing more oil per acre from cottonseed or soybeans, and number of farms in the sample.

IV. SOYBEANS VERSUS COTTONSEED FOR MEAL PRODUCTION

In the areas growing cotton and soybeans, the per acre meal outturn from soybeans is approximately 1-3/4 times that from cottonseed (table 60). The computed meal outturn from cottonseed in the Mississippi River Delta is 448 pounds per acre, compared with 774 pounds from soybeans. In the Red River Delta and in Texas, soybeans produced more than twice as much meal per acre as cottonseed. Nevertheless, approximately one-fifth of the farms produced more meal per acre from cottonseed than from soybeans.

Further details, by counties and areas, are given in tables 60 to 66.

Table 1.- Yield of oil and meal per 100 pounds of seed from peanuts, soybeans, and cottonseed

| State | Peanuts 1/ | | | Soybeans 2/ | | | Cottonseed 3/ | | |
|----------------|------------|------|--------|-------------|------|--------|---------------|------|--------|
| | Oil | Meal | Pounds | Oil | Meal | Pounds | Oil | Meal | Pounds |
| | | | Pounds | | | Pounds | | | Pounds |
| Alabama | 30 | 43 | | | | | 16.1 | | 43.5 |
| Arkansas | 30 | 50 | | 12.7 | | 78.9 | | 15.4 | 44.3 |
| Georgia | 30 | 43 | | | | | 16.2 | | 43.6 |
| Florida | 30 | 43 | | | | | 16.2 | | 43.6 |
| Louisiana | 30 | 50 | | 13.7 | | 79.9 | | 15.8 | 43.5 |
| Mississippi | 30 | 43 | | | 14.0 | | 79.6 | | 16.8 |
| Oklahoma | 30 | 50 | | | | | 14.0 | | 43.7 |
| South Carolina | 30 | 43 | | | | | 13.5 | | 46.6 |
| Texas | 30 | 50 | | 14.4 | | 83.1 | | 14.4 | 44.7 |

1/ Estimates from table 5, The Fats and Oils Situation, March 1943. The published estimates on meal for Arkansas, Louisiana, Oklahoma, and Texas are 54 pounds per 100 pounds of seed, rather than the 50 pounds estimated herein, which is a compromise between the published estimate and unpublished data for the 1942 crop for certain mills in those States.

2/ Based on information furnished the Southern Division, AAA, by mills crushing the 1942 crop of soybeans produced in these States.

3/ Based on turnout as reported by the Bureau of the Census for August 1942 to July 1943.

Table 2.—Factors applied to peanut, soybean, and cotton lint yields to convert such yields to oil and meal yields per acre.

22

Table 3.- Comparative data on oil yields per acre from cottonseed and peanuts,
by areas, Southern Region, 1942

| Number and name of area 1/ | Number of farms in sample: | Yield per acre, 1942 | | Ratio of oil yield: producing more oil per acre, to peanuts: | | Percent of farms | |
|--|----------------------------|----------------------|------------------------------------|--|----------|------------------|----------------------|
| | | Cotton: | Computed oil cutturn from Peanuts: | Cotton: | Peanuts: | Cotton seed: | Peanuts: cottonseed: |
| | | in lint: | to seed: | in cotton: | to seed: | in cotton: | to seed: |
| 1. High Plains | 289 | 222 | 602 | 50 | 181 | 362 | 9 |
| 3. Rio Grande Plain | 309 | 63 | 479 | 14 | 144 | 1,029 | 5 |
| 4. Edwards Plateau | 295 | 103 | 503 | 23 | 151 | 657 | 2 |
| 5. Rolling Plains | 1,389 | 184 | 579 | 43 | 174 | 405 | 7 |
| 6. Oklahoma Central Prairies | 509 | 162 | 511 | 39 | 154 | 395 | 10 |
| 8. East Oklahoma Prairies | 376 | 148 | 477 | 38 | 143 | 376 | 9 |
| 9. Cross Timbers | 2,114 | 133 | 547 | 33 | 164 | 497 | 5 |
| 10. Grand Prairie | 84 | 102 | 350 | 25 | 105 | 420 | 8 |
| 13. Coastal Plain (South Central) | 3,623 | 144 | 537 | 37 | 161 | 435 | 14 |
| 14. Ozark-Ouachita Highlands | 1,117 | 160 | 370 | 45 | 111 | 247 | 12 |
| 17. Brown Loams | 1,211 | 277 | 315 | 80 | 94 | 118 | 48 |
| 18. Sand-Clay Hills | 704 | 276 | 341 | 80 | 102 | 128 | 39 |
| 19. Black Belt | 555 | 187 | 236 | 53 | 71 | 134 | 43 |
| 20. Upper Coastal Plain | 979 | 267 | 383 | 72 | 115 | 160 | 36 |
| 21. Limestone Basin | 300 | 311 | 710 | 80 | 213 | 266 | 2 |
| 22. Sand Mountain | 597 | 490 | 846 | 126 | 254 | 202 | 16 |
| 23. Appalachian Highlands | 133 | 250 | 305 | 64 | 92 | 144 | 37 |
| 25. Piedmont Plateau | 1,607 | 251 | 355 | 67 | 106 | 158 | 32 |
| 26. Fall Line Sand Hills | 711 | 181 | 520 | 52 | 155 | 298 | 10 |
| 27. Coastal Plain - Red Belt | 299 | 229 | 577 | 63 | 173 | 275 | 2 |
| 28. Middle Coastal Plain | 6,259 | 192 | 662 | 52 | 197 | 379 | 12 |
| 29. Lower Coastal Plain | 114 | 223 | 446 | 60 | 131 | 218 | 30 |
| 31. Rolling Sandy Lands and Flatwoods | 133 | 153 | 431 | 42 | 125 | 298 | 13 |
| Southern Region | 23,707 | 188 | 584 | 50 | 175 | 350 | 17 |

1/ Numbers correspond with area numbers on map in this report.

Southern Division, AAA
November 12, 1943

Table 4.- Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas, Southern Region, 1942

| Oil yield per acre (pounds) | Percent | | | | | | | | | | |
|-----------------------------------|-----------------------|----------------------------|--------------------|---------------------|--------------------|----------------------|------------------|-------------------|---------------------|----------------|----------------|
| | Rio High Plains | Edwards Grande Plain | Rolling Plateau | Oklahoma: Plains | Central Prairie | Oklahoma: Timbers | Grand Prairie | Cross Prairies | Ozark: Highlands | Brown Loams | Sand: Hills |
| 0-19 | 15 | 2 | 82 | 6 | 41 | 0 | 22 | 2 | 13 | 3 | 26 |
| 20-39 | 23 | 4 | 17 | 12 | 48 | 4 | 34 | 6 | 47 | 9 | 35 |
| 40-59 | 27 | 9 | 1 | 14 | 11 | 5 | 20 | 8 | 28 | 9 | 18 |
| 60-79 | 20 | 5 | * | 8 | 4 | 13 | 7 | 8 | 10 | 11 | 9 |
| 80-99 | 13 | 5 | 9 | 16 | 7 | 8 | 2 | 7 | 6 | 10 | 1 |
| 100-119 | 2 | 5 | 9 | 8 | 3 | 8 | 1 | 9 | 2 | 9 | 1 |
| 120-139 | * | 6 | 12 | 8 | 1 | 6 | 1 | 11 | * | 10 | * |
| 140-159 | * | 5 | 7 | 12 | * | 7 | 1 | 7 | * | 9 | 0 |
| 160-179 | 5 | 6 | 13 | * | 7 | 6 | 5 | * | 7 | * | 5 |
| 180-199 | 6 | 5 | 7 | 7 | 7 | 4 | 5 | * | 4 | 4 | * |
| 200-219 | 6 | 5 | 5 | 5 | 5 | 4 | 4 | 5 | 0 | 6 | * |
| 220-239 | 5 | 1 | 6 | 4 | 4 | 4 | 3 | * | 5 | 0 | * |
| 240-259 | 6 | 1 | 4 | 5 | 4 | 4 | 3 | * | 0 | 3 | * |
| 260-279 | 5 | 2 | 3 | 5 | 4 | 3 | 4 | * | 1 | 2 | * |
| 280-299 | 5 | 1 | 1 | 3 | 2 | 1 | 2 | * | 1 | 0 | * |
| 300-319 | 2 | * | 1 | 2 | 2 | 1 | 1 | 2 | 1 | 2 | * |
| 320-339 | 2 | 1 | 1 | 2 | 2 | 1 | 1 | 1 | 0 | 1 | * |
| 340-359 | 3 | * | 2 | 2 | 1 | 1 | 1 | * | 1 | 0 | * |
| 360-379 | 4 | 0 | * | 1 | 2 | 1 | 1 | * | 0 | 0 | * |
| 380-399 | 3 | * | 1 | * | 1 | 1 | 1 | * | 0 | 0 | * |
| 400 and over | 7 | 1 | * | 3 | 3 | 2 | 2 | * | * | * | * |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 289 | 309 | 295 | 1,389 | 509 | 376 | 2,114 | 84 | 3,623 | 1,117 | 704 |

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 12, 1943

— continued

Table 4.- Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas, Southern Region, 1942, (cont.)

| Oil yield per acre (pounds) | Black Belt | Upper Coastal Plain | 'Limestone' Basin | Sand Mountains | 'Appalachian' Highlands | 'Piedmont' Plateau | 'Fall Line' Sand Hills | Coastal Plain | Middle Red Belt | Lower Plain | Rolling Lands and Flatwoods | Percent | | | | | | | | | | | | |
|-----------------------------------|---------------|---------------------------|----------------------|-------------------|----------------------------|-----------------------|---------------------------|------------------|--------------------|----------------|-----------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--|
| | | | | | | | | | | | | Cot- ton- nuts; seed; | |
| 0- 19 | : | 15 | 17 | 2 | 4 | 0 | 0 | 0 | 2 | 10 | 9 | * | 3 | 0 | 9 | 2 | 4 | 1 | 16 | 2 | | | | |
| 20- 39 | : | 28 | 19 | 11 | 13 | 3 | 0 | * | 22 | 13 | 24 | 15 | 22 | 3 | 40 | 0 | 26 | 5 | 18 | 14 | 37 | 8 | | |
| 40- 59 | : | 21 | 23 | 20 | 12 | 19 | * | 2 | 1 | 24 | 16 | 21 | 15 | 33 | 7 | 42 | 2 | 28 | 6 | 31 | 14 | 30 | 9 | |
| 60- 79 | : | 16 | 13 | 25 | 16 | 30 | 0 | 8 | 3 | 27 | 15 | 17 | 17 | 19 | 11 | 9 | 7 | 19 | 7 | 24 | 11 | 8 | 9 | |
| 80- 99 | : | 10 | 8 | 23 | 12 | 26 | 3 | 12 | 5 | 15 | 10 | 13 | 12 | 14 | 9 | 5 | 10 | 10 | 7 | 10 | 8 | 4 | 12 | |
| 100-119 | : | 6 | 5 | 12 | 9 | 18 | 14 | 22 | 6 | 5 | 4 | 9 | 9 | 4 | 11 | 0 | 13 | 5 | 8 | 8 | 9 | 4 | 11 | |
| 120-139 | : | 2 | 5 | 5 | 9 | 4 | 9 | 23 | 7 | 2 | 7 | 4 | 6 | 5 | 13 | 1 | 16 | 2 | 9 | 3 | 4 | 1 | 16 | |
| 140-159 | : | 1 | 3 | 1 | 6 | 10 | 17 | 7 | 2 | 10 | 2 | 6 | 1 | 11 | 1 | 16 | 1 | 8 | 2 | 6 | 2 | | | |
| 160-179 | : | 1 | 2 | 1 | 4 | 7 | 9 | 5 | 0 | 3 | 1 | 4 | 1 | 7 | 10 | * | 7 | 7 | 7 | 7 | 4 | 3 | 4 | |
| 180-199 | : | * | 2 | * | 4 | 5 | 4 | 8 | 0 | 3 | 0 | 2 | 0 | 6 | 7 | * | 7 | 7 | 7 | 7 | 3 | 4 | | |
| 200-219 | : | 1 | * | 2 | 7 | 3 | 8 | 0 | 4 | * | 2 | * | 6 | * | 6 | 7 | * | 7 | * | 7 | 5 | 4 | | |
| 220-239 | : | 1 | 2 | 2 | 9 | 5 | 1 | 2 | * | 2 | 0 | 4 | 1 | * | 3 | 2 | 1 | 4 | 1 | 6 | 4 | 1 | | |
| 240-259 | : | * | 2 | 1 | 6 | 6 | 6 | 2 | * | 1 | * | 3 | 1 | * | 1 | 1 | 1 | 1 | 4 | 3 | 5 | 2 | | |
| 260-279 | : | 1 | 1 | 6 | 6 | 5 | 0 | 1 | 1 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 4 | 3 | 4 | | |
| 280-299 | : | * | 1 | 1 | 6 | 3 | 1 | 1 | 0 | 2 | 1 | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 3 | 3 | 3 | 0 | | |
| 300-319 | : | 2 | 5 | 6 | 5 | 3 | 5 | 6 | * | 0 | 2 | * | 1 | * | 1 | 1 | 1 | 1 | 2 | 3 | 2 | 1 | | |
| 320-339 | : | 1 | 1 | 3 | 4 | 3 | 5 | 5 | * | * | 1 | * | 1 | * | 1 | 1 | 1 | 1 | 1 | 2 | 0 | 1 | | |
| 340-359 | : | 0 | 1 | 3 | 5 | 3 | 5 | 5 | * | * | 1 | * | 1 | * | 1 | 0 | 1 | 1 | 1 | 1 | 3 | 0 | | |
| 360-379 | : | * | 1 | 3 | 2 | 2 | 2 | 2 | * | 1 | * | 1 | * | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | | |
| 380-399 | : | * | 2 | 1 | 12 | * | 1 | 12 | * | * | 1 | * | 1 | * | 1 | 2 | * | 2 | 1 | 1 | 1 | 1 | | |
| 400 and over | : | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * | | |
| Total | : | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | |
| Number of farms in sample | : | 555 | 979 | 300 | 597 | 133 | 1,607 | 711 | 182 | 6,259 | 114 | 133 | | | | | | | | | | | | |

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 12, 1945

Table 5.- Comparative data on oil yields per acre from cottonseed and peanuts,
by States, Southern Region, 1942

| State | Number of farms in sample | Yield per acre, 1942 | | | Ratio of : Percent of farms oil yield:producing more oil per acre,: per acre from | | |
|-----------------|---------------------------------------|----------------------|---------|----------------------------------|---|-----------------------|-------------------------|
| | | Cotton | Peanuts | outturn from lint ; sample | peanuts | Cotton- to seed | Peanuts ;cottonseed: |
| | Number | Pounds | Pounds | Pounds | Percent | Percent | Percent |
| Alabama | 3,875 | 179 | 593 | 48 | 174 | 362 | 22 |
| Arkansas | 1,150 | 188 | 396 | 56 | 119 | 212 | 17 |
| Florida | 876 | 151 | 535 | 42 | 159 | 379 | 4 |
| Georgia | 4,054 | 203 | 637 | 55 | 191 | 347 | 9 |
| Louisiana | 1,302 | 171 | 306 | 48 | 92 | 192 | 22 |
| Mississippi | 2,750 | 279 | 347 | 81 | 104 | 128 | 45 |
| Oklahoma | 3,762 | 147 | 528 | 37 | 158 | 427 | 7 |
| South Carolina | 1,742 | 215 | 437 | 60 | 130 | 217 | 21 |
| Texas | 4,196 | 105 | 469 | 24 | 141 | 588 | 8 |
| Southern Region | 23,707 | 188 | 584 | 50 | 175 | 350 | 17 |

Table 6.- Frequency distribution of farms by oil yields per acre from cottonseed and peanuts,
by States, Southern Region, 1942

| Oil yield per acre (pounds) | Percent | | | | | | | | | | Southern Region |
|-----------------------------------|---------|----------|---------|---------|-----------|-------------|----------|----------------|-------|--------|--------------------|
| | Alabama | Arkansas | Georgia | Florida | Louisiana | Mississippi | Oklahoma | South Carolina | Texas | South | |
| 0- 19 | 9 | 6 | 8 | 2 | 5 | 1 | 18 | 10 | 7 | 7 | 2 |
| 20- 39 | 22 | 10 | 24 | 9 | 26 | 4 | 35 | 4 | 25 | 14 | 39 |
| 40- 59 | 19 | 10 | 31 | 11 | 31 | 5 | 25 | 10 | 16 | 10 | 25 |
| 60- 79 | 17 | 9 | 22 | 12 | 21 | 6 | 14 | 6 | 22 | 17 | 10 |
| 80- 99 | 12 | 8 | 10 | 11 | 10 | 7 | 5 | 7 | 11 | 13 | 24 |
| 100-119 | 9 | 8 | 4 | 10 | 4 | 2 | 9 | 5 | 9 | 7 | 10 |
| 120-139 | 5 | 7 | 1 | 12 | 2 | 9 | 1 | 10 | 1 | 6 | 8 |
| 140-159 | 3 | 7 | 4 | 10 | 1 | 9 | * | 10 | 1 | 6 | 4 |
| 160-179 | 2 | 2 | 5 | * | 5 | * | 8 | 0 | 7 | * | 8 |
| 180-199 | 1 | 1 | 5 | 0 | 5 | 0 | 7 | * | 5 | * | 6 |
| 200-219 | 1 | 4 | * | 3 | * | 8 | * | 5 | * | 3 | 4 |
| 220-239 | 1 | 4 | 2 | 2 | 6 | 0 | 5 | * | 2 | 0 | 5 |
| 240-259 | 1 | 4 | 5 | 5 | 6 | * | 6 | 2 | * | 4 | 0 |
| 260-279 | 1 | 5 | 2 | 4 | 5 | 4 | 5 | 1 | * | 4 | 0 |
| 280-299 | 1 | 2 | * | 2 | 5 | 4 | 1 | 0 | * | 2 | 0 |
| 300-319 | 2 | 3 | 3 | 3 | 3 | 3 | 3 | 1 | 0 | 2 | 0 |
| 320-339 | 2 | * | 2 | 3 | 2 | * | 0 | * | 0 | * | 1 |
| 340-359 | 1 | * | 1 | 1 | 2 | * | 0 | * | 0 | 2 | * |
| 360-379 | 1 | * | 1 | 1 | 1 | * | 0 | * | 1 | 1 | 1 |
| 380-399 | 1 | 0 | 1 | 1 | 1 | * | 0 | * | 1 | 1 | 1 |
| 400 and over | 1 | * | 1 | 1 | 1 | * | * | * | 3 | 2 | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms: In sample | 3,875 | 1,150 | 4,054 | 876 | 1,302 | 2,750 | 3,762 | 1,742 | 4,196 | 23,707 | |

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 10, 1943

Table 7.- Rank of Southern Region States by various factors related to oil yields per acre from cottonseed and peanuts, 1942

| | | | Percent of oil yield per acre from cottonseed | Percent of oil yield per acre from peanuts | farms producing more oil per acre from cottonseed | farms producing more oil per acre from peanuts | Man-labor requirements per acre ² (High to low) |
|-------|--|---|---|---|---|--|--|
| Rank: | Computed oil outturn per acre from cottonseed | Computed oil outturn per acre from peanuts | Ratio of oil yield per acre; peanuts to cottonseed | Ratio of oil yield per acre; peanuts to peanuts ¹ | Ratio of oil yield per acre; cottonseed to cottonseed ¹ | Ratio of oil yield per acre; peanuts than from cotton- seed ¹ | Ratio of oil yield per acre; peanuts to cotton ¹ |
| 1 | Mississippi | Georgia | : Texas | : Mississippi | : Florida | : South Carolina; Mississippi | : Oklahoma |
| 2 | South Carolina | Alabama | : Oklahoma | : Alabama | : Oklahoma | : Alabama | : Arkansas |
| 3 | Arkansas | Florida | : Louisiana | : Florida | : Texas | : Georgia | : Florida |
| 4 | Georgia | Oklahoma | : Alabama | : South Carolina; Georgia | : South Carolina; Georgia | : Mississippi | : Alabama |
| 5 | Alabama | Texas | : Georgia | : Arkansas | : Arkansas | : Florida | : South Carolina; Louisiana |
| 6 | Louisiana | South Carolina | : South Carolina; Georgia | : Georgia | : South Carolina; Arkansas | : Georgia | : Mississippi |
| 7 | Florida | Arkansas | : Texas | : Texas | : Louisiana | : Louisiana | : Alabama |
| 8 | Oklahoma | Mississippi | : Louisiana | : Oklahoma | : Alabama | : Oklahoma | : Georgia |
| 9 | Texas | Louisiana | : Mississippi | : Florida | : Mississippi | : Texas | : South Carolina |

¹/ Based on 23,707 sample farms from 136 counties in which cotton and peanuts compete.

²/ Based on Labor Requirements for Crops and Livestock, Bureau of Agricultural Economics, May 1943; Louisiana Bulletin 361, February 1943; Mississippi Bulletin 376, March 1943; and Peanuts: A War Crop for Arkansas, Arkansas Mimeo., March 1943.

Table 8.- Comparative data on oil yields per acre from cottonseed and peanuts,
Selected Alabama Counties, 1942.

| County and area | Number of farms in sample | Yield per acre, 1942 Cotton; Peanuts; lint; Cotton- seed; | Computed oil outturn from peanuts : to : Peanuts : seed ; cottonseed; | Ratio of : Percent of farms oil yield producing more oil per acre : per acre from | | |
|-----------------------|---------------------------------------|--|---|---|-------------------------|---------|
| | | | | peanuts | Cotton- to : seed | Peanuts |
| | | | | Percent | Percent | Percent |
| Limestone Basin | | | | | | |
| Limestone | 300 | 311 | 710 | 80 | 213 | 266 |
| Upper Coastal Plain | | | | | | |
| Elmore | 752 | 249 | 332 | 64 | 100 | 156 |
| Franklin | 251 | 206 | 303 | 53 | 91 | 172 |
| Lamar | 206 | 334 | 437 | 86 | 131 | 152 |
| Sand Mountain | | | | | | |
| Cullman | 597 | 490 | 846 | 126 | 254 | 202 |
| DeKalb | 299 | 497 | 1,023 | 128 | 307 | 240 |
| Appalachian Highlands | | | | | | |
| Calhoun | 133 | 250 | 305 | 64 | 92 | 144 |
| Shelby | 74 | 268 | 313 | 69 | 94 | 136 |
| Piedmont Plateau | | | | | | |
| Lee | 59 | 217 | 290 | 56 | 87 | 155 |
| Randolph | 621 | 227 | 283 | 58 | 85 | 147 |
| Black Belt | | | | | | |
| Hale | 350 | 125 | 190 | 32 | 57 | 178 |
| Lowndes | 271 | 254 | 307 | 65 | 92 | 142 |
| Sand-Clay Hills | | | | | | |
| Clarke | 496 | 178 | 227 | 50 | 68 | 136 |
| Middle Coastal Plain | | | | | | |
| Coffee | 242 | 149 | 247 | 42 | 74 | 176 |
| Conecuh | 87 | 138 | 240 | 39 | 72 | 185 |
| Henry | 293 | 156 | 821 | 42 | 198 | 471 |
| Totals | 3,875 | 179 | 593 | 48 | 174 | 362 |

Table 9.- Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Alabama, 1942

| Oil yield per acre (pounds) | -Percent- | | | | | | | |
|-----------------------------------|--|---|-----|-----|-----|-----|-----|-------|
| | Limestone : Upper : Sand : Appalachian : Piedmont : Black : Sand-Clay : Middle : Basin of : Coastal : Mountain : Highland : Plateau : Belt : Hills : Coastal : State Alabama 1/ : Plain 2/ : 3/ : 4/ : 5/ : 6/ : 7/ : Plain 8/ : | Cot-: Pea-: Cot-: Pea-: Cot-: Pea-: Cot-: Pea-: Cot-: Pea-: Cotton-: ton-: nuts: ton-: nuts: ton-: nuts: ton-: nuts: seed: nuts: seed: nuts: seed: nuts: seed: | | | | | | |
| 0-19 | 1 | 0 | 0 | 2 | 4 | 0 | 2 | 10 |
| 20-39 | 1 | 3 | 0 | 14 | 15 | * | 22 | 13 |
| 40-59 | 1 | 18 | * | 22 | 14 | 2 | 24 | 16 |
| 60-79 | 1 | 30 | 0 | 25 | 15 | 8 | 3 | 27 |
| 80-99 | 1 | 26 | 3 | 22 | 13 | 12 | 5 | 15 |
| 100-119 | 1 | 18 | 14 | 10 | 8 | 22 | 6 | 10 |
| 120-139 | 1 | 4 | 9 | 4 | 8 | 23 | 7 | 2 |
| 140-159 | 1 | 10 | 1 | 5 | 17 | 7 | 2 | * |
| 160-179 | 1 | 7 | * | 4 | 9 | 5 | 0 | 3 |
| 180-199 | 1 | 5 | * | 3 | 4 | 8 | 0 | 3 |
| 200-219 | 1 | 7 | 2 | 3 | 8 | 0 | 4 | 1 |
| 220-239 | 1 | 9 | 2 | 5 | 5 | 1 | 2 | 1 |
| 240-259 | 1 | 6 | 2 | 6 | 6 | 2 | * | 0 |
| 260-279 | 1 | 6 | 1 | 5 | 5 | 0 | 0 | 1 |
| 280-299 | 1 | 6 | 1 | 3 | 3 | 1 | * | 0 |
| 300-319 | 1 | 5 | 1 | 6 | 0 | 0 | 0 | 3 |
| 320-339 | 1 | 3 | 1 | 5 | * | 0 | 0 | 2 |
| 340-359 | 1 | 4 | * | 3 | 3 | 0 | 0 | 1 |
| 360-379 | 1 | 3 | * | 3 | * | 3 | * | 1 |
| 380-399 | 1 | 2 | * | 2 | * | 2 | * | 1 |
| 400 and over | 1 | 1 | 12 | 1 | 12 | 1 | 2 | 1 |
| Total | 1 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms: in sample | 1 | — | — | — | — | — | — | — |
| | 300 | 752 | 597 | 133 | 621 | 496 | 87 | 889 |
| | | | | | | | | 3,875 |

Sample counties: 1/ Limestone. 2/ Elmore, Franklin, and Lamar. 3/ Cullman and DeKalb. 4/ Calhoun and Shelby.
5/ Lee and Randolph. 6/ Hale and Lowndes. 7/ Clarke. 8/ Coffee, Conecuh, and Henry.

* Less than 5 tenths of 1 percent.
Southern Division, AAA October 22, 1943

Table 10.- Relative advantage of peanuts over cottonseed in per acre oil production,
by areas in Alabama, 1942

| Area | Percent of farms producing more than times : times : times : time : per acre from : cottonseed than: sample as much oil per acre from peanuts as from cottonseed | | | | | | Percent of farms produc- ing more oil : per acre from : cottonseed than: from peanuts : | Number of farms |
|-------------------------|---|----|----|----|----|----|--|-----------------------|
| | 6 | 5 | 4 | 3 | 2 | 1 | | |
| Limestone Basin 1/ | 3 | 5 | 16 | 39 | 69 | 98 | - | 2 |
| Upper Coastal Plain 2/ | 2 | 3 | 8 | 15 | 30 | 65 | 35 | 752 |
| Sand Mountain 3/ | 1 | 2 | 5 | 16 | 45 | 85 | 15 | 597 |
| Appalachian Highland 4/ | 2 | 2 | 5 | 14 | 29 | 63 | 37 | 133 |
| Piedmont Plateau 5/ | 4 | 6 | 10 | 18 | 35 | 71 | 29 | 619 |
| Black Belt 6/ | 7 | 9 | 14 | 20 | 31 | 57 | 43 | 496 |
| Sand-Clay Hills 7/ | 16 | 19 | 23 | 30 | 40 | 69 | 31 | 87 |
| Middle Coastal Plain 8/ | 36 | 46 | 59 | 75 | 88 | 97 | 3 | 889 |
| Total | 11 | 15 | 21 | 31 | 49 | 78 | 22 | 3,873 |

Sample counties: 1/ Limestone
2/ Elmore, Franklin, and Lamar
3/ Cullman and Dekalb
4/ Calhoun and Shelby

5/ Lee and Randolph
6/ Hale and Lowndes
7/ Clarke
8/ Coffee, Conecuh, and Henry

Southern Division, AAA
October 26, 1943

Table 11.- Frequency distribution of farms by ratio of peanut oil yield per acre to cottonseed oil yield per acre, by areas in Alabama, 1942

| Ratio Percent | Limestone : Basin of : Coastal : Mountain : Highland : Plateau : Belt | | | AppalachianPiedmont : Black : Sand-Clay: Middle : Hills : Coastal : Plain | | | State | |
|-------------------------------|---|----------|-----|---|-----|-----|-------|-------|
| | Alabama 1/ | Plain 2/ | 3/ | 4/ | 5/ | 6/ | 7/ | |
| Under 60 | 0 | 15 | 3 | 20 | 14 | 28 | 9 | 11 |
| 60-79 | 1 | 11 | 5 | 8 | 7 | 6 | 6 | 5 |
| 80-99 | 1 | 9 | 7 | 9 | 8 | 9 | 16* | 1 |
| Under 100 | 2 | 35 | 15 | 37 | 29 | 43 | 31 | 22 |
| 100-119 | 5 | 10 | 7 | 10 | 8 | 5 | 7 | 6 |
| 120-139 | 7 | 9 | 7 | 8 | 9 | 6 | 7 | 7 |
| 140-159 | 4 | 6 | 10 | 5 | 6 | 5 | 2 | 6 |
| 160-179 | 6 | 6 | 10 | 5 | 6 | 7 | 2 | 6 |
| 180-199 | 7 | 4 | 6 | 6 | 7 | 3 | 2 | 4 |
| 200-199 | 29 | 35 | 40 | 34 | 36 | 26 | 29 | 29 |
| 200-219 | 9 | 3 | 6 | 4 | 4 | 3 | 4 | 2 |
| 220-239 | 5 | 4 | 6 | 2 | 4 | 3 | 3 | 4 |
| 240-259 | 6 | 3 | 8 | 5 | 3 | 2 | 1 | 3 |
| 260-279 | 5 | 4 | 6 | 0 | 4 | 1 | 2 | 4 |
| 280-299 | 5 | 1 | 3 | 4 | 2 | 2 | 0 | 2 |
| 200-299 | 30 | 17 | 29 | 15 | 17 | 11 | 10 | 13 |
| 300-319 | 4 | 4 | 3 | 3 | 2 | 2 | 1 | 3 |
| 320-339 | 6 | 2 | 2 | 2 | 2 | 1 | 2 | 2 |
| 340-359 | 5 | 1 | 2 | 2 | 1 | 1 | 2 | 5 |
| 360-379 | 4 | 2 | 2 | 2 | 2 | 1 | 0 | 2 |
| 380-399 | 4 | 1 | 1 | 0 | 1 | * | 2 | 3 |
| 300-399 | 23 | 7 | 11 | 9 | 6 | 6 | 7 | 10 |
| 400-419 | 4 | 2 | 1 | 1 | 1 | 1 | 2 | 2 |
| 420-439 | 2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 440-459 | 2 | 1 | * | 1 | 1 | 1 | 0 | 3 |
| 460-479 | 2 | * | 1 | 0 | 1 | 0 | 2 | 1 |
| 480-499 | 1 | 1 | * | 0 | 1 | 0 | 2 | 1 |
| 400-499 | 11 | 5 | 3 | 3 | 4 | 5 | 4 | 6 |
| 500-519 | 1 | * | * | 0 | 1 | 1 | 0 | 1 |
| 520-539 | 0 | * | * | 0 | 0 | 1 | 0 | 2 |
| 540-559 | 1 | * | * | 0 | 0 | 1 | 1 | 2 |
| 560-579 | 0 | 0 | * | 0 | 0 | 1 | 1 | 0 |
| 580-599 | 0 | * | 0 | 0 | 0 | 1 | 1 | 0 |
| 500-599 | 2 | 1 | 1 | 0 | 2 | 2 | 3 | 4 |
| 600 and over | 3 | 2 | 1 | 2 | 4 | 7 | 16 | 36 |
| Number of farms: in sample | 300 | 752 | 597 | 133 | 619 | 496 | 87 | 889 |
| | | | | | | | | 8,873 |

Sample counties: 1/ Limestone. 2/ Elmore, Franklin, and Lamar. 3/ Cullman and DeKalb. 4/ Calhoun and Shelby. 5/ Lee and Randolph. 6/ Hale and Lowndes. 7/ Clarke. 8/ Coffee, Conecuh, and Henry.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
October 26, 1943

Table 12.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected Arkansas counties, 1942

| County and area | Number of farms | Yield per acre, 1942 | | | Ratio of oil yield: producing more oil per acre, per acre from | | |
|-------------------------|-----------------------|----------------------|----------|------------------------------|---|------------|------------|
| | | Cotton: | Peanuts: | Computed oil outturn from | Peanuts: | Cotton: | Peanuts: |
| | | in lint | sample | seed | to seed | to seed | to seed |
| | | Number | Pounds | Pounds | Pounds | Pounds | Percent |
| Ozark-Ouachita Highland | 687 | 184 | 427 | 54 | 128 | 237 | 13 |
| Faulkner | 162 | 226 | 537 | 66 | 161 | 244 | 1 |
| Garland | 29 | 137 | 330 | 40 | 99 | 248 | 17 |
| Izard | 40 | 256 | 390 | 75 | 117 | 156 | 40 |
| Logan | 209 | 150 | 363 | 44 | 109 | 248 | 14 |
| Montgomery | 69 | 130 | 507 | 38 | 152 | 400 | 4 |
| Searcy | 40 | 130 | 433 | 38 | 130 | 342 | 18 |
| Sebastian | 67 | 140 | 437 | 41 | 131 | 320 | 9 |
| Sharp | 35 | 294 | 487 | 86 | 146 | 170 | 46 |
| Stone | 36 | 174 | 413 | 51 | 124 | 243 | 19 |
| Coastal Plains | 463 | 191 | 367 | 58 | 110 | 190 | 77 |
| Columbia | 121 | 208 | 393 | 61 | 118 | 193 | 17 |
| Little River | 52 | 137 | 307 | 40 | 92 | 230 | 19 |
| Miller | 57 | 185 | 337 | 54 | 101 | 187 | 25 |
| Guachita | 89 | 198 | 337 | 58 | 101 | 174 | 28 |
| Union | 144 | 212 | 437 | 62 | 131 | 211 | 27 |
| Total | 1,150 | 188 | 396 | 56 | 119 | 212 | 17 |
| | | | | | | | 83 |

Table 13.- Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas in Arkansas, 1942

| Oil yield per acre (pounds) | Ozark-Ouachita | | Highlands-1/ ¹ | | Coastal Plains-2/ ² | | State | |
|--------------------------------|----------------|---------|---------------------------|---------|--------------------------------|---------|------------|---------|
| | Cottonseed | Peanuts | Cottonseed | Peanuts | Cottonseed | Peanuts | Cottonseed | Peanuts |
| 0- 19 | 10 | * | 4 | 4 | 4 | 8 | 2 | |
| 20- 39 | 27 | 7 | 21 | 12 | 24 | 9 | | |
| 40- 59 | 30 | 10 | 34 | 12 | 31 | 11 | | |
| 60- 79 | 20 | 11 | 25 | 12 | 22 | 12 | | |
| 80- 99 | 9 | 13 | 10 | 8 | 10 | 11 | | |
| 100-119 | 3 | 8 | 4 | 12 | 4 | 10 | | |
| 120-139 | 1 | 12 | 2 | 15 | 1 | 12 | | |
| 140-159 | * | 12 | * | 8 | * | 10 | | |
| 160-179 | * | * | 6 | 4 | * | 5 | | |
| 180-199 | 0 | 7 | 4 | 4 | | 5 | | |
| 200-219 | * | * | 3 | 2 | * | 5 | | |
| 220-239 | * | * | 2 | 2 | 2 | 2 | | |
| 240-259 | * | * | 3 | 2 | 2 | 3 | | |
| 260-279 | * | * | 2 | 1 | 1 | 2 | | |
| 280-299 | * | * | * | 1 | * | * | | |
| 300-319 | * | * | 3 | 2 | 2 | 3 | | |
| 320-339 | * | * | * | * | * | ** | | |
| 340-359 | * | * | * | * | * | ** | | |
| 360-379 | * | * | 1 | * | * | ** | | |
| 380-399 | * | * | 0 | * | * | ** | | |
| 400 and over | * | * | * | 1 | 1 | * | | |
| Total | 1 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 687 | — | — | 463 | — | — | 1,150 | — |

Sample counties: 1/ Faulkner, Garland, Izard, Logan, Montgomery, Searcy, Sebastian, Sharp, and Stone.

2/ Columbia, Little River, Miller, Ouachita, and Union.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
September 15, 1943

Table 14.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected Florida counties, 1942

| County and area | Number of farms | Yield per acre, 1942 | | Ratio of oil yield:producing more oil per acre,: per acre from | Percent of farms |
|--------------------------------------|-----------------------|----------------------|---------|--|------------------|
| | | Cotton | Peanuts | | |
| | | outturn from | | | |
| | | | | | |
| Middle Coastal Plain | 743 | 150 | 569 | 41 | 170 |
| Jackson | 300 | 149 | 557 | 41 | 167 |
| Leon | 143 | 91 | 321 | 25 | 93 |
| Santa Rosa | 300 | 185 | 803 | 51 | 241 |
| Rolling Sandy Lands and Flatwoods | | | | | |
| Swannee | 133 | 153 | 431 | 42 | 125 |
| Total | 876 | 151 | 535 | 42 | 159 |
| | | | | | |
| | | | | | |

Table 15.- Frequency distribution of farms by oil yield per acre from cottonseed and peanuts,
selected Florida counties, 1942

| Oil yield per acre (pounds) | Middle Coastal Plain 1/ Cottonseed; Peanuts; | | | Rolling Sandy Lands: and Flatwoods 2/ Cottonseed; Peanuts; | | | State Peanuts | | |
|--------------------------------|---|-----|-----|--|-----|-----|------------------|-----|-----|
| | | | | Percent | | | | | |
| 0- 19 | 18 | 1 | 16 | 2 | 18 | 1 | | | |
| 20- 39 | 35 | 3 | 37 | 8 | 35 | 4 | | | |
| 40- 59 | 24 | 4 | 30 | 9 | 25 | 5 | | | |
| 60- 79 | 16 | 5 | 8 | 9 | 14 | 6 | | | |
| 80- 99 | 5 | 6 | 4 | 12 | 5 | 7 | | | |
| 100-119 | 2 | 8 | 4 | 11 | 2 | 9 | | | |
| 120-139 | * | 9 | 1 | 16 | 1 | 10 | | | |
| 140-159 | * | * | 9 | 12 | * | 10 | | | |
| 160-179 | * | 0 | 9 | 3 | 0 | 7 | | | |
| 180-199 | * | * | 6 | 4 | * | 5 | | | |
| 200-219 | * | * | 6 | 4 | * | 5 | | | |
| 220-239 | * | 0 | 6 | 1 | 0 | 5 | | | |
| 240-259 | * | * | 7 | 2 | | 6 | | | |
| 260-279 | * | * | 5 | 4 | | 5 | | | |
| 280-299 | * | * | 5 | 0 | | 4 | | | |
| 300-319 | * | * | * | 1 | | 3 | | | |
| 320-339 | * | * | * | 1 | | 3 | | | |
| 340-359 | * | * | 2 | 0 | | 2 | | | |
| 360-379 | * | * | 1 | 0 | | 1 | | | |
| 380-399 | * | * | 1 | 1 | | 1 | | | |
| 400 and over | * | * | 1 | | | 1 | | | |
| Total | | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | | 743 | — | — | 133 | — | — | — | 876 |

Sample counties: 1/ Jackson, Leon, and Santa Rosa. 2/ Suwannee.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
October 2, 1943

Table 16.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected Georgia Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | Ratio of oil yield; producing more oil per acre, per acre, | | Percent of farms producing more oil per acre, per acre, to cotton-seed, Peanuts ; Cotton-seed, Peanuts |
|--------------------------|---------------------------|----------------------|---------|--|---------|--|
| | | Cotton | Peanuts | Cotton | Peanuts | |
| Piedmont Plateau | 893 | 245 | 347 | 67 | 104 | 155 |
| Baldwin | 173 | 218 | 320 | 60 | 96 | 160 |
| Coweta | 120 | 276 | 463 | 76 | 139 | 183 |
| McDuffle | 23 | 214 | 327 | 59 | 98 | 166 |
| Morgan | 301 | 352 | 333 | 97 | 100 | 103 |
| Talbot | 276 | 149 | 283 | 41 | 85 | 207 |
| Fall Line Sand Hills | | | | | | |
| Crawford | 182 | 163 | 523 | 45 | 157 | 349 |
| Coastal Plain - Red Belt | | | | | | |
| Sumter | 299 | 229 | 577 | 63 | 173 | 275 |
| Middle Coastal Plain | | | | | | |
| Dooloch | 2,643 | 193 | 680 | 52 | 204 | 392 |
| Burke | 193 | 183 | 900 | 49 | 270 | 551 |
| Coffee | 249 | 239 | 507 | 64 | 152 | 238 |
| Colquitt | 301 | 168 | 743 | 45 | 223 | 496 |
| Early | 295 | 217 | 653 | 58 | 196 | 338 |
| Laurens | 301 | 187 | 757 | 50 | 227 | 454 |
| Lowndes | 297 | 206 | 573 | 55 | 172 | 313 |
| Toombs | 252 | 198 | 710 | 53 | 213 | 402 |
| Wilcox | 196 | 168 | 813 | 45 | 244 | 542 |
| Worth | 259 | 172 | 583 | 46 | 175 | 380 |
| Lower Coastal Plain | 300 | 187 | 687 | 50 | 206 | 412 |
| Pierce | 37 | 236 | 653 | 63 | 196 | 311 |
| Total | 4,054 | 203 | 637 | 55 | 191 | 347 |
| | | | | | | 91 |

Table 17.- Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Georgia, 1942

| Oil yield per acre (pounds) | Piedmont | Fall Line | Coastal Plain | Middle Coastal | Lower Coastal | Plain 4/ Red Belt 3/ | Plain 5/ Cotton: Pea- seed : nuts | Cotton: Pea- seed : nuts | Cotton: Pea- seed : nuts | Cotton: Pea- seed : nuts | State |
|-----------------------------------|---|--|---|---|---|-------------------------|---|-----------------------------|-----------------------------|-----------------------------|-------|
| | Plateau 1/ Cotton: Pea- seed : nuts | Sand Hills 2/ Cotton: Pea- seed : nuts | Plain 4/ Cotton: Pea- seed : nuts | Plain 5/ Cotton: Pea- seed : nuts | Plain 5/ Cotton: Pea- seed : nuts | Percent | | | | | |
| 0- 19 | 5 | 4 | 3 | 0 | 3 | 0 | 5 | * | 5 | 0 | 5 |
| 20- 39 | 20 | 13 | 25 | * | 40 | 0 | 27 | 1 | 16 | 0 | 26 |
| 40- 59 | 21 | 15 | 18 | 2 | 42 | 2 | 35 | 2 | 38 | 0 | 31 |
| 60- 79 | 18 | 15 | 28 | 3 | 9 | 7 | 23 | 4 | 16 | 3 | 21 |
| 80- 99 | 15 | 13 | 17 | 5 | 5 | 10 | 8 | 5 | 5 | 10 | 10 |
| 100-119 | 12 | 11 | 8 | 11 | 0 | 13 | 2 | 6 | 14 | 13 | 4 |
| 120-139 | 6 | 7 | 1 | 17 | 1 | 16 | * | 8 | 3 | 5 | 2 |
| 140-159 | 2 | 7 | * | 11 | * | 16 | * | 9 | 3 | 11 | 1 |
| 160-179 | 1 | 4 | * | 9 | * | 10 | * | 8 | * | 11 | * |
| 180-199 | 0 | 3 | 8 | * | 7 | 0 | 9 | 3 | 0 | 3 | 0 |
| 200-219 | * | * | 2 | * | 13 | 7 | * | 9 | 3 | * | 8 |
| 220-239 | * | * | 2 | 7 | 1 | 7 | 7 | 7 | 8 | 6 | 6 |
| 240-259 | * | * | 2 | 4 | 2 | 2 | 7 | 7 | 11 | 11 | 6 |
| 260-279 | * | * | 1 | 5 | 1 | 1 | 5 | 5 | 5 | 5 | 4 |
| 280-299 | * | * | 1 | 3 | 1 | 1 | 4 | 4 | 3 | 3 | 3 |
| 300-319 | * | * | 2 | * | 2 | 2 | 4 | 4 | 5 | 5 | 3 |
| 320-339 | * | * | * | * | 1 | 1 | 3 | 3 | 0 | 0 | 2 |
| 340-359 | * | * | * | * | * | 1 | 2 | 2 | 5 | 5 | 1 |
| 360-379 | * | * | * | * | * | 0 | 2 | 2 | 3 | 3 | 1 |
| 380-399 | * | * | * | * | 0 | 1 | 3 | 1 | 3 | 1 | 1 |
| 400 and over: | * | * | * | * | * | 2 | 4 | 4 | 1 | 1 | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 893 | 299 | 182 | 2,643 | 37 | 4,054 | | | | | |

Sample counties: 1/ Baldwin, Coweta, McDuffie, Morgan, and Talbot. 2/Crauford, 3/ Sumter.
 4/ Bulloch, Burke, Coffee, Colquitt, Early, Laurens, Lowndes, Toombs, Wilcox, and Worth. 5/ Pierce.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
 October 22, 1943

Table 18.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected Louisiana parishes, 1942

| Parish and area | Number of farms | Yield per acre, 1942 | | Ratio of oil yield:producing more oil per acre,; per acre from | |
|----------------------|-----------------------|----------------------|---------|---|-------------------|
| | | Cotton | Peanuts | outturn from sample | Cotton peanuts |
| | | | | | |
| Coastal Plain | 997 | 164 | 304 | 47 | 91 |
| Allen | 28 | 218 | 590 | 62 | 177 |
| Caddo | 316 | 113 | 193 | 32 | 58 |
| Rapides | 29 | 264 | 487 | 75 | 146 |
| Sabine | 168 | 179 | 420 | 51 | 126 |
| Union | 245 | 250 | 547 | 71 | 164 |
| Webster | 143 | 183 | 243 | 52 | 73 |
| Winn | 68 | 172 | 453 | 49 | 130 |
| Middle Coastal Plain | 305 | 255 | 334 | 68 | 100 |
| St. Helena | 107 | 231 | 337 | 52 | 101 |
| Washington | 198 | 261 | 335 | 70 | 100 |
| Total | 1,302 | 171 | 306 | 48 | 92 |
| | | | | | 192 |
| | | | | | 22 |
| | | | | | 78 |

Table 19.- Frequency distribution of farms by oil yields per acre from cottonseed and peanuts, by areas in Louisiana, 1942

| Oil yield per acre (pounds) | Coastal Plain 1/ | | Middle Coastal Plain 2/ | | State Cottonseed ³ , Peanuts ⁴ , Cottonseed ³ , Peanuts ⁴ , Cottonseed ³ , Peanuts ⁴ |
|--------------------------------|------------------|---------|----------------------------|---------|---|
| | Percent | Percent | Percent | Percent | |
| 0- 19 | 12 | 8 | 5 | 6 | 10 7 |
| 20- 59 | 28 | 9 | 20 | 14 | 25 10 |
| 40- 59 | 26 | 8 | 25 | 14 | 25 10 |
| 60- 79 | 20 | 21 | 26 | 17 | 22 21 |
| 80- 99 | 11 | 13 | 12 | 11 | 11 13 |
| 100-119 | 5 | 9 | 7 | 10 | 5 9 |
| 120-139 | 1 | 7 | 2 | 6 | 1 6 |
| 140-159 | * | 6 | 2 | 7 | 1 6 |
| 160-179 | * | 5 | 0 | 2 | * |
| 180-199 | * | 5 | 0 | 1 | 0 4 |
| 200-219 | * | * | 1 | 4 | * |
| 220-239 | * | * | 2 | 2 | 2 2 |
| 240-259 | * | 1 | 1 | 1 | 1 1 |
| 260-279 | * | 1 | 1 | 1 | 1 1 |
| 280-299 | * | * | 1 | 1 | 1 1 |
| 300-319 | * | * | 1 | 1 | 1 1 |
| 320-339 | * | * | * | 1 | 1 * |
| 340-359 | * | * | * | 1 | 1 * |
| 360-379 | * | * | * | 0 | 0 * |
| 380-399 | * | * | * | 0 | 0 * |
| 400 and over | * | * | * | * | * |
| Total | 100 | 100 | 100 | 100 | 100 100 |
| Number of farms in sample: | 997 | — | — | 505 | — 1,302 |

Sample parishes: 1/ Allen, Caddo, Rapides, Sabine, Union, Webster, and Winn.

2/ St. Helena and Washington.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 9, 1945

Table 20.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected Mississippi counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of oil yield:producing more oil per acre; per acre | | | Percent of farms producing more oil per acre; per acre | | |
|----------------------|---------------------------|----------------------|---------|--------------|--|---------|---------|--|------------|---------|
| | | Cotton | Peanuts | outturn from | Cotton | Peanuts | Percent | Cotton | Peanuts | Percent |
| | | in | lint | sample | seed | seed | Percent | seed | cottonseed | Percent |
| Brown Loams | 1,211 | 277 | 315 | 80 | 94 | 118 | 48 | 52 | 52 | 52 |
| Amite | 270 | 266 | 297 | 76 | 89 | 117 | 48 | 52 | 52 | 52 |
| Hinds | 360 | 319 | 363 | 91 | 109 | 120 | 43 | 57 | 57 | 57 |
| Holmes | 185 | 259 | 233 | 74 | 70 | 95 | 59 | 41 | 41 | 41 |
| Montgomery | 181 | 270 | 307 | 77 | 92 | 119 | 47 | 53 | 53 | 53 |
| Yalobusha | 215 | 298 | 400 | 90 | 120 | 133 | 48 | 52 | 52 | 52 |
| Sand-Clay Hills | 617 | 279 | 343 | 81 | 103 | 127 | 40 | 60 | 60 | 60 |
| Clarke | 165 | 245 | 280 | 70 | 84 | 120 | 47 | 53 | 53 | 53 |
| Neshoba | 307 | 298 | 377 | 85 | 113 | 133 | 35 | 65 | 65 | 65 |
| Pontotoc | 145 | 301 | 390 | 91 | 117 | 129 | 45 | 55 | 55 | 55 |
| Black Belt | 59 | 203 | 253 | 58 | 76 | 131 | 42 | 58 | 58 | 58 |
| Lowndes | | | | | | | | | | |
| Upper Coastal Plain | 227 | 291 | 450 | 88 | 135 | 153 | 35 | 65 | 65 | 65 |
| Itawamba | | | | | | | | | | |
| Middle Coastal Plain | 636 | 294 | 363 | 84 | 109 | 130 | 47 | 53 | 53 | 53 |
| Covington | 293 | 291 | 370 | 83 | 111 | 134 | 41 | 59 | 59 | 59 |
| Greene | 44 | 235 | 393 | 67 | 118 | 176 | 25 | 75 | 75 | 75 |
| Simpson | 299 | 357 | 327 | 102 | 98 | 96 | 56 | 44 | 44 | 44 |
| Total | 2,750 | 279 | 347 | 81 | 104 | 128 | 45 | 55 | 55 | 55 |

Table 21.- Frequency distribution of farms by oil yield per acre from cottonseed and peanuts,
by areas in Mississippi, 1942

| Oil yield per acre (pounds) | Brown : Loams 1/ | | Sand-Clay : Hills 2/ | | Black : Belt 3/ | | Upper Coastal Plain 4/ | | Middle Coastal Plain 5/ | | State | |
|--------------------------------|------------------|-------|----------------------|-----|-----------------|-----|------------------------|-----|-------------------------|-----|-------|-------|
| | Cotton | Pea | Cotton | Pea | Cotton | Pea | Cotton | Pea | Cotton | Pea | seed | nuts |
| 0- 19 | 1 | 9 | 2 | 4 | 7 | 7 | 0 | 2 | 1 | 7 | 1 | 7 |
| 20- 39 | 8 | 15 | 6 | 12 | 22 | 23 | 2 | 10 | 7 | 14 | 7 | 14 |
| 40- 59 | 18 | 12 | 14 | 11 | 25 | 14 | 14 | 6 | 13 | 9 | 16 | 10 |
| 60- 79 | 23 | 18 | 26 | 16 | 27 | 24 | 25 | 11 | 18 | 14 | 23 | 17 |
| 80- 99 | 23 | 9 | 26 | 13 | 10 | 3 | 26 | 9 | 24 | 11 | 24 | 10 |
| 100-119 | 14 | 6 | 15 | 10 | 7 | 10 | 21 | 10 | 18 | 7 | 15 | 7 |
| 120-139 | 8 | 10 | 7 | 11 | 2 | 7 | 7 | 12 | 10 | 11 | 8 | 11 |
| 140-159 | 4 | 7 | 2 | 7 | 7 | 3 | 1 | 9 | 6 | 8 | 4 | 7 |
| 160-179 | 1 | 1 | 1 | 1 | 3 | 3 | 2 | 5 | 2 | 3 | 1 | 3 |
| 180-199 | * | 5 | 1 | 4 | 4 | 2 | 1 | 6 | 1 | 4 | 1 | 4 |
| 200-219 | * | 3 | * | 1 | 2 | 1 | 2 | 1 | * | 3 | 2 | * |
| 220-239 | 0 | 1 | 0 | 2 | 2 | 2 | 2 | 3 | 3 | 2 | 0 | 2 |
| 240-259 | * | 2 | 2 | 0 | * | 1 | 1 | 3 | 3 | 1 | * | 2 |
| 260-279 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 3 | 3 | 2 | * | 1 |
| 280-299 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 0 | * |
| 300-319 | 0 | 1 | 1 | 0 | 0 | 0 | 1 | 5 | 2 | 0 | 2 | * |
| 320-339 | 0 | * | 0 | * | 0 | 0 | 0 | 0 | 0 | 0 | 0 | * |
| 340-359 | * | 0 | * | 0 | * | 0 | 0 | 0 | 0 | 0 | 0 | * |
| 360-379 | * | 0 | * | 0 | * | 1 | 1 | 1 | * | 0 | 0 | 1 |
| 380-399 | * | 0 | * | 0 | * | 0 | 0 | 0 | 0 | * | 0 | * |
| 400 and over | * | * | * | * | * | * | * | 2 | 2 | 1 | * | * |
| Total | : | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | | 1,211 | 617 | 59 | 227 | 636 | | | | | | 2,750 |

Sample counties: 1/ Amite, Hinds, Holmes, Montgomery, and Yalobusha. 2/ Clarke, Neshoba, and Pontotoc.

3/ Lowndes. 4/ Itawamba. 5/ Covington, Greene, and Simpson.

* Less than 5 tenths of 1 percent.

Table 222.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected Oklahoma counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | Ratio of oil yield; producing more oil per acre, per acre from peanuts : Cotton- to : Peanuts | |
|------------------|---------------------------------------|---|--------|---|--------|
| | | Computed oil outturn from Cotton- seed : | | Cotton- seed : cottonseed; | |
| | | Pounds | Pounds | Pounds | Pounds |
| Rolling Plains | 560 | 192 | 622 | 47 | 187 |
| Caddo | 261 | 235 | 807 | 57 | 242 |
| Greer | 169 | 173 | 587 | 42 | 176 |
| Hartman | 24 | 132 | 383 | 32 | 115 |
| Jackson | 106 | 202 | 577 | 49 | 173 |
| Central Prairies | 509 | 162 | 611 | 39 | 153 |
| Grady | 300 | 166 | 523 | 40 | 157 |
| McClain | 209 | 156 | 490 | 38 | 147 |
| Gross Timbers | 1,599 | 140 | 553 | 36 | 166 |
| Carter | 202 | 97 | 433 | 25 | 130 |
| Hughes | 165 | 144 | 620 | 37 | 186 |
| Lincoln | 179 | 133 | 570 | 34 | 171 |
| Logan | 54 | 144 | 530 | 35 | 159 |
| Love | 286 | 133 | 390 | 34 | 117 |
| Oklfuskee | 268 | 162 | 573 | 39 | 172 |
| Payne | 136 | 218 | 460 | 53 | 138 |
| Seminole | 309 | 152 | 647 | 39 | 194 |
| Eastern Prairies | 376 | 148 | 477 | 38 | 143 |
| Muskogee | 222 | 129 | 457 | 33 | 137 |
| Tulsa | 77 | 234 | 537 | 60 | 161 |
| Wagoner | 77 | 148 | 497 | 38 | 149 |
| Ozark-Ouachita | 430 | 152 | 362 | 39 | 106 |
| Lefler | 210 | 136 | 327 | 35 | 98 |
| McCurtain | 220 | 156 | 357 | 40 | 107 |
| Coastal Plains | | | | | |
| Bryan | 288 | 133 | 640 | 34 | 192 |
| Total | 5,762 | 147 | 528 | 37 | 158 |
| | | | | | |
| | | | | | |

Table 23.—Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Oklahoma, 1942

| Oil yield per acre (pounds) | -Percent- | | | | | | State |
|-----------------------------------|----------------------|-----------------------|----------------------|----------------------|-----------------------|----------------------|-------|
| | Rolling Plains 1/ | Central Timbers 2/ | Cross Prairies 3/ | Eastern Timber 4/ | Ozark- Ouachita 5/ | Coastal Plains 6/ | |
| 0- 19 | 19 | 1 | 13 | 3 | 21 | 2 | 24 |
| 20- 39 | 24 | 4 | 47 | 9 | 42 | 4 | 0 |
| 40- 59 | 22 | 5 | 28 | 9 | 26 | 5 | 59 |
| 60- 79 | 17 | 6 | 8 | 10 | 8 | 8 | 6 |
| 80- 99 | 11 | 7 | 2 | 7 | 9 | 7 | 25 |
| 100-119 | 3 | 5 | 1 | 9 | 1 | 24 | 7 |
| 120-139 | 2 | 6 | 1 | 8 | 9 | 11 | 9 |
| 140-159 | 1 | 8 | * | 7 | 8 | 10 | 9 |
| 160-179 | 1 | 5 | * | 6 | 7 | 6 | 8 |
| 180-199 | 1 | 7 | * | 4 | 6 | 5 | 8 |
| 200-219 | 1 | 5 | * | 4 | 0 | 6 | 5 |
| 220-239 | 1 | 5 | 4 | 4 | 0 | 5 | 0 |
| 240-259 | 1 | 7 | 4 | 4 | 3 | 3 | 4 |
| 260-279 | 1 | 4 | 3 | 3 | 3 | 1 | 4 |
| 280-299 | 1 | 4 | 2 | 2 | 2 | 1 | 2 |
| 300-319 | 1 | 3 | 2 | 2 | 2 | 1 | 6 |
| 320-339 | 1 | 4 | 2 | 2 | 1 | 0 | 2 |
| 340-359 | 1 | 2 | 1 | 1 | 1 | * | 2 |
| 360-379 | 1 | 2 | 2 | 1 | 1 | 0 | 1 |
| 380-399 | 1 | 2 | 1 | 1 | 1 | * | 1 |
| 400 and over | 1 | 8 | 3 | 3 | 2 | 1 | 1 |
| Total | 109 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 560 | 509 | 1,599 | 376 | 430 | 288 | 3,762 |

Sample counties: 1/ Caddo, Greer, Harmon, and Jackson. 2/ Grady and McClain. 3/ Carter, Hughes, Lincoln, Logan, Love, Okfuskee, Payne, and Seminole. 4/ Muskogee, Tulsa, and Wagoner.

5/ Latimer and McCurtain. 6/ Bryan.

* Less than 5 tenths of 1 percent.

Table 24.- Comparative data on oil yields per acre from cottonseed and peanuts,
selected South Carolina counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of oil yield:producing more oil per acre, per acre: | | |
|----------------------|---------------------------|----------------------|---------|--------------|---|-------------------|-----------------------|
| | | Cotton | Peanuts | Computed oil | Peanuts | Cotton to Peanuts | Cottonseed to Peanuts |
| Piedmont Plateau | 93 | 310 | 443 | 90 | 128 | 142 | 25 |
| Anderson | 42 | 357 | 700 | 103 | 203 | 197 | 10 |
| Edgefield | 51 | 294 | 355 | 85 | 103 | 121 | 37 |
| Fall Line Sand Hills | 529 | 219 | 512 | 63 | 152 | 241 | 14 |
| Aiken | 233 | 215 | 517 | 62 | 155 | 250 | 8 |
| Chesterfield | 129 | 256 | 463 | 74 | 139 | 188 | 22 |
| Lexington | 167 | 232 | 503 | 67 | 146 | 218 | 16 |
| Middle Coastal Plain | 1,043 | 202 | 412 | 55 | 128 | 233 | 23 |
| Allendale | 267 | 173 | 300 | 47 | 90 | 191 | 25 |
| Barnwell | 291 | 158 | 417 | 43 | 125 | 291 | 11 |
| Clarendon | 120 | 305 | 357 | 83 | 107 | 129 | 47 |
| Horry | 69 | 331 | 910 | 90 | 273 | 303 | 9 |
| Lee | 130 | 206 | 310 | 56 | 93 | 166 | 27 |
| Marion | 166 | 448 | 650 | 122 | 195 | 160 | 29 |
| Lower Coastal Plain | 77 | 218 | 367 | 59 | 106 | 180 | 57 |
| Colleton | 46 | 209 | 383 | 57 | 111 | 195 | 33 |
| Dorchester | 31 | 253 | 310 | 69 | 90 | 130 | 58 |
| Total | 1,742 | 215 | 437 | 60 | 130 | 217 | 21 |
| | | | | | | | 79 |

Table 25.- Frequency distribution of farms by oil yield per acre from cottonseed and peanuts,
selected South Carolina Counties, 1942

| Oil yield per acre (pounds) | Piedmont | Fall Line | Sand Hills 1/ | Plain 3/ | Middle Coastal | Lower Coastal | State |
|--------------------------------|----------------------------|--|----------------|--|-----------------------------|----------------|----------------|
| | Cotton-peanuts | Cotton-peanuts | Cotton-peanuts | Cotton-peanuts | Cotton-peanuts | Cotton-peanuts | Cotton-peanuts |
| | seed | seed | seed | seed | seed | seed | seed |
| 0- 19 | 0 | 0 | 3 | * | 9 | 4 | 1 |
| 20- 39 | 10 | 3 | 16 | 4 | 25 | 12 | 20 |
| 40- 59 | 13 | 7 | 30 | 8 | 22 | 13 | 22 |
| 60- 79 | 14 | 14 | 23 | 12 | 13 | 29 | 16 |
| 80- 99 | 18 | 14 | 16 | 9 | 12 | 10 | 14 |
| 100-119 | 24 | 9 | 6 | 10 | 10 | 9 | 5 |
| 120-139 | 7 | 2 | 4 | 11 | 5 | 8 | 4 |
| 140-159 | 11 | 9 | 1 | 9 | 3 | 6 | 4 |
| 160-179 | 5 | 9 | 1 | 6 | 1 | 5 | 1 |
| 180-199 | 3 | 5 | 0 | 6 | * | 3 | 3 |
| 200-219 | 12 | * | 6 | * | 6 | * | 4 |
| 220-239 | 4 | 0 | 5 | 0 | 3 | 3 | 0 |
| 240-259 | 1 | * | 3 | * | 2 | 3 | 3 |
| 260-279 | 3 | 0 | 1 | 3 | 2 | 3 | 0 |
| 280-299 | 9 | 0 | 1 | 3 | 1 | 3 | 2 |
| 300-319 | 0 | 0 | * | 2 | 1 | 0 | 0 |
| 320-339 | 0 | 0 | * | 1 | 1 | 0 | 1 |
| 340-359 | 0 | 0 | * | 1 | 1 | 1 | 1 |
| 360-379 | 1 | * | 1 | * | 1 | 0 | 1 |
| 380-399 | 0 | * | 1 | 1 | 0 | 0 | 1 |
| 400 and over | * | * | 2 | 3 | 1 | 2 | 2 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | — | — | — | — | — | — | — |
| Sample counties: | 1/ Anderson and Edgefield. | 2/ Aiken, Chesterfield, and Lexington. | 3/ Allendale, | Barnwell, Clarendon, Horry, Lee, and Marion. | 4/ Colleton and Dorchester. | | 1,742 |

* Less than 5 tenths of 1 percent.

Southern Division, AAA, October 20, 1942

Table 26.- Comparative data on oil yields per acre from cottonseed and peanuts, selected Texas Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of oil yield per acre; peanuts to cottonseed | | | Percent of farms producing more oil per acre; peanuts to cottonseed | | |
|------------------|------------------------------------|-------------------------|--------|--------|---|--------|--------|--|---------|---------|
| | | Cotton : Lint : Peanuts | | | Computed oil output from: Cottonseed : Peanuts | | | Peanuts | | |
| | | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Percent | Percent | Percent |
| High Plains | 289 | 222 | 602 | 50 | 181 | 362 | 8 | 92 | | |
| Bailey | 51 | 202 | 680 | 45 | 204 | 453 | 12 | 88 | | |
| Dawson | 80 | 220 | 377 | 49 | 113 | 231 | 16 | 84 | | |
| Lamb | 53 | 186 | 860 | 42 | 258 | 614 | 2 | 98 | | |
| Lubbock | 105 | 260 | 827 | 58 | 248 | 428 | 5 | 95 | | |
| Rio Grande Plain | 309 | 63 | 479 | 14 | 144 | 1,028 | 2 | 98 | | |
| Atascosa | 158 | 63 | 527 | 14 | 158 | 1,129 | 0 | 100 | | |
| Duval | 108 | 63 | 250 | 14 | 75 | 536 | 6 | 94 | | |
| Starr | 45 | 63 | 157 | 14 | 47 | 336 | 16 | 84 | | |
| Edwards Plateau | 295 | 103 | 503 | 23 | 151 | 656 | 2 | 98 | | |
| Gillespie | 63 | 81 | 445 | 18 | 133 | 739 | 2 | 98 | | |
| San Saba | 232 | 116 | 640 | 26 | 162 | 623 | 2 | 98 | | |
| Rolling Plains | 829 | 161 | 454 | 36 | 136 | 378 | 8 | 92 | | |
| Callahan | 203 | 121 | 597 | 27 | 119 | 441 | 6 | 94 | | |
| Cottle | 50 | 202 | 437 | 45 | 131 | 291 | 13 | 87 | | |
| Garza | 102 | 255 | 570 | 57 | 171 | 300 | 9 | 91 | | |
| Mitchell | 150 | 130 | 393 | 29 | 118 | 407 | 5 | 95 | | |
| Stonewall | 169 | 108 | 470 | 24 | 141 | 587 | 5 | 97 | | |
| Wichita | 33 | 291 | 490 | 65 | 147 | 226 | 18 | 82 | | |
| Wilbarger | 122 | 314 | 510 | 70 | 153 | 219 | 15 | 85 | | |
| Cross Timbers | 515 | 92 | 510 | 21 | 153 | 728 | 1 | 99 | | |
| Comanche | 293 | 90 | 510 | 20 | 153 | 765 | 0 | 100 | | |
| Jack | 68 | 108 | 427 | 24 | 128 | 533 | 7 | 93 | | |
| Wise | 154 | 99 | 527 | 22 | 158 | 718 | 2 | 98 | | |
| Grand Prairie | | | | | | | | | | |
| Bosque | 84 | 102 | 350 | 25 | 105 | 420 | 8 | 92 | | |
| Coastal Plain | 1,875 | 125 | 423 | 29 | 127 | 438 | 11 | 89 | | |
| Anderson | 298 | 108 | 293 | 24 | 88 | 367 | 13 | 87 | | |
| Brazos | 75 | 148 | 250 | 33 | 75 | 227 | 20 | 80 | | |
| Franklin | 299 | 148 | 360 | 33 | 108 | 327 | 12 | 88 | | |
| Gonzales | 122 | 74 | 300 | 18 | 90 | 500 | 7 | 93 | | |
| Grayson | 299 | 159 | 747 | 39 | 224 | 574 | 0 | 100 | | |
| Harrison | 270 | 112 | 213 | 25 | 64 | 256 | 21 | 79 | | |
| Lamar | 175 | 139 | 407 | 34 | 122 | 359 | 7 | 93 | | |
| Montgomery | 39 | 76 | 227 | 17 | 68 | 400 | 10 | 90 | | |
| Nacogdoches | 298 | 116 | 297 | 26 | 89 | 342 | 8 | 92 | | |
| Total | 4,196 | 105 | 469 | 24 | 141 | 508 | 6 | 92 | | |

Table 27.- Frequency distribution of farms by oil yield per acre from cottonseed and peanuts, by areas in Texas, 1942

Semple counties: 1/ Bailey, Dawson, Lamb, and Lubbock. 2/ Atascosa, Duval, and Starr.

3 / Gillespi and San Saba. 4 / Callahan, Cottle, Gerza, Mitchell, Stonewall, Wichita, and Wilbarger.
5 / Comanche, Jack, and Wise. 6 / Bosque. 7 / Anderson, Brazos, Franklin, Gonzales, Grayson, Harrison.

Lamar, Montgomery, and Nacogdoches.

Table 28.- Relative advantage of peanuts over cottonseed in per acre oil production,
by areas in Texas, 1942

| Area | Percent of farms producing more than times : times : times : time : time : in as much oil per acre from peanuts as from cottonseed | | | Percent of farms produc- ing more oil : per acre from : cottonseed than : sample from peanuts : | | | Number 289 |
|---------------------|---|----|----|--|----|----|---------------|
| | 6 | 5 | 4 | 3 | 2 | 1 | |
| High Plains 1/ | 30 | 36 | 43 | 60 | 77 | 91 | 9 |
| Rio Grande Plain 2/ | 68 | 75 | 80 | 85 | 90 | 97 | 3 |
| Edwards Plateau 3/ | 57 | 70 | 77 | 84 | 93 | 97 | 3 |
| Rolling Plain 4/ | 26 | 34 | 45 | 61 | 78 | 94 | 6 |
| Cross Timbers 5/ | 64 | 73 | 84 | 91 | 95 | 99 | 1 |
| Grand Prairie 6/ | 26 | 37 | 46 | 64 | 77 | 93 | 7 |
| Coastal Plain 7/ | 26 | 35 | 45 | 58 | 74 | 91 | 9 |
| Total | 36 | 44 | 53 | 64 | 79 | 93 | 7 |
| | | | | | | | 4,196 |

Sample counties: 1/ Bailey, Dawson, Lamb, and Lubbock. 2/ Atascosa, Duval, and Starr.
 3/ Gillespie and San Saba. 4/ Callahan, Cottle, Garza, Mitchell, Stonewall,
 Wichita, and Wilbarger. 5/ Comanche, Jack, and Wise. 6/ Bosque.
 7/ Anderson, Brazos, Franklin, Grayson, Harrison, Gonzales, Lamar,
 Montgomery, and Nacogdoches.

Table 29.- Frequency distribution of farms by ratio of peanut oil yield per acre to cottonseed oil yield per acre, by areas in Texas, 1942

| Ratio Percent | High Plains 1/ Plain 2/ | | Rio Grande Plateau 3/ | | Edwards Plateau 3/ | | Rolling Plains 4/ Cross Timbers 5/ | | Grand Prairie 6/ Coastal Plain 7/ | | State | |
|---------------------------|----------------------------|-----|--------------------------|-----|-----------------------|----|---------------------------------------|-------|--------------------------------------|----|-----------|----|
| | | | | | | | | | | | Percent - | |
| Under 60 | 6 | * | * | * | 3 | * | * | 5 | 5 | 5 | 5 | 5 |
| 60-79 | 2 | 1 | 1 | 2 | 2 | 1 | 2 | 2 | 2 | 3 | 3 | 2 |
| 80-99 | 1 | * | * | 1 | 1 | 1 | 0 | 0 | 5 | 5 | 5 | 2 |
| Under 100 | 9 | 3 | 3 | 5 | 6 | 1 | 7 | 7 | 9 | 7 | 7 | 7 |
| 100-119 | 3 | 2 | 1 | 1 | 3 | 1 | 0 | 4 | 4 | 4 | 4 | 3 |
| 120-139 | 2 | 1 | 1 | 1 | 3 | 1 | 7 | 7 | 4 | 4 | 4 | 3 |
| 140-159 | 5 | 4 | 1 | 1 | 4 | 1 | 2 | 3 | 3 | 3 | 3 | 3 |
| 160-179 | 4 | 2 | 2 | 1 | 3 | 1 | 5 | 5 | 5 | 5 | 5 | 2 |
| 180-199 | 2 | 1 | 1 | 1 | 3 | * | 2 | 3 | 3 | 3 | 3 | 2 |
| 100-199 | 14 | 7 | 4 | 16 | 16 | 4 | 4 | 16 | 17 | 17 | 14 | 14 |
| 200-219 | 4 | 1 | 2 | 5 | 5 | 1 | 1 | 1 | 4 | 4 | 4 | 3 |
| 220-239 | 5 | 2 | 2 | 5 | 5 | 1 | 1 | 1 | 5 | 5 | 5 | 3 |
| 240-259 | 4 | 2 | 2 | 2 | 3 | 1 | 1 | 5 | 5 | 5 | 5 | 3 |
| 260-279 | 1 | 1 | 1 | 1 | 3 | 1 | 5 | 5 | 5 | 5 | 5 | 3 |
| 280-299 | 3 | 1 | 2 | 3 | 2 | 1 | 2 | 2 | 3 | 3 | 3 | 3 |
| 200-299 | 17 | 5 | 9 | 17 | 17 | 4 | 4 | 13 | 16 | 16 | 15 | 15 |
| 300-319 | 4 | 1 | 1 | 1 | 5 | 2 | 4 | 4 | 4 | 4 | 4 | 2 |
| 320-339 | 3 | 1 | 2 | 2 | 5 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 340-359 | 3 | 2 | 1 | 1 | 5 | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| 360-379 | 4 | 1 | 1 | 1 | 5 | 1 | 1 | 4 | 4 | 4 | 4 | 2 |
| 380-399 | 3 | * | 2 | 2 | 5 | 1 | 1 | 4 | 4 | 4 | 4 | 2 |
| 300-399 | 17 | 5 | 7 | 16 | 16 | 7 | 7 | 18 | 18 | 18 | 11 | 11 |
| 400-419 | 2 | 1 | 1 | 1 | 5 | 2 | 2 | 5 | 5 | 5 | 2 | 2 |
| 420-439 | 1 | 1 | 1 | 1 | 5 | 2 | 2 | 0 | 0 | 0 | 2 | 2 |
| 440-459 | 2 | 1 | 1 | 1 | 5 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 460-479 | 1 | 1 | 1 | 1 | 2 | 1 | 5 | 0 | 0 | 0 | 1 | 1 |
| 480-499 | 1 | * | 1 | * | 4 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
| 400-499 | 7 | 6 | 7 | 11 | 11 | 11 | 9 | 10 | 9 | 9 | 9 | 9 |
| 500-519 | 1 | 1 | 1 | 1 | 5 | 2 | 5 | 0 | 0 | 0 | 3 | 2 |
| 520-539 | 1 | 1 | 1 | 1 | 5 | 1 | 1 | 3 | 1 | 1 | 2 | 2 |
| 540-559 | 2 | 2 | 2 | 4 | 2 | 1 | 1 | 4 | 1 | 1 | 2 | 2 |
| 560-579 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 | 2 | 1 |
| 580-599 | 1 | 2 | 1 | 1 | 5 | 1 | 1 | 2 | 2 | 2 | 1 | 1 |
| 500-599 | 6 | 7 | 13 | 8 | 9 | 9 | 11 | 9 | 9 | 8 | 8 | 8 |
| 600 and over | 30 | 68 | 57 | 26 | 64 | 26 | 26 | 26 | 26 | 36 | 36 | 36 |
| Number of farms in sample | 289 | 309 | 295 | 629 | 515 | 84 | 1,875 | 4,196 | | | | |

Sample counties: 1/ Bailey, Cottle, Garza, Mitchell,
2/ Atascosa, Dawson, Lamb, and Lubbock.
3/ Stonewall, Wichita, and Wilbarger.

* Less than 5 tenths of one percent.

Southern Division, AAA
November 3, 1943

4/ Callahan, Erath, Franklin, Grayson, Harrison,
5/ Anderson, Brazos, Franklin, Grayson, Harrison,
6/ Bosque.
7/ Comanche, Jack, and Wise.
Gillespie and San Saba.
Concho, Lamar, Montague, and Nacogdoches.

Table 30.—Comparative data on meal yields per acre from cottonseed and peanuts,
by areas, Southern Region, 1942

| Number and name of area 1/ | Number of farms in sample | Yield per acre, 1942 | | | Ratio of meal yield producing more meal per acre, per acre from | | |
|---------------------------------------|---------------------------|----------------------|------|---------------|---|-----------------|--------------------------|
| | | Cotton Peanuts | | Computed meal | Cotton Peanuts | Peanuts to seed | Cottonseed to cottonseed |
| | | Cotton | Lint | outturn from | Cotton Peanuts | Peanuts | Peanuts |
| 1. High Plains | 289 | 222 | 602 | 154 | 301 | 195 | 18 |
| 2. Rio Grande Plain | 309 | 63 | 479 | 44 | 240 | 545 | 82 |
| 3. Edwards Plateau | 295 | 103 | 503 | 71 | 252 | 355 | 89 |
| 4. Rolling Plains | 1,390 | 184 | 579 | 134 | 290 | 216 | 7 |
| 5. Oklahoma Central Prairies | 509 | 162 | 511 | 127 | 256 | 202 | 80 |
| 6. East Oklahoma Prairies | 382 | 148 | 477 | 123 | 238 | 193 | 22 |
| 7. Cross Timbers | 2,114 | 133 | 547 | 105 | 274 | 261 | 22 |
| 8. Grand Prairie | 84 | 102 | 350 | 78 | 175 | 224 | 78 |
| 9. Coastal Plain | 3,623 | 144 | 537 | 109 | 268 | 246 | 77 |
| 10. Ozark-Ouachita Highlands | 1,117 | 160 | 370 | 134 | 185 | 138 | 77 |
| 11. Brown Loams | 1,212 | 277 | 315 | 208 | 135 | 19 | 86 |
| 12. Sand-Clay Hills | 704 | 276 | 341 | 209 | 147 | 70 | 86 |
| 13. Black Belt | 555 | 187 | 236 | 142 | 101 | 71 | 86 |
| 14. Upper Coastal Plain | 979 | 267 | 385 | 192 | 165 | 86 | 81 |
| 15. Limestone Basin | 300 | 311 | 710 | 216 | 305 | 141 | 70 |
| 16. Sand Mountain | 597 | 490 | 846 | 341 | 364 | 107 | 66 |
| 17. Appalacian Highlands | 133 | 250 | 305 | 174 | 131 | 75 | 67 |
| 18. Piedmont Plateau | 1,606 | 251 | 355 | 185 | 153 | 83 | 67 |
| 19. Fall Line Sand Hills | 711 | 181 | 520 | 144 | 224 | 156 | 66 |
| 20. Coastal Plain - Red Belt | 299 | 229 | 677 | 165 | 248 | 150 | 65 |
| 21. Middle Coastal Plain | 6,259 | 192 | 662 | 142 | 285 | 201 | 65 |
| 22. Lower Coastal Plain | 114 | 225 | 446 | 168 | 192 | 114 | 65 |
| 23. Rolling Sandy Lands and Flatwoods | 133 | 153 | 431 | 115 | 185 | 164 | 65 |
| Southern Region | 23,715 | 188 | 584 | 141 | 269 | 191 | 74 |
| | | | | | | | 64 |

1/ Numbers correspond with area numbers on map in this report.

Table 31.- Comparative data on meal yields per acre from cottonseed and peanuts,
by States, Southern Region, 1942

| State | Number of farms | Yield per acre, 1942 | | | Ratio of : Percent of farms meal yield producing more meal per acre,: | | |
|-----------------|-----------------------|--------------------------------|------------------------|---------------------------|---|----------------------------------|---------|
| | | Cotton-Peanuts: in sample : | outturn from lint : | Cotton-Peanuts: Seed : | peanuts : to seed : | Cotton- seed : cottonseed: | Percent |
| Alabama | 3,873 | 179 | 593 | 130 | 255 | 196 | 47 |
| Arkansas | 1,150 | 188 | 396 | 158 | 198 | 125 | 38 |
| Florida | 876 | 151 | 535 | 112 | 230 | 205 | 15 |
| Georgia | 4,054 | 203 | 637 | 147 | 274 | 186 | 25 |
| Louisiana | 1,302 | 171 | 306 | 132 | 153 | 116 | 41 |
| Mississippi | 2,751 | 279 | 347 | 211 | 149 | 71 | 76 |
| Oklahoma | 3,769 | 147 | 528 | 120 | 264 | 220 | 20 |
| South Carolina | 1,742 | 215 | 437 | 169 | 188 | 111 | 51 |
| Texas | 4,196 | 105 | 469 | 74 | 234 | 316 | 19 |
| Southern Region | 23,713 | 188 | 584 | 141 | 269 | 191 | 36 |
| | | | | | | | 64 |

Table 32.—Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by States, Southern Region, 1942

* Less than 5 tenths of 1 percent.

Southern Division, AIA
November 10, 1943

Table 330.- Comparative data on meal yields per acre from cottonseed and peanuts,
selected Alabama Counties, 1942

| County and area | Number of farms | Yield per acre, 1942 | | Meal yield producing more meal per acre from cottonseed | | Ratio of meal yield to peanut yield; Cottonseed : Peanuts | | Percent of farms per acre from cottonseed | |
|-------------------------------|-----------------|----------------------|---------|---|---------------------|---|------------|---|---------|
| | | Cotton | Peanuts | Computed meal | outturn from cotton | Cotton | Peanuts | Cotton | Peanuts |
| | | in sample | lint | sample | seed | to seed | cottonseed | seed | Percent |
| | | Number | Pounds | Pounds | Pounds | Pounds | Pounds | Percent | Percent |
| Limestone Basin Limestone | 300 | 311 | 710 | 216 | 305 | 141 | 26 | 74 | |
| Upper Coastal Plain Elmore | 752 | 249 | 332 | 173 | 143 | 83 | 67 | 33 | |
| Franklin | 251 | 205 | 503 | 143 | 130 | 91 | 59 | 41 | |
| Lamar | 206 | 334 | 437 | 232 | 188 | 81 | 67 | 33 | |
| Sand Mountain Cullman | 295 | 295 | 333 | 205 | 143 | 70 | 75 | 25 | |
| DeKalb | 597 | 490 | 846 | 341 | 364 | 107 | 51 | 49 | |
| Appalachian Highlands Calhoun | 250 | 250 | 305 | 174 | 131 | 75 | 69 | 31 | |
| Shelby | 74 | 268 | 313 | 187 | 135 | 72 | 69 | 31 | |
| Piedmont Plateau Lee | 59 | 217 | 290 | 151 | 125 | 83 | 68 | 32 | |
| Randolph | 619 | 227 | 283 | 163 | 122 | 75 | 66 | 34 | |
| Black Belt Hale | 350 | 125 | 190 | 90 | 82 | 91 | 58 | 42 | |
| Lowndes | 269 | 254 | 307 | 182 | 132 | 73 | 71 | 29 | |
| Sand-Clay Hills Clarke | 496 | 178 | 227 | 135 | 98 | 73 | 71 | 29 | |
| Middle Coastal Plain Coffee | 254 | 231 | 190 | 176 | 82 | 47 | 84 | 16 | |
| Conecuh | 242 | 149 | 247 | 113 | 106 | 94 | 48 | 52 | |
| Total | 3,873 | 179 | 593 | 130 | 255 | 196 | 48 | 52 | |

Southern Division, AAA
October 19, 1943

Table 34.—Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, by areas in Alabama, 1942

| Meal yield per acre (pounds) | Limestone | | Upper | | Sand | | Appalachian | | Piedmont | | Black | | Sand-Clay | | Middle | | State | |
|------------------------------------|-----------|---------|----------|----------|---------|-------|-------------|-------|----------|-------|-------|------|-----------|------|--------|------|-------|------|
| | Basin | Coastal | Mountain | Highland | Plateau | Hills | Belt | Hills | Coastal | Plain | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- |
| 0- 49 | 0 | 0 | 2 | 16 | 0 | * | 2 | 19 | 13 | 28 | 14 | 32 | 22 | 30 | 17 | 1 | 9 | 13 |
| 50- 99 | 2 | 0 | 13 | 24 | * | 2 | 11 | 27 | 32 | 31 | 27 | 36 | 36 | 31 | 30 | 4 | 19 | 17 |
| 100-149 | 16 | 5 | 18 | 22 | 1 | 9 | 30 | 20 | 22 | 19 | 22 | 14 | 21 | 15 | 26 | 11 | 19 | 14 |
| 150-199 | 27 | 20 | 22 | 15 | 6 | 12 | 23 | 9 | 16 | 10 | 14 | 8 | 12 | 18 | 16 | 13 | 17 | 13 |
| 200-249 | 24 | 16 | 24 | 8 | 9 | 10 | 20 | 14 | 8 | 7 | 10 | 6 | 3 | 2 | 8 | 15 | 13 | 10 |
| 250-299 | 20 | 9 | 11 | 6 | 19 | 12 | 9 | 4 | 6 | 2 | 6 | 2 | 3 | 1 | 2 | 14 | 9 | 8 |
| 300-349 | 8 | 16 | 6 | 4 | 22 | 12 | 2 | 4 | 2 | 1 | 2 | 4 | 2 | 1 | 1 | 13 | 6 | 7 |
| 350-399 | 2 | 11 | 3 | 2 | 17 | 9 | 1 | 2 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 9 | 4 | 6 |
| 400-449 | 1 | 10 | 1 | 2 | 13 | 8 | 1 | 7 | 9 | 0 | 1 | 1 | 1 | 0 | 0 | 3 | 2 | 4 |
| 450-499 | 7 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | 1 | 1 | 1 |
| 500-549 | * | 4 | * | * | * | * | * | * | * | * | * | * | * | * | * | 1 | 2 | 1 |
| 550-599 | * | 2 | * | * | * | * | * | * | * | * | * | * | * | * | * | 1 | 1 | 1 |
| 600-649 | * | 2 | * | * | * | * | * | * | * | * | * | * | * | * | * | 0 | * | * |
| 650-699 | * | 2 | * | * | * | * | * | * | * | * | * | * | * | * | * | 0 | * | * |
| 700-749 | * | 2 | * | * | * | * | * | * | * | * | * | * | * | * | * | 0 | * | * |

| | | | | | | | | | | | | | | | |
|------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-------|-----|
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms: | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — |
| in sample : | 300 | 752 | 597 | — | 133 | — | 619 | 496 | — | 87 | — | 889 | — | 5,873 | — |

* Less than 5 tenths of 1 percent.

Southern Division, AAA
October 29, 1943

Table 35.- Comparative data on meal yields per acre from cottonseed and peanuts,
selected Arkansas Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of : Percent of farms meal yield:producing more meal per acre,: per acre from | | |
|--------------------------|---------------------------------------|----------------------|---------|--|---|-----------------------|---------|
| | | Cotton | Peanuts | Computed meal outturn from Cotton- seed | Peanuts to cottonseed: | Cotton- to seed | Peanuts |
| Ozark-Ouachita Highlands | 687 | 184 | 427 | 155 | 214 | 138 | 30 |
| Faulkner | 162 | 226 | 537 | 190 | 268 | 141 | 17 |
| Garland | 29 | 137 | 330 | 115 | 165 | 143 | 48 |
| Izard | 40 | 256 | 390 | 215 | 195 | 91 | 67 |
| Logan | 209 | 150 | 363 | 126 | 182 | 144 | 33 |
| Montgomery | 69 | 130 | 507 | 109 | 254 | 233 | 17 |
| Searcy | 40 | 130 | 433 | 109 | 216 | 198 | 32 |
| Sebastian | 67 | 140 | 437 | 118 | 218 | 185 | 21 |
| Sharp | 35 | 294 | 487 | 247 | 244 | 99 | 63 |
| Stone | 36 | 174 | 413 | 146 | 206 | 141 | 31 |
| Coastal Plains | 463 | 191 | 367 | 161 | 184 | 114 | 49 |
| Columbia | 121 | 208 | 393 | 175 | 196 | 112 | 50 |
| Little River | 52 | 137 | 307 | 115 | 154 | 134 | 38 |
| Miller | 57 | 185 | 337 | 156 | 168 | 108 | 56 |
| Ouachita | 89 | 198 | 337 | 167 | 168 | 101 | 55 |
| Union | 144 | 212 | 437 | 178 | 218 | 122 | 45 |
| Total | 1,150 | 188 | 396 | 158 | 198 | 125 | 38 |
| | | | | | | 62 | |

Table 36.- Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, by areas in Arkansas, 1942

| Meal yield per acre (pounds) | Ozark-Ouachita : Highlands 1/ : | | | Coastal Plains 2/ : | | | State | | |
|---|---|---------|--------------------|---------------------|---------|--------------------|------------|---------|--------------------|
| | Cottonseed | Peanuts | Cottonseed:Peanuts | Cottonseed | Peanuts | Cottonseed:Peanuts | Cottonseed | Peanuts | Cottonseed:Peanuts |
| 0- 49 | : | 8 | 1 | 3 | 9 | 6 | 4 | | |
| 50- 99 | : | 25 | 17 | 15 | 19 | 20 | 18 | | |
| 100-149 | : | 24 | 14 | 30 | 15 | 26 | 15 | | |
| 150-199 | : | 22 | 18 | 24 | 18 | 23 | 17 | | |
| 200-249 | : | 13 | 16 | 17 | 17 | 14 | 16 | | |
| 250-299 | : | 7 | 13 | 7 | 8 | 7 | 11 | | |
| 300-349 | : | 2 | 9 | 5 | 5 | 5 | 7 | | |
| 350-399 | : | 1 | 3 | 1 | 2 | 1 | 3 | | |
| 400-449 | : | * | 4 | * | 3 | * | 3 | | |
| 450-499 | : | * | 1 | 1 | 1 | * | 2 | | |
| 500-549 | : | * | 3 | 2 | * | | 3 | | |
| 550-599 | : | * | * | 1 | * | | 1 | | |
| 600-649 | : | * | 1 | * | * | | * | | |
| 650-699 | : | * | * | * | * | | * | | |
| 700-749 | : | * | * | * | * | | * | | |
| 750 and over | : | * | * | * | * | | * | | |
| Total | : | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | | — | — | — | — | — | — | — | — |
| Sample counties: | 1/ Faulkner, Carroll, Izard, Logan, Montgomery, Searcy, Sebastian, Sharp, and Stone. | 687 | — | 463 | — | — | — | 1,150 | — |
| 2/ Columbia, Little River, Miller, Ouachita, and Union. | | | | | | | | | |

* Less than 5 tenths of 1 percent.

Table 37.- Comparative data on meal yields per acre from cottonseed and peanuts, selected Florida Counties, 1942

| County and area | | Number of farms in sample | Yield per acre, 1942 | Computed meal | outturn from cotton; Peanuts; Cotton-lint; Peanuts; seed | Peanuts; Cotton-to; Peanuts; seed | Percent | Percent | Percent | Ratio of : Percent of farms : meal yield; producing more meal per acre; per acre from |
|-----------------------------------|------------|---------------------------|----------------------|---------------|--|-----------------------------------|---------|---------|---------|---|
| | | Number | Pounds | Pounds | Pounds | Pounds | Percent | Percent | Percent | |
| Middle Coastal Plain- | Jackson | 743 | 150 | 569 | 111 | 245 | 221 | 10 | 90 | |
| | Leon | 300 | 149 | 557 | 110 | 240 | 218 | 7 | 93 | |
| | Santa Rosa | 143 | 91 | 321 | 67 | 138 | 206 | 24 | 76 | |
| | | 300 | 185 | 803 | 137 | 345 | 252 | 6 | 94 | |
| Rolling Sandy Lands and Flatwoods | Swankee | 133 | 153 | 431 | 113 | 185 | 164 | 26 | 74 | |
| Total | | 876 | 151 | 535 | 112 | 230 | 205 | 15 | 85 | |

Table 38.- Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, selected Florida Counties, 1942

| Meal yield per acre (pounds) | Middle Coastal Plain 1/ | | | Rolling Sandy Lands: and Flatwoods 2/ | | | State | | |
|---------------------------------|-------------------------|---------|------------|--|------------|---------|------------|---------|------------|
| | Cottonseed | Peanuts | Cottonseed | Peanuts | Cottonseed | Peanuts | Cottonseed | Peanuts | Cottonseed |
| 0- 49 | 16 | 3 | 14 | 8 | 16 | 4 | 4 | 4 | 4 |
| 50- 99 | 33 | 7 | 34 | 14 | 33 | 8 | 8 | 8 | 8 |
| 100-149 | 23 | 11 | 30 | 20 | 24 | 12 | 12 | 12 | 12 |
| 150-199 | 18 | 15 | 13 | 21 | 17 | 16 | 16 | 16 | 16 |
| 200-249. | 6 | 16 | 2 | 19 | 6 | 16 | 16 | 16 | 16 |
| 250-299 | 3 | 11 | 4 | 5 | 3 | 11 | 11 | 11 | 11 |
| 300-349 | 1 | 11 | 2 | 4 | 1 | 10 | 10 | 10 | 10 |
| 350-399 | * | 9 | 1 | 5 | * | 8 | 8 | 8 | 8 |
| 400-449 | * | * | 8 | 1 | * | 7 | 7 | 7 | 7 |
| 450-499 | * | 0 | 5 | 2 | 0 | 5 | 5 | 5 | 5 |
| 500-549 | * | * | 2 | 0 | * | 2 | 2 | 2 | 2 |
| 550-599 | * | * | 1 | 1 | * | 1 | 1 | 1 | 1 |
| 600-649 | * | 0 | * | * | 0 | 0 | * | * | * |
| 650-699 | * | * | * | * | * | * | * | * | * |
| 700-749 | * | * | 1 | * | * | * | * | * | * |
| 750 and over | * | * | * | * | * | * | * | * | * |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 743 | — | — | — | 133 | — | — | — | 876 |

Sample counties: 1/ Jackson, Leon, and Santa Rosa.

2/ Suwannee.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
October 2, 1943

Table 39.- Comparative data on meal yields per acre from cottonseed and peanuts,
selected Georgia Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of : Percent of farms meal yield:producing more meal per acre, ! per acre from peanuts : | | |
|--------------------------|---------------------------------------|---|--------|-------------------|---|-------------------|--------------------------|
| | | Computed meal outturn from Cotton;Peanuts ; | | Cotton- seed ; | Peanuts ; Peanuts seed ; | Cotton- seed ; | Peanuts ; Cottonseed; |
| | | Number | Pounds | Pounds | Pounds | Percent | Percent |
| Piedmont Plateau | 893 | 245 | 347 | 182 | 149 | 82 | 66 |
| Baldwin | 173 | 218 | 320 | 162 | 138 | 85 | 66 |
| Coweta | 120 | 276 | 463 | 205 | 199 | 97 | 58 |
| McDuffie | 23 | 214 | 327 | 159 | 141 | 87 | 70 |
| Morgan | 301 | 352 | 333 | 261 | 143 | 55 | 86 |
| Talbot | 276 | 149 | 283 | 110 | 122 | 111 | 48 |
| Fall Line Sand Hills | 182 | 163 | 523 | 121 | 225 | 186 | 12 |
| Crawford | | | | | | | 88 |
| Coastal Plain - Red Belt | | | | | | | |
| Sumter | 299 | 229 | 577 | 165 | 248 | 150 | 23 |
| Middle Coastal Plain | 2,643 | 193 | 680 | 139 | 292 | 210 | 12 |
| Bullock | 193 | 183 | 900 | 132 | 387 | 298 | 2 |
| Burke | 249 | 239 | 507 | 172 | 218 | 127 | 39 |
| Coffee | 301 | 168 | 743 | 121 | 319 | 264 | 5 |
| Colquitt | 295 | 217 | 653 | 156 | 281 | 180 | 15 |
| Early | 301 | 187 | 757 | 135 | 326 | 241 | 4 |
| Laurens | 297 | 206 | 573 | 148 | 246 | 166 | 22 |
| Lowndes | 252 | 198 | 710 | 142 | 305 | 215 | 14 |
| Toombs | 196 | 168 | 813 | 121 | 350 | 289 | 7 |
| Wilcox | 259 | 172 | 583 | 124 | 251 | 202 | 11 |
| Worth | 300 | 187 | 687 | 135 | 295 | 219 | 5 |
| Lower Coastal Plain | | | | | | | |
| Pierce | 37 | 236 | 653 | 170 | 281 | 165 | 19 |
| Total | 4,054 | 203 | 637 | 147 | 274 | 186 | 25 |

Table 40.—Frequency distribution of meal yields per acre from cottonseed and peanuts,
by areas in Georgia, 1942

| Meal yield per acre (pounds) | Piedmont | Fall Line | Coastal Plain | Middle Coastal | Lower Coastal | State |
|---|-----------------------------|-------------|---------------|----------------|---------------|-------|
| | Plateau 1/ Sand Hills 2/ | Red Belt 3/ | Plain 4/ | Plain 5/ | | |
| Cotton: Pea- seed : nuts : seed : nuts | | | | | | |
| | | | | | Percent | |
| 0- 49 | 4 | 13 | 0 | 2 | * | 4 |
| 50- 99 | 18 | 27 | 5 | 21 | 4 | 4 |
| 100-149 | 19 | 23 | 42 | 17 | 9 | 9 |
| 150-199 | 19 | 14 | 15 | 26 | 24 | 12 |
| 200-249 | 15 | 10 | 4 | 25 | 19 | 14 |
| 250-299 | 13 | 5 | 2 | 12 | 11 | 14 |
| 300-349 | 8 | 4 | 1 | 6 | 2 | 11 |
| 350-399 | 3 | 2 | 2 | 1 | 9 | 8 |
| 400-449 | 1 | 1 | 2 | 1 | * | 1 |
| 450-499 | * | 1 | 1 | 1 | * | 5 |
| 500-549 | * | * | * | 1 | * | 3 |
| 550-599 | * | * | * | 1 | * | 2 |
| 600-649 | * | 0 | 0 | 1 | 1 | 1 |
| 650-699 | * | 0 | 1 | 1 | 1 | 1 |
| 700-749 | * | 0 | 0 | 0 | * | |
| 750 and over | * | 1 | | 1 | 1 | |
| Total | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 893 | 182 | 299 | 2,643 | 37 | 4,054 |

Sample counties: 1/ Baldwin, Coweta, McDuffie, Morgan, and Talbot. 2/ Crawford. 3/ Sumter.
4/ Bulloch, Burke, Coffee, Colquitt, Early, Laurens, Lowndes, Toombs, Wilcox, and Worth. 5/ Pierce.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 1, 1943

Table 41.—Comparative data on meal yields per acre from cottonseed and peanuts,
selected Louisiana parishes, 1942

| Parish and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of : Percent of farms meal yield:producing more meal per acre,: per acre from | | |
|----------------------|---------------------------------------|----------------------|------|--|---|-----------------|-----------------|
| | | Cotton: Peanuts | | Computed meal outturn from Cotton: Peanuts | Peanuts | Cotton: Peanuts | Cotton: Peanuts |
| | | Lint | Seed | Seed | Seed | Cottonseed | Cottonseed |
| Coastal Plain | 997 | 164 | 304 | 128 | 152 | 119 | 36 |
| Allen | 28 | 218 | 590 | 171 | 295 | 173 | 18 |
| Caddo | 316 | 113 | 193 | 88 | 96 | 109 | 36 |
| Rapides | 29 | 264 | 487 | 207 | 244 | 118 | 41 |
| Sabine | 168 | 179 | 420 | 140 | 210 | 150 | 34 |
| Union | 245 | 250 | 547 | 196 | 274 | 140 | 25 |
| Webster | 143 | 183 | 243 | 143 | 122 | 85 | 63 |
| Winn | 68 | 172 | 433 | 135 | 216 | 160 | 31 |
| Middle Coastal Plain | 305 | 255 | 334 | 189 | 167 | 88 | 57 |
| St. Helena | 107 | 231 | 337 | 171 | 168 | 98 | 43 |
| Washington | 198 | 261 | 333 | 193 | 166 | 86 | 65 |
| Total | 1,302 | 171 | 306 | 132 | 153 | 116 | 41 |
| | | | | | | | 59 |

Table 42.- Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Louisiana, 1942.

| Meal yield per acre (pounds) | Percent | | |
|---------------------------------|------------------|-------------------------|---------|
| | Cottonseed | | Peanuts |
| | Coastal Plain 1/ | Middle Coastal Plain 2/ | State |
| 0- 49 | 11 | 11 | 9 |
| 50- 99 | 25 | 13 | 14 |
| 100-149 | 23 | 24 | 20 |
| 150-199 | 23 | 18 | 21 |
| 200-249 | 11 | 8 | 17 |
| 250-299 | 6 | 10 | 9 |
| 300-349 | 1 | 6 | 4 |
| 350-399 | * | 4 | 4 |
| 400-449 | * | 3 | 1 |
| 450-499 | * | 1 | 0 |
| 500-549 | * | 1 | 1 |
| 550-599 | * | * | 1 |
| 600-649 | * | * | 0 |
| 650-699 | * | * | 0 |
| 700-749 | * | 0 | * |
| 750 and over | * | 1 | * |
| Total | 100 | 100 | 100 |
| Number of farms in sample: | 997 | 305 | 1,302 |

Sample parishes: 1/ Allen, Caddo, Sabine, Union, Webster, and Winn.
2/ St. Helena and Washington.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 10, 1943

Table 43.- Comparative data on meal yields per acre from cottonseed and peanuts,
selected Mississippi Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | Ratio of meal yields:producing more meal per acre, per acre from | | Percent of farms producing more meal |
|----------------------|---------------------------|----------------------|---------|--|---------|--------------------------------------|
| | | Cotton | Peanuts | Cotton | Peanuts | |
| | | lint | seed | outturn from cotton | to seed | |
| Brown Loams | 1,212 | 277 | 315 | 209 | 135 | 65 |
| Amitite | 270 | 266 | 297 | 198 | 128 | 65 |
| Hinds | 360 | 319 | 363 | 238 | 156 | 78 |
| Holmes | 185 | 259 | 235 | 193 | 100 | 66 |
| Montgomery | 182 | 270 | 307 | 201 | 132 | 52 |
| Yalobusha | 215 | 298 | 400 | 235 | 172 | 66 |
| Send-Clay Hills | 617 | 279 | 343 | 211 | 147 | 73 |
| Clarke | 165 | 245 | 280 | 182 | 120 | 74 |
| Neshoba | 307 | 298 | 377 | 222 | 162 | 75 |
| Pontotoc | 145 | 301 | 390 | 237 | 168 | 71 |
| Black Belt | | | | | | |
| Lowndes | 59 | 203 | 253 | 151 | 109 | 72 |
| Upper Coastal Plain | | | | | | |
| Itawamba | 227 | 291 | 450 | 229 | 194 | 85 |
| Middle Coastal Plain | | | | | | |
| Covington | 636 | 294 | 363 | 219 | 156 | 71 |
| Greene | 293 | 291 | 370 | 217 | 159 | 73 |
| Simpson | 44 | 235 | 393 | 179 | 169 | 94 |
| Total | 2,751 | 279 | 347 | 211 | 149 | 71 |
| | | | | | | 76 |
| | | | | | | 24 |

Table 44.- Frequency distribution of farms by meal yields per acre from cottonseed and peanuts,
by areas in Mississippi, 1942

| Meal yield per acre (pounds) | Brown | | Sand-Clay | | Black | | Upper Coastal: Middle Coastal: | | State |
|---------------------------------|----------|----------|-----------|--|----------|----------|--------------------------------|-------------|-------|
| | Loams 1/ | Hills 2/ | Belt 3/ | Cotton-; Pea-; Cotton-; Pea-; Cotton-; Pea-; | Plain 4/ | Plain 5/ | nuts : seed | nuts : seed | |
| Percent - - - - - | | | | | | | | | |
| 0- 49 | 1 | 20 | 1 | 14 | 7 | 20 | 0 | 1 | 18 |
| 50- 99 | 7 | 25 | 6 | 22 | 19 | 35 | 2 | 16 | 24 |
| 100-149 | 17 | 18 | 12 | 24 | 29 | 19 | 12 | 20 | 19 |
| 150-199 | 21 | 14 | 24 | 17 | 25 | 14 | 23 | 16 | 17 |
| 200-249 | 24 | 9 | 25 | 9 | 10 | 7 | 27 | 13 | 22 |
| 250-299 | 15 | 6 | 18 | 6 | 8 | 3 | 22 | 8 | 25 |
| 300-349 | 9 | 5 | 8 | 4 | 0 | 2 | 9 | 6 | 10 |
| 350-399 | 4 | 1 | 5 | 1 | 2 | | 2 | 3 | 4 |
| 400-449 | 2 | 1 | 1 | 2 | * | | 1 | 6 | 5 |
| 450-499 | * | * | 1 | 1 | * | | 1 | 0 | 2 |
| 500-549 | * | * | 1 | 1 | 1 | | 1 | * | * |
| 550-599 | * | * | 0 | * | 0 | * | 1 | * | * |
| 600-649 | * | * | 0 | 0 | * | | 1 | * | * |
| 650-699 | * | * | 0 | 0 | 0 | | 1 | * | 0 |
| 700-749 | * | 0 | * | 0 | 0 | | * | * | * |
| 750 and over | * | * | * | * | * | | * | * | * |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample : | 1,212 | — | 617 | — | 59 | — | 227 | 636 | 2,751 |

Sample counties: 1/ Amite, Hinds, Holmes, Montgomery, and Yalobusha.

2/ Clarke, Neshoba, and Pontotoc.

3/ Lowndes.

4/ Itawamba.

5/ Covington, Greene, and Simpson.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 9, 1943

Table 45. Comparative data on meal yields per acre from cottonseed and peanuts,
selected Oklahoma Counties, 1942

| County and area | Number: | | | Yield per acre, 1942 | | | Ratio of : Percent of farms of farms in : Cotton: Peanuts: outturn from sample : lint : cotton-peanuts : seed : cottonseed: | | | meal yield: producing more meal per acre, per acre from peanuts : cotton-to : seed : peanuts : cottonseed: | | |
|------------------|---------|--------|--------|----------------------|--------|--------|--|---------|---------|--|---------|---------|
| | | | | | | | | | | | | |
| | Number | Pounds | Pounds | Pounds | Pounds | Pounds | Percent | Percent | Percent | Percent | Percent | Percent |
| Rolling Plains | 561 | 192 | 622 | 151 | 311 | 206 | 16 | 84 | | | | |
| Caddo | 262 | 235 | 807 | 185 | 404 | 218 | 14 | 86 | | | | |
| Greer | 169 | 173 | 587 | 136 | 294 | 216 | 18 | 82 | | | | |
| Harmon | 24 | 132 | 383 | 104 | 192 | 185 | 8 | 92 | | | | |
| Jackson | 106 | 202 | 577 | 159 | 288 | 181 | 21 | 79 | | | | |
| Central Prairies | 509 | 162 | 511 | 127 | 256 | 202 | 22 | 78 | | | | |
| Grady | 300 | 165 | 523 | 130 | 262 | 202 | 22 | 78 | | | | |
| McClain | 209 | 156 | 490 | 123 | 245 | 199 | 22 | 78 | | | | |
| Cross Timbers | 1,599 | 140 | 553 | 116 | 276 | 238 | 18 | 82 | | | | |
| Carter | 202 | 97 | 433 | 81 | 216 | 267 | 11 | 89 | | | | |
| Hughes | 166 | 144 | 620 | 120 | 310 | 258 | 11 | 89 | | | | |
| Lincoln | 178 | 133 | 570 | 110 | 285 | 259 | 13 | 87 | | | | |
| Logan | 54 | 144 | 530 | 113 | 265 | 235 | 22 | 78 | | | | |
| Love | 286 | 133 | 390 | 110 | 195 | 177 | 28 | 72 | | | | |
| Oklfuskee | 268 | 152 | 573 | 126 | 286 | 227 | 14 | 86 | | | | |
| Payne | 136 | 218 | 460 | 171 | 230 | 134 | 38 | 62 | | | | |
| Seminole | 309 | 152 | 647 | 126 | 324 | 257 | 13 | 87 | | | | |
| Eastern Prairies | 382 | 148 | 477 | 123 | 238 | 193 | 23 | 77 | | | | |
| Muskogee | 228 | 129 | 457 | 107 | 228 | 213 | 21 | 79 | | | | |
| Tulsa | 77 | 234 | 537 | 194 | 268 | 138 | 32 | 68 | | | | |
| Wagoner | 77 | 148 | 497 | 123 | 248 | 202 | 21 | 79 | | | | |
| Ozark-Ouachita | 430 | 152 | 352 | 126 | 176 | 140 | 33 | 67 | | | | |
| Latimer | 210 | 136 | 327 | 113 | 164 | 145 | 32 | 68 | | | | |
| McCurtain | 220 | 156 | 357 | 130 | 178 | 137 | 34 | 66 | | | | |
| Coastal Plains | 288 | 133 | 640 | 110 | 320 | 291 | 8 | 92 | | | | |
| Bryan | 3,769 | 147 | 528 | 120 | 264 | 220 | 20 | 80 | | | | |
| Total | | | | | | | | | | | | |

Table 46.- Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Oklahoma, 1942

| Meal yield per acre (pounds) | Rolling Plains 1/ Cotton-Pea- | Central Prairies 2/ Cotton-Pea- | Gross Timbers 3/ Cotton-Pea- | Eastern ;Cotton-Pea- | Ozark- 5/: Pea-Cotton- :nuts: seed | Coastal- 6/: Pea-Cotton- :nuts: seed | State |
|------------------------------------|-------------------------------------|---------------------------------------|------------------------------------|-------------------------|--|--|-------|
| | | | | | Percent | Percent | |
| 0- 49 | 15 | 2 | 8 | 6 | 14 | 4 | 19 |
| 50- 99 | 18 | 7 | 27 | 14 | 29 | 11 | 32 |
| 100-149 | 19 | 10 | 39 | 13 | 32 | 13 | 18 |
| 150-199 | 18 | 8 | 15 | 13 | 16 | 14 | 13 |
| 200-249 | 12 | 10 | 6 | 11 | 6 | 13 | 7 |
| 250-299 | 11 | 9 | 3 | 9 | 2 | 10 | 6 |
| 300-349 | 4 | 10 | 1 | 7 | 1 | 9 | 3 |
| 350-399 | 2 | 7 | 1 | 7 | * | 7 | 0 |
| 400-449 | 1 | 10 | * | 5 | * | 4 | 1 |
| 450-499 | * | 6 | * | 4 | * | 4 | 1 |
| 500-549 | 0 | 4 | 3 | 0 | 3 | * | 2 |
| 550-599 | * | 5 | 2 | * | 2 | * | 2 |
| 600-649 | * | 3 | 2 | * | 2 | * | 2 |
| 650-699 | * | 2 | 1 | 1 | 1 | * | 1 |
| 700-749 | * | 1 | 1 | 1 | 1 | * | 1 |
| 750 and over | 6 | 2 | 2 | 2 | 2 | 1 | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 561 | 509 | 1,599 | 382 | 430 | 288 | 3,769 |

Sample counties: 1/ Caddo, Greer, Harmon, and Jackson. 2/ Grady and McClain. 3/ Carter, Hughes, Lincoln, Logan, Love, Okfuskee, Payne, and Seminole. 4/ Muskogee, Tulsa, and Wagoner.

5/ Latimer and McCurtain. 6/ Bryan.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 3, 1943

Table 47.- Comparative data on meal yields per acre from cottonseed and peanuts,
selected South Carolina Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | Ratio of meal yields:producing more meal | | Percent of farms producing more meal per acre, : per acre, : per acre from | | |
|----------------------|---------------------------|----------------------|---------|--|---------|--|---------|---------|
| | | Cotton | Peanuts | Cotton | Peanuts | | | |
| | | Number | Pounds | Pounds | Pounds | Percent | Percent | Percent |
| Piedmont Plateau | 93 | 310 | 443 | 253 | 190 | 75 | 59 | 41 |
| Anderson | 42 | 357 | 700 | 291 | 301 | 103 | 40 | 60 |
| Edgefield | 51 | 294 | 355 | 240 | 153 | 64 | 75 | 25 |
| Fall Line Sand Hills | 529 | 219 | 512 | 179 | 220 | 123 | 43 | 57 |
| Aiken | 233 | 215 | 517 | 175 | 222 | 127 | 35 | 65 |
| Chesterfield | 129 | 256 | 463 | 209 | 199 | 95 | 55 | 45 |
| Lexington | 167 | 232 | 503 | 189 | 216 | 114 | 46 | 54 |
| Middle Coastal Plain | 1,043 | 202 | 412 | 155 | 177 | 114 | 52 | 48 |
| Allendale | 267 | 173 | 300 | 133 | 129 | 97 | 52 | 48 |
| Barnwell | 291 | 158 | 417 | 121 | 179 | 148 | 33 | 67 |
| Clarendon | 120 | 305 | 357 | 235 | 154 | 66 | 80 | 20 |
| Horry | 69 | 331 | 910 | 255 | 391 | 153 | 32 | 68 |
| Lee | 130 | 206 | 310 | 158 | 133 | 84 | 60 | 40 |
| Marion | 166 | 448 | 650 | 344 | 280 | 81 | 70 | 30 |
| Lower Coastal Plain | 77 | 218 | 367 | 168 | 158 | 94 | 66 | 34 |
| Colleton | 46 | 209 | 383 | 161 | 165 | 102 | 61 | 39 |
| Dorchester | 31 | 253 | 310 | 195 | 133 | 68 | 74 | 26 |
| Total | 1,742 | 215 | 437 | 169 | 188 | 111 | 51 | 49 |

Table 48.- Frequency distribution of farms by meal yield per acre from cottonseed and peanuts, by areas in South Carolina, 1942

| Meal yield per acre (pounds) | Piedmont | Fall Line | Middle Coastal: | Lower Coastal: | State |
|---------------------------------|---|--|---|-----------------------------------|-----------------------------------|
| | Plateau 1/ Cotton-:Peanuts: seed : seed : | Sand Hills 2/ Cotton-:Peanuts: seed : seed : | Plain 3/ Cotton-:Peanuts: seed : seed : | Cotton-:Peanuts: seed : seed : | Cotton-:Peanuts: seed : seed : |
| 0- 49 | 0 | 2 | 2 | 8 | 16 |
| 50- 99 | 8 | 13 | 14 | 11 | 6 |
| 100-149 | 11 | 23 | 24 | 20 | 33 |
| 150-199 | 14 | 11 | 24 | 21 | 17 |
| 200-249 | 13 | 11 | 16 | 14 | 24 |
| 250-299 | 18 | 11 | 10 | 10 | 17 |
| 300-349 | 16 | 15 | 5 | 7 | 17 |
| 350-399 | 6 | 3 | 3 | 4 | 8 |
| 400-449 | 11 | 10 | 1 | 4 | 7 |
| 450-499 | 2 | 0 | 1 | 1 | 4 |
| 500-549 | 1 | 1 | 0 | 1 | 4 |
| 550-599 | | | 0 | 1 | 3 |
| 600-649 | | * | * | 0 | 1 |
| 650-699 | | * | * | 0 | 1 |
| 700-749 | | 0 | * | 1 | 1 |
| 750 and over | | * | 1 | 0 | * |
| Total | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 93 | 529 | 1,043 | 77 | 1,742 |

Sample counties: 1/ Anderson and Edgefield.

2/ Aiken, Chesterfield, and Lexington.

3/ Allendale, Barnwell, Clarendon, Horry, Lee, and Marion.

4/ Colleton and Dorchester.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
October 20, 1943

Table 49.- Comparative data on meal yields per acre from cottonseed and peanuts, selected Texas Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | Ratio of meal : yield per acre, more meal per acre from | |
|------------------|---------------------------|----------------------|---------|---|---|
| | | Cotton | Peanuts | Computed meal outturn from: peanuts to cottonseed | Percent of farms producing cottonseed ; Peanuts to cottonseed : |
| High Plains | 289 | 222 | 602 | 154 | 301 |
| Bailey | 51 | 202 | 680 | 140 | 340 |
| Dawson | 80 | 220 | 377 | 152 | 188 |
| Jamb | 53 | 188 | 860 | 130 | 430 |
| Lubbock | 105 | 260 | 827 | 180 | 414 |
| Rio Grande Plain | 309 | 63 | 479 | 44 | 240 |
| Atascosa | 158 | 63 | 527 | 44 | 264 |
| Daval | 108 | 63 | 250 | 44 | 125 |
| Starr | 43 | 63 | 157 | 44 | 78 |
| Edwards Plateau | 295 | 103 | 503 | 71 | 252 |
| Gillespie | 63 | 81 | 443 | 56 | 222 |
| San Saba | 232 | 116 | 540 | 80 | 270 |
| Rolling Plains | 829 | 161 | 454 | 112 | 227 |
| Gallahan | 203 | 121 | 397 | 84 | 198 |
| Cottle | 30 | 202 | 437 | 140 | 218 |
| Garza | 102 | 255 | 570 | 177 | 285 |
| Mitchell | 150 | 130 | 393 | 90 | 196 |
| Stonewall | 189 | 108 | 470 | 75 | 235 |
| Wichita | 53 | 291 | 490 | 202 | 245 |
| Wilbarger | 122 | 314 | 510 | 218 | 255 |
| Cross Timbers | 515 | 92 | 510 | 64 | 255 |
| Comanche | 293 | 90 | 510 | 62 | 255 |
| Jack | 68 | 108 | 427 | 75 | 214 |
| Wise | 154 | 99 | 527 | 69 | 264 |
| Grand Prairie | | | | | |
| Bosque | 84 | 102 | 350 | 78 | 175 |
| Coastal Plain | 1,875 | 125 | 423 | 89 | 212 |
| Anderson | 298 | 108 | 293 | 75 | 146 |
| Brazos | 75 | 148 | 250 | 103 | 125 |
| Franklin | 299 | 148 | 360 | 103 | 180 |
| Gonzales | 122 | 74 | 300 | 56 | 150 |
| Grayson | 299 | 159 | 747 | 121 | 374 |
| Harrison | 270 | 112 | 213 | 78 | 106 |
| Lamar | 175 | 139 | 407 | 106 | 204 |
| Montgomery | 39 | 76 | 227 | 53 | 114 |
| Nacogdoches | 298 | 116 | 297 | 80 | 148 |
| Total | 4,196 | 105 | 469 | 74 | 234 |
| | | | | | 316 |
| | | | | | 19 |
| | | | | | 81 |

Table 50.- Frequency distribution of farms by meal yields per acre from cottonseed and peanuts, by areas in Texas, 1942

| Meal yield per acre (pounds) | High Plains 1/ Cot- ton- nuts :seed: | Rio Grande: Plain 2/ Cot- Peanut- nuts :seed: | Edwards: Plateau 3/ Cot- Peanut- nuts :seed: | Rolling: Plains 4/ Cot- Peanut- nuts :seed: | Cross: Timbers 5/ Cot- Peanut- nuts :seed: | Grand Prairie 6/ Cot- Peanut- nuts :seed: | Coastal: Plains 7/ Cot- Peanut- nuts :seed: | State |
|------------------------------------|---|--|---|--|---|--|--|-------|
| 0- 49 | 11 | 4 | 69 | 12 | 32 | 3 | 17 | 7 |
| 50- 99 | 17 | 11 | 27 | 20 | 43 | 6 | 33 | 14 |
| 100-149 | 20 | 7 | 3 | 12 | 21 | 13 | 24 | 14 |
| 150-199 | 19 | 9 | 1 | 15 | 4 | 16 | 11 | 15 |
| 200-249 | 18 | 10 | * | 16 | 15 | 7 | 12 | * |
| 250-299 | 12 | 6 | 9 | 17 | 4 | 12 | 17 | 8 |
| 300-349 | 2 | 10 | 7 | 8 | 3 | 9 | 10 | 4 |
| 350-399 | 1 | 7 | 3 | 9 | 1 | 5 | 7 | 2 |
| 400-449 | 0 | 9 | 2 | 5 | * | 4 | 6 | * |
| 450-499 | * | 7 | 2 | 3 | * | 4 | 2 | * |
| 500-549 | 3 | 1 | 2 | 1 | 2 | 1 | 2 | 1 |
| 550-599 | 4 | * | 2 | 2 | 2 | 1 | 1 | 1 |
| 600-649 | 4 | * | 1 | * | 1 | * | 1 | 1 |
| 650-699 | 3 | 0 | * | 1 | * | 0 | * | 1 |
| 700-749 | 2 | 0 | 0 | 0 | * | * | * | * |
| 750 and over | 4 | 1 | * | * | * | * | * | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 289 | 309 | 295 | 829 | 515 | 84 | 1,875 | 4,196 |

Sample counties: 1/Bailey, Dawson, Lamb, and Lubbock. 2/Atascosa, Duval, and Starr. 3/Gillespie and San Saba. 4/Callahan, Cottle, Garza, Mitchell, Stonewall, Wichita, and Wilbarger. 5/Comanche, Jack and Wise. 6/Bosque. 7/Anderson, Brazos, Franklin, Gonzales, Grayson, Harrison, Lamar, Montgomery, and Nacogdoches.

* Less than 5 tenths of 1 percent.

Table 51.—Comparative data on oil yields per acre from cottonseed and soybeans,
State and area summary, Southern Region, 1942

| State and area | Number of farms | Yield per acre, 1942 | | Ratio of oil yields: producing more oil per acre, per acre from | | Percent oil yields: producing more oil per acre |
|----------------------------|-----------------------|----------------------|----------------|--|----------------|--|
| | | Cotton: | Soy- beans: | Computed oil outturn from Cotton: | Soy- beans: | |
| | | in lint | sample | seed | seed | |
| | | Number | Pounds | Bushels | Pounds | Percent |
| | | | | | | Percent |
| Arkansas | 2,135 | 518 | 17.0 | 167 | 130 | 78 |
| Louisiana | 815 | 386 | 11.5 | 116 | 95 | 82 |
| Mississippi | 1,037 | 447 | 15.8 | 148 | 133 | 90 |
| Texas 1/ | 70 | 291 | 8.7 | 65 | 75 | 115 |
| Total | 4,057 | 440 | 15.4 | 136 | 123 | 90 |
| Mississippi River Delta | 3,129 | 493 | 16.3 | 161 | 130 | 81 |
| Arkansas 2/ | 2,096 | 531 | 17.1 | 172 | 130 | 76 |
| Louisiana 3/ | 261 | 445 | 12.2 | 134 | 100 | 75 |
| Mississippi 4/ | 772 | 465 | 16.3 | 156 | 137 | 88 |
| Red River Delta | 298 | 326 | 12.7 | 97 | 100 | 103 |
| Arkansas 5/ | 39 | 230 | 13.4 | 67 | 102 | 152 |
| Louisiana 6/ | 259 | 352 | 11.8 | 106 | 97 | 92 |
| Other Louisiana areas 7/ | 295 | 290 | 11.2 | 87 | 92 | 106 |
| Other Mississippi areas 8/ | 265 | 324 | 10.1 | 97 | 85 | 88 |

Sample counties: 1/ Bailey, Lamb, Lubbock, and Wilbarger.

2/ Chicot, Clay, Craighead, Crittenden, Lee, and Mississippi.

3/ Concordia, Madison, and Morehouse.

4/ Coahoma, Holmes, Sharkey, and Sunflower.

5/ Little River and Miller.

6/ Caddo and Rapides.

7/ Saint Landry.

8/ Amite, Hinds, Itawamba, Montgomery, Pontotoc, and Yalobusha.

Table 52.—Relative advantage (or disadvantage) of soybeans over cottonseed in per acre oil production, by areas in Arkansas and Mississippi, 1942

| State and area | Percent of farms producing | | | Number of farms in sample | | |
|----------------|---|-------------|---------------------|---|-------------|---------------------|
| | Less than 50 percent | 100 percent | 200 percent or more | Less than 50 percent | 100 percent | 200 percent or more |
| | as much oil per acre from soybeans as from cottonseed | | | as much oil per acre from cottonseed as from soybeans | | |
| | | | | Percent | Percent | Number |
| Delta | | | | | | |
| Arkansas | 21 | 76 | 24 | 5 | 1 | 2,096 |
| Mississippi | 2/ | 62 | 38 | 11 | 5 | 773 |
| Other areas | | | | | | |
| Arkansas | 3 | 25 | 75 | 52 | 33 | 39 |
| Mississippi | 4/ | 74 | 26 | 14 | 6 | 265 |
| Total | | | | | | |
| Arkansas | 20 | 75 | 25 | 5 | 2 | 2,135 |
| Mississippi | 15 | 66 | 34 | 11 | 5 | 1,038 |
| Grand total | 18 | 71 | 29 | 8 | 3 | 3,173 |

Sample counties: 1/ Chicot, Clay, Crittenden, Lee, and Mississippi.

2/ Coahoma, Holmes, Sharkey, and Sunflower.

3/ Little River and Miller.

4/ Amite, Hinds, Itawamba, Montgomery, Pontotoc, and Yalobusha.

Table 53.- Frequency distribution of farms by ratio of soybean oil yield per acre to cottonseed oil yield per acre,
by areas in Arkansas and Mississippi, 1942

| Ratio Percent | Delta | | | Other areas | | | Grand total |
|----------------------------|------------------|-----|---------------------|-------------|------------------|---------------------|----------------|
| | Arkansas : 1/ | | Mississippi : 2/ | Total : | Arkansas : 3/ | Mississippi : 4/ | |
| | | | | Percent | | | - - - |
| 0- 9 | : | 0 | 0 | 0 | 0 | 4 | 3 |
| 10- 19 | : | 1 | * | 1 | 0 | 9 | 8 |
| 20- 29 | : | 4 | 1 | 5 | 0 | 9 | 8 |
| 30- 39 | : | 7 | 1 | 6 | 0 | 7 | 6 |
| 40- 49 | : | 9 | 5 | 8 | 3 | 10 | 8 |
| 0- 49 | : | 21 | 7 | 18 | 5 | 39 | 35 |
| 50- 59 | : | 10 | 10 | 10 | 3 | 8 | 7 |
| 60- 69 | : | 13 | 12 | 15 | 5 | 5 | 12 |
| 70- 79 | : | 13 | 11 | 15 | 4 | 8 | 12 |
| 80- 89 | : | 10 | 11 | 10 | 8 | 8 | 10 |
| 90- 99 | : | 9 | 11 | 9 | 4 | 6 | 9 |
| 50- 99 | : | 55 | 55 | 55 | 22 | 35 | 34 |
| 100-109 | : | 7 | 8 | 7 | 8 | 5 | 7 |
| 110-119 | : | 4 | 6 | 5 | 9 | 2 | 5 |
| 120-129 | : | 3 | 6 | 4 | 3 | 5 | 4 |
| 130-139 | : | 3 | 4 | 3 | 5 | 2 | 3 |
| 140-149 | : | 2 | 3 | 2 | 0 | 2 | 1 |
| 100-149 | : | 19 | 27 | 21 | 23 | 12 | 12 |
| 150-159 | : | 2 | 1 | 2 | 13 | 1 | 2 |
| 160-169 | : | 2 | 2 | 1 | 3 | 3 | 1 |
| 170-179 | : | * | 1 | * | 3 | 2 | 1 |
| 180-189 | : | * | 1 | 1 | 0 | 1 | 1 |
| 190-199 | : | * | 2 | * | 0 | 1 | * |
| 150-199 | : | 4 | 6 | 4 | 19 | 8 | 5 |
| 200 and over | : | 1 | 5 | 2 | 33 | 6 | 10 |
| Number of farms in sample: | 2,096 | 773 | 2,869 | 39 | 265 | 304 | 3,173 |

Sample counties:
 1/ Chicot, Clay, Craighead, Crittenden, Lee, and Mississippi.

2/ Coahoma, Holmes, Sharkey, and Sunflower.

3/ Little River and Miller.

4/ Amite, Hinds, Itawamba, Montgomery, Pontotoc, and Yalobusha.

* Less than 5 tenths of one percent.

Table 54.- Comparative data on oil yields per acre from cottonseed and soybeans,
selected Arkansas Counties, 1942

| County and area | Number of farms | Yield per acre, 1942 | | Computed oil outturn from sample : | Soybeans : Lint : beans : sample : | Ratio of : Percent of farms oil yield:producing more oil per acre, per acre from | |
|-------------------------|-----------------------|----------------------|------|--|--|--|----------------------------------|
| | | Cotton | Soy- | | | Cotton- to beans | Cotton- seed : cottonseed: |
| Red River Delta | 39 | 230 | 13•4 | 67 | 102 | 152 | 26 |
| Little River | 15 | 249 | 13•4 | 73 | 102 | 140 | 20 |
| Miller | 24 | 219 | 13•4 | 64 | 102 | 159 | 29 |
| Mississippi River Delta | 2,096 | 531 | 17•1 | 172 | 130 | 76 | 77 |
| Chicot | 68 | 396 | 12•2 | 128 | 93 | 73 | 72 |
| Clay | 176 | 411 | 13•8 | 133 | 105 | 79 | 61 |
| Craighead | 357 | 519 | 16•4 | 168 | 125 | 74 | 77 |
| Crittenden | 264 | 588 | 16•3 | 190 | 124 | 65 | 85 |
| Lee | 114 | 430 | 11•9 | 139 | 91 | 65 | 82 |
| Mississippi | 1,117 | 600 | 19•3 | 194 | 147 | 76 | 77 |
| Total | 2,135 | 518 | 17•0 | 167 | 130 | 78 | 76 |
| | | | | | | | 24 |

Table 55.- Frequency distribution of farms by oil yield per acre from cottonseed and soybeans
by sample counties, Arkansas Delta, 1942

| Oil yield per acre (pounds) | Chicot | | Clay | | 'Craighead | | 'Crittenden | | Lee | | Mississipi | | Delta | |
|--------------------------------|--------|------|------|------|------------|-------|-------------|------|------|------|------------|------|-------|------|
| | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- | Cot- |
| Under 60 | 5 | 16 | 5 | 16 | 1 | 9 | 10 | 1 | 22 | * | 3 | 1 | 7 | |
| 60- 79 | 7 | 32 | 6 | 21 | 1 | 14 | 14 | 3 | 24 | * | 8 | 1 | 12 | |
| 80- 99 | 21 | 15 | 13 | 15 | 3 | 13 | 5 | 6 | 17 | 2 | 10 | 4 | 11 | |
| 100-119 | 18 | 15 | 17 | 14 | 6 | 13 | 3 | 20 | 20 | 15 | 3 | 12 | 6 | 14 |
| 120-139 | 13 | 5 | 13 | 10 | 13 | 9 | 8 | 11 | 25 | 8 | 7 | 10 | 10 | |
| 140-159 | 13 | 13 | 19 | 8 | 17 | 22 | 13 | 25 | 18 | 8 | 13 | 27 | 14 | 23 |
| 160-179 | 9 | 3 | 12 | 6 | 23 | 6 | 19 | 3 | 17 | 2 | 16 | 7 | 17 | 6 |
| 180-199 | 7 | 7 | 5 | 16 | 7 | 15 | 6 | 3 | 2 | 16 | 9 | 14 | 7 | |
| 200-219 | 3 | 3 | 1 | 13 | 1 | 18 | 1 | 4 | 4 | 16 | 2 | 14 | 2 | |
| 220-239 | 1 | 2 | 1 | 4 | 2 | 12 | 2 | 1 | 2 | 12 | 7 | 9 | 4 | |
| 240-259 | 3 | 1 | 2 | 2 | 1 | 5 | 1 | 2 | 1 | 6 | 1 | 4 | 1 | |
| 260 and over | | | 1 | 1 | 1 | 3 | 7 | 2 | | 9 | 4 | 6 | 3 | |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 68 | 176 | 357 | 264 | 114 | 1,117 | 2,096 | | | | | | | |

* Less than 5 tenths of 1 percent.

Southern Division, AAA
September 15, 1943

Table 56.- Comparative data on oil yields per acre from cottonseed and soybeans,
selected Louisiana Parishes, 1942

| Parish and area | Number : | | | Yield per acre, 1942 | | | Ratio of oil yields:producing more oil per acre, per acre from | | |
|---------------------------------|----------|---------|---------|----------------------|--------------|--------------|--|---------|---------|
| | farms | Cotton: | Soy- | Computed oil | outturn from | Cotton- Soy- | Cotton: | Soy- | beans |
| sample | lint | beans | Cotton: | beans | Soy- | to | seed | seed | beans |
| | Number | Pounds | Bushels | Pounds | Pounds | Percent | Percent | Percent | Percent |
| Mississippi River Delta | 261 | 445 | 12.2 | 134 | 100 | 75 | 75 | 75 | 25 |
| Concordia | 96 | 480 | 9.6 | 144 | 79 | 55 | 87 | 87 | 13 |
| Madison | 66 | 436 | 16.8 | 131 | 138 | 105 | 47 | 47 | 53 |
| Morehouse | 99 | 436 | 10.8 | 131 | 89 | 68 | 82 | 82 | 18 |
| Red River Delta | 259 | 352 | 11.8 | 106 | 97 | 92 | 69 | 69 | 31 |
| Caddo | 32 | 340 | 17.0 | 102 | 140 | 137 | 44 | 44 | 56 |
| Rapides | 227 | 393 | 11.1 | 118 | 91 | 77 | 73 | 73 | 27 |
| Central Louisiana Mixed Farming | | | | | | | | | |
| St. Landry | 295 | 290 | 11.2 | 87 | 92 | 106 | 49 | 51 | |
| Total | 815 | 386 | 11.5 | 116 | 95 | 82 | 64 | 64 | 36 |

Table 57.- Frequency distribution of farms by oil yields per acre from cottonseed and soybeans, by areas in Louisiana, 1942

| Oil yield per acre (pounds) | Mississippi | River Delta | Delta 1/ 2/ | Red River | Central Louisiana: | |
|-----------------------------------|-------------|-------------|-------------------|-----------|-------------------------------|-------------------------------|
| | | | | | Cotton- seed ; Soybeans | Cotton- seed ; Soybeans |
| Under 60 | 2 | 15 | 5 | 21 | 21 | 32 |
| 60-79 | 5 | 20 | 9 | 16 | 24 | 10 |
| 80-99 | 15 | 32 | 17 | 39 | 27 | 13 |
| 100-119 | 14 | 9 | 24 | 4 | 16 | 30 |
| 120-139 | 23 | 11 | 21 | 5 | 5 | 20 |
| 140-159 | 16 | 4 | 14 | 2 | 5 | 15 |
| 160-179 | 14 | 6 | 8 | 7 | 2 | 33 |
| 180-199 | 9 | * | 1 | 2 | 0 | 12 |
| 200-219 | 3 | 3 | 1 | 2 | * | 8 |
| 220-239 | 1 | 0 | * | 1 | * | 5 |
| 240-259 | * | * | * | * | 1 | 1 |
| 260 and over | * | * | 1 | 1 | 1 | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 261 | 259 | 259 | 295 | 295 | 815 |

Sample parishes: 1/ Concordia, Madison, and Morehouse.

2/ Caddo and Rapides.

3/ St. Landry.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 10, 1943

Table 58.- Comparative data on oil yields per acre from cottonseed and soybeans,
selected Mississippi Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of : Percent of farms oil yield:producing more oil per acre,: per acre from | | |
|-----------------|---------------------------------------|----------------------|---------------|---|---|--|---------------|
| | | Cotton | Soy- beans | Computed oil outturn from Cotton-; Soy- seed : beans | soybeans : | Cotton- to : Cotton- seed : seed | Soy- beans |
| | | Pounds | Bushels | Pounds | Pounds | Percent | Percent |
| Delta | 772 | 465 | 16.3 | 156 | 137 | 88 | 62 |
| Coahoma | 298 | 423 | 16.4 | 142 | 138 | 97 | 52 |
| Holmes | 48 | 438 | 13.0 | 125 | 109 | 87 | 60 |
| Sharkey | 126 | 485 | 17.9 | 163 | 150 | 96 | 60 |
| Sunflower | 300 | 497 | 16.8 | 167 | 141 | 84 | 72 |
| Other | 265 | 324 | 10.1 | 97 | 85 | 88 | 70 |
| Amito | 6 | 270 | 6.8 | 77 | 57 | 74 | 50 |
| Hinds | 10 | 336 | 19.9 | 96 | 167 | 174 | 20 |
| Itawamba | 111 | 331 | 9.2 | 100 | 77 | 77 | 68 |
| Montgomery | 48 | 284 | 7.3 | 81 | 61 | 75 | 62 |
| Pontotoc | 20 | 350 | 5.7 | 106 | 48 | 45 | 85 |
| Yalobusha | 70 | 331 | 6.8 | 100 | 57 | 57 | 81 |
| Total | 1,037 | 447 | 15.8 | 148 | 133 | 90 | 64 |
| | | | | | | | 36 |

Table 59.- Frequency distribution of farms by oil yield per acre from cottonseed and soybeans, selected Mississippi Counties, 1942

| Oil yield per acre (pounds) | Coahoma | Holmes | Sharkey | Sunflower | Delta | Other | State total | Areas 1/ total | |
|--------------------------------|---------|--------|---------|-----------|-------|-------|----------------|----------------------------------|----------------------------------|
| | | | | | | | | Cot-:Soy- ton-:beans seed: | Cot-:Soy- ton-:beans seed: |
| Under 60 | 3 | 0 | 8 | 27 | 3 | 1 | 0 | 2 | 12 |
| 60-79 | 6 | 0 | 17 | 8 | 1 | 4 | 0 | 3 | 16 |
| 80-99 | 12 | 3 | 21 | 8 | 6 | 1 | 23 | 7 | 11 |
| 100-119 | 14 | 27 | 7 | 10 | 9 | 9 | 6 | 18 | 20 |
| 120-139 | 15 | 26 | 8 | 10 | 9 | 15 | 15 | 14 | 19 |
| 140-159 | 14 | 25 | 10 | 0 | 14 | 8 | 21 | 12 | 17 |
| 160-179 | 14 | 15 | 4 | 17 | 17 | 44 | 21 | 19 | 21 |
| 180-199 | 10 | 1 | 8 | 4 | 21 | 7 | 21 | 1 | 16 |
| 200-219 | 7 | 1 | 13 | 10 | 11 | 3 | 9 | 9 | 5 |
| 220-239 | 3 | 0 | 4 | 0 | 6 | 0 | 3 | * | 0 |
| 240-259 | 1 | 1 | 1 | 3 | 1 | 2 | 1 | 2 | * |
| 260 and over | 1 | 1 | 3 | 2 | 1 | 1 | * | 1 | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 298 | 48 | 126 | 300 | 772 | 265 | 1,037 | | |

1/ Includes Amite, Hinds, Itawamba, Montgomery, Pontotoc, and Yalobusha Counties.

* Less than 5 tenths of 1 percent.

Table 60.-- Comparative data on meal yields per acre from cottonseed and soybeans,
State and area summary, Southern Region, 1942

| State and area | Number of farms | Yield per acre, 1942 | | Computed meal outturn from sample | Cotton: Soy- bean | Soy- bean | Ratio of meal yields: producing more meal per acre, per acre from soybeans to cottonseed: | | Percent of farms producing more meal per acre from soybeans |
|----------------------------|---|----------------------|---------|---|----------------------|--------------|---|---------|---|
| | | Pounds | Bushels | | | | Pounds | Percent | |
| Arkansas | 2,134 | 518 | 17.0 | 481 | | 805 | 167 | 24 | 76 |
| Louisiana | 815 | 386 | 11.5 | 319 | | 551 | 173 | 22 | 78 |
| Mississippi | 1,038 | 447 | 15.8 | 380 | | 755 | 199 | 14 | 86 |
| Texas 1/ | 70 | 291 | 8.7 | 202 | | 434 | 215 | 17 | 83 |
| Total | 4,057 | 440 | 15.4 | 389 | | 733 | 188 | 21 | 79 |
| Mississippi River Delta | 3,129 | 493 | 16.3 | 448 | | 774 | 173 | 20 | 80 |
| Arkansas 2/ | 2,095 | 531 | 17.1 | 494 | | 810 | 164 | 24 | 76 |
| Louisiana 3/ | 261 | 445 | 12.2 | 368 | | 585 | 159 | 27 | 73 |
| Mississippi 4/ | 773 | 465 | 16.3 | 407 | | 778 | 191 | 5 | 95 |
| Red River Delta | 298 | 326 | 12.7 | 270 | | 608 | 225 | 8 | 92 |
| Arkansas 5/ | 39 | 230 | 13.4 | 194 | | 634 | 327 | 3 | 97 |
| Louisiana 6/ | 259 | 352 | 11.8 | 291 | | 566 | 194 | 22 | 78 |
| Other Louisiana areas 7/ | 295 | 290 | 11.2 | 240 | | 537 | 224 | 18 | 82 |
| Other Mississippi areas 8/ | 265 | 324 | 10.1 | 252 | | 482 | 191 | 37 | 63 |
| Sample counties: | 1/ Bailey, Lamb, Lubbock, and Willbarger. 2/ Chicot, Clay, Craighead, Crittenden, Lee, and Mississippi. 3/ Concordia, Madison, and Morehouse. 4/ Coahoma, Holmes, Sharkey, and Sunflower. 5/ Little River and Miller. 6/ Caddo and Rapides. 7/ St. Landry. 8/ Amite, Hinds, Itawamba, Montgomery, Pontotoc, and Yalobusha. | | | | | | | | |

Table 61.- Comparative data on meal yields per acre from cottonseed and soybeans,
selected Arkansas Counties, 1942

| County and area | Number of farms in sample | Yield per acre, 1942 | | | Ratio of meal yield:producing more meal per acre, per acre from soybeans : to cottonseed: | | | Percent of farms producing more meal per acre from soybeans : cottonseed : |
|-------------------------|---------------------------|----------------------|-----------|--|---|---------------|------------------|--|
| | | Cotton | Soy-beans | Computed meal outturn from cotton : soybeans | Cotton : soybeans | Cotton : seed | Soy-beans : seed | |
| | Number | Pounds | Bushels | Pounds | Pounds | Bushels | Pounds | Percent |
| Red River Delta | 39 | 230 | 13.4 | 194 | 634 | 327 | 3 | 97 |
| Little River | 15 | 249 | 13.4 | 210 | 634 | 302 | 0 | 100 |
| Miller | 24 | 219 | 13.4 | 184 | 634 | 345 | 4 | 96 |
| Mississippi River Delta | 2,096 | 531 | 17.1 | 494 | 810 | 164 | 24 | 76 |
| Chicot | 68 | 396 | 12.2 | 368 | 578 | 157 | 18 | 82 |
| Clay | 176 | 411 | 13.8 | 382 | 653 | 171 | 18 | 82 |
| Craighead | 357 | 519 | 16.4 | 483 | 776 | 161 | 19 | 81 |
| Crittenden | 264 | 588 | 16.3 | 547 | 772 | 141 | 78 | 22 |
| Lee | 114 | 430 | 11.9 | 400 | 563 | 141 | 27 | 73 |
| Mississippi | 1,117 | 600 | 19.3 | 558 | 914 | 164 | 14 | 86 |
| Total | 2,135 | 518 | 17.0 | 480 | 805 | 168 | 24 | 76 |

Table 62.- Frequency distribution of farms by meal yield per acre from cottonseed and soybeans, by sample counties, Arkansas Delta, 1942.

| Meal yield per acre (pounds) | Chicot | Clay | Craighead | Crittenden | Lee | Mississippi | Delta total |
|---------------------------------|--------|------|-----------|------------|-----|-------------|----------------|
| | * | * | * | * | * | * | * |
| 0- 99 | 1 | 1 | 1 | 2 | 1 | 1 | 1 |
| 100-199 | 3 | 7 | 1 | 5 | 3 | 3 | 4 |
| 200-299 | 32 | 24 | 6 | 1 | 14 | 6 | 6 |
| 300-399 | 25 | 22 | 12 | 5 | 11 | 9 | 14 |
| 400-499 | 21 | 28 | 17 | 12 | 7 | 37 | 5 |
| 500-599 | 12 | 10 | 13 | 31 | 11 | 21 | 27 |
| 600-699 | 9 | 5 | 8 | 11 | 6 | 2 | 10 |
| 700-799 | 3 | 10 | 2 | 15 | 2 | 12 | 17 |
| 800-899 | 6 | 6 | 8 | 1 | 4 | 7 | 10 |
| 900-999 | 14 | 6 | 6 | * | 22 | 9 | 14 |
| 1000-1099 | | | | | 10 | 8 | 8 |
| 1100-1199 | | | | | 30 | 29 | 26 |
| 1200-1299 | | | | | 5 | 21 | 16 |
| 1300-1399 | | | | | 22 | 4 | 5 |
| 1400-1499 | | | | | 4 | 13 | 6 |
| 1500-1599 | | | | | 10 | 7 | 14 |
| 1600-1699 | | | | | 4 | 4 | 8 |
| 1700-1799 | | | | | 1 | 1 | 2 |
| 1800 and over | | | | | 1 | 1 | 2 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 63 | — | 176 | — | 357 | — | 1,117 |
| | | | | | 264 | — | 2,096 |

* Less than 5 tenths of 1 percent.

Southern Division, AAA
September 17, 1943

Table 63.- Comparative data on meal yields per acre from cottonseed and soybeans,
selected Louisiana Parishes, 1942

| Parish and area | Number of farms in sample | Yield per acre, 1942 | | Ratio of meal yields:producing more meal per acre, per acre from | | Percent of farms producing more meal per acre |
|------------------------------------|---------------------------------------|----------------------|--------------|--|--------------|--|
| | | Cotton | Soy- bean | outturn from Cotton-beans | Soy- bean | |
| | Number | Pounds | Bushels | Pounds | Percent | Percent |
| Mississippi River Delta | 261 | 445 | 12.2 | 368 | 585 | 159 |
| Concordia | 96 | 480 | 9.6 | 397 | 460 | 116 |
| Madison | 66 | 436 | 16.8 | 360 | 805 | 224 |
| Morehouse | 99 | 436 | 10.8 | 360 | 518 | 144 |
| Red River Delta | 259 | 352 | 11.8 | 291 | 566 | 194 |
| Caddo | 32 | 340 | 17.0 | 281 | 815 | 290 |
| Rapides | 227 | 393 | 11.1 | 325 | 532 | 164 |
| Central Louisiana Mixed Farming | | | | | | |
| St. Landry | 295 | 290 | 11.2 | 240 | 537 | 224 |
| Total | 815 | 386 | 11.5 | 319 | 551 | 173 |
| | | | | | | 22 |
| | | | | | | 78 |

Table 64.- Frequency distribution of farms by meal yields per acre from cottonseed and soybeans, by areas in Louisiana, 1942

| Meal yield per acre (pounds) | Mississippi | | | Red River | | | Central Louisiana: | | |
|------------------------------------|-----------------------|----------------|-----------------------|----------------|-----------------------|----------------|--------------------|----------|----------|
| | River Delta | | | Acacia | | | Fixed Farming: | | |
| | 1/ Cotton- seed | 2/ Soybeans | 3/ Cotton- seed | 1/ Soybeans | 2/ Cotton- seed | 3/ Soybeans | Cotton- seed | Soybeans | Soybeans |
| 0- 99 | 0 | 1 | 2 | 2 | 5 | * | 2 | 2 | 1 |
| 100- 199 | 5 | 2 | 7 | 7 | 55 | 1 | 15 | 3 | 3 |
| 200- 299 | 25 | 8 | 31 | 7 | 42 | 3 | 33 | 6 | 6 |
| 300- 399 | 34 | 14 | 39 | 15 | 15 | 5 | 29 | 11 | 11 |
| 400- 499 | 27 | 30 | 17 | 33 | 5 | 6 | 16 | 21 | 21 |
| 500- 599 | 12 | 15 | 2 | 12 | * | 8 | 5 | 12 | 12 |
| 600- 699 | 1 | 6 | * | 4 | * | 8 | * | 6 | 6 |
| 700- 799 | | | 11 | 5 | | 9 | | 8 | 8 |
| 800- 899 | | | 4 | 2 | | 2 | | 3 | 3 |
| 900- 999 | | | 6 | 7 | | 21 | | 12 | 12 |
| 1000-1099 | | | * | 1 | | | | 1 | 1 |
| 1100-1199 | | | 2 | 2 | | | 8 | 4 | 4 |
| 1200-1299 | | | 1 | 1 | | | 1 | 1 | 1 |
| 1300-1399 | | | 0 | 1 | | | 2 | 1 | 1 |
| 1400-1499 | | | * | * | | | 5 | 2 | 2 |
| 1500 and over | | | | 1 | | | 20 | 8 | 8 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample | 261 | 259 | 259 | 259 | 259 | 259 | 259 | 259 | 259 |

Sample parishes: 1/ Concordia, Madison, and Morehouse.

2/ Caddo and Rapides.

3/ St. Landry.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 10, 1943

Table 65.- Comparative data on meal yields per acre from cottonseed and soybeans,
selected Mississippi Counties, 1942

| County and area | Number of farms | Yield per acre, 1942 | | Ratio of meal yields:producing more meal per acre, : per acre from | | Percent of farms producing more meal | |
|-----------------|-----------------------|----------------------|----------------------------|--|--------------------------------|---|---------------|
| | | Cotton: | Soy- in lint : beans | Cotton-: Soy- seed : beans | Cotton-: to seed : beans | Cotton-: seed : beans | Soy- beans |
| | | Number | Pounds | Bushels | Pounds | Pounds | Percent |
| Delta | 773 | 465 | 16.3 | 407 | 778 | 191 | 5 |
| Coahoma | 298 | 423 | 16.4 | 371 | 783 | 211 | 1 |
| Holmes | 48 | 438 | 13.0 | 384 | 621 | 162 | 19 |
| Sharkey | 126 | 485 | 17.9 | 425 | 855 | 201 | 5 |
| Sunflower | 301 | 497 | 16.3 | 435 | 802 | 184 | 8 |
| Other Areas | | | | | | | |
| Amite | 6 | 270 | 6.8 | 201 | 325 | 162 | 33 |
| Hinds | 10 | 336 | 19.9 | 250 | 950 | 380 | 10 |
| Itawamba | 111 | 331 | 9.2 | 261 | 439 | 163 | 26 |
| Montgomery | | 48 | 284 | 7.3 | 211 | 349 | 165 |
| Pontotoc | | 20 | 350 | 5.7 | 276 | 272 | 99 |
| Yalobusha | | 70 | 331 | 6.8 | 261 | 325 | 125 |
| Total | 1,038 | 447 | 15.8 | 380 | 755 | 199 | 14 |
| | | | | | | | 86 |

Table 66.—Frequency distribution of farms by meal yields per acre from cottonseed and soybeans,
selected Mississippi Counties, 1942

| Meal yield per acre (pounds) | Coahoma | | Holmes | | Sharkey | | Sunflower | | Delta | | Other | | State total | |
|---------------------------------|---------|------|--------|------|---------|------|-----------|------|-------|------|-------|------|----------------|------|
| | Cot- | Soy- | Cot- | Soy- | Cot- | Soy- | Cot- | Soy- | Cot- | Soy- | Cot- | Soy- | Cot- | Soy- |
| | beans | seed | beans | seed | beans | seed | beans | seed | beans | seed | beans | seed | beans | seed |
| 0- 99 | 1 | 0 | 2 | 2 | 0 | 0 | 1 | 0 | 1 | * | 2 | 13 | 1 | 3 |
| 100- 199 | 7 | 0 | 19 | 13 | 4 | 1 | 0 | 0 | 5 | 1 | 22 | 18 | 9 | 5 |
| 200- 299 | 23 | 0 | 29 | 8 | 13 | 1 | 5 | 0 | 15 | 1 | 51 | 16 | 24 | 5 |
| 300- 399 | 28 | 0 | 19 | 11 | 21 | 2 | 29 | 0 | 27 | 1 | 20 | 14 | 25 | 4 |
| 400- 499 | 26 | 1 | 10 | 11 | 35 | 6 | 45 | 19 | 33 | 10 | 5 | 17 | 27 | 12 |
| 500- 599 | 11 | 5 | 19 | 6 | 23 | 9 | 17 | 9 | 16 | 7 | 0 | 6 | 12 | 7 |
| 600- 699 | 3 | 23 | 2 | 2 | 2 | 2 | 1 | 11 | 2 | 14 | * | 3 | 2 | 11 |
| 700- 799 | 1 | 27 | 10 | 2 | 16 | 1 | 15 | 1 | 19 | * | 5 | 5 | * | 16 |
| 800- 899 | 25 | 0 | 7 | 0 | 13 | 0 | 16 | * | 0 | 16 | * | 0 | 12 | * |
| 900- 999 | 12 | 19 | 41 | 1 | 17 | * | 19 | * | 19 | 3 | * | 3 | * | 15 |
| 1000-1099 | 4 | 4 | 7 | 7 | 2 | 4 | 7 | 2 | 4 | * | 3 | 5 | 5 | * |
| 1100-1199 | 1 | 10 | 4 | 9 | 1 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | * | * |
| 1200-1299 | 0 | 0 | 0 | 2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | * |
| 1300-1399 | 0 | 0 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | * |
| 1400-1499 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | * |
| 1500 and over | 1 | 2 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | 1 | 1 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| Number of farms in sample: | 298 | 48 | 126 | 301 | 773 | 265 | 298 | 48 | 126 | 301 | 773 | 265 | 298 | 48 |

1/ Includes Amite, Hinds, Itawamba, Montgomery, Pontotoc, and Yalobusha Counties.

* Less than 5 tenths of 1 percent.

Southern Division, AAA
November 9, 1943

