



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.



ADAPTATION RESOURCES WORKBOOK FOR CALIFORNIA SPECIALTY CROPS

A Guide for Adaptation Planning



AUTHORS

Lauren E. Parker, Devon Johnson, Tapan B. Pathak, Michael Wolff, Virginia Jameson, and Steven M. Ostoja

Napa Valley wine country





Authors

Lauren E. Parker^{1,2}, Devon Johnson^{1,2}, Tapan B. Pathak^{3,4}, Michael Wolff⁵, Virginia Jameson⁵, and Steven M. Ostoja^{1,2,6}

¹ USDA California Climate Hub, Davis, California

² Institute of the Environment, University of California Davis, Davis, California

³ University of California Agriculture and Natural Resources, Merced, California

⁴ University of California Merced, Merced, California

⁵ California Department of Food and Agriculture, Sacramento, California

⁶ USDA-ARS Sustainable Agricultural Water Systems Research Unit, Davis, California

Lauren Parker is the research program manager at the USDA California Climate Hub and a postdoctoral fellow at the U.C. Davis Institute of the Environment, Davis, California, leparkr@ucdavis.edu.

Devon Johnson was the coordinator of the USDA California Climate Hub and an assistant specialist in climate change adaptation in agriculture at the U.C. Davis Institute of the Environment, Davis, California. Devon is now a senior policy analyst at the Center for Resource Solutions, San Francisco, California.

Tapan Pathak is an extension specialist in climate adaptation in agriculture at U.C. Agriculture and Natural Resources and is affiliated with the Department of Civil and Environmental Engineering at U.C. Merced, Merced, California, tbp@ucanr.edu.

Michael Wolff is a senior environmental scientist at the California Department of Food and Agriculture, Sacramento, California, michael.wolff@cdfa.ca.gov.

Virginia Jameson is the deputy secretary for climate and working lands at the California Department of Food and Agriculture, Sacramento, California, virginia.jameson@cdfa.ca.gov.

Steven Ostoja is the director of the USDA California Climate Hub, a scientist at the USDA-ARS Sustainable Agricultural Water Systems research unit, and a fellow at the U.C. Davis Institute of the Environment, Davis, California, steven.ostoja@usda.gov.

Preferred citation:

Parker, L. E., Johnson, D., Pathak, T. B., Wolff, M., Jameson, V., and Ostoja, S. M. (2023). Adaptation Resources Workbook for California Specialty Crops. USDA California Climate Hub Technical Report CACH-2023-1. Davis, CA: U.S. Department of Agriculture, Climate Hubs. 55 p.

Acknowledgments

Many individuals contributed their knowledge and expertise to the development of this workbook. The authors would especially like to thank the following agriculture professionals and members of the technical service provider community who served as expert reviewers, providing feedback and edits that improved the quality of this workbook.

Mark Battany, PhD, UC Cooperative Extension

Liz Colby, P.E., USDA-NRCS

Darcy Cook, Mission Resource Conservation District

Ben Faber, PhD, UC Cooperative Extension

Norm Groot, Monterey County Farm Bureau

Amy Lofting, Meridian Farm

Karen Lowell, PhD, CCA, USDA-NRCS

Wendy Rash, USDA-NRCS

Margaret Smither-Kopperl, USDA-NRCS

Cover photos only courtesy of USDA-NRCS.

All remaining photos courtesy istockphoto.com.

The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the United States Department of Agriculture or the Agricultural Research Service of any product or service to the exclusion of others that may be suitable.

Table of Contents

A GUIDE FOR ADAPTATION PLANNING	6
Climate change and California agriculture.....	6
What is this workbook and who is it for?	6
What does "climate adaptation" mean?.....	6
How do I use this workbook?	7
THE 5-STEP ADAPTATION PROCESS.....	7
Step 1: Define Goals and Objectives	8
Step 2: Assess Climate Impacts and Vulnerabilities	10
Step 3: Re-Evaluate Goals and Objectives Given Projected Impacts and Vulnerabilities	12
Step 4: Identify Adaptation Strategies, Approaches, and Practices	14
Step 5: Monitor and Evaluate	17
A MENU OF ADAPTATION STRATEGIES AND APPROACHES FOR CALIFORNIA SPECIALTY CROPS	20
Strategy 1: Support and Maintain Soil Health, Soil Biological Services, and Water Quality	20
Strategy 2: Cope with Uncertain Water Availability	23
Strategy 3: Manage Biological Crop Stressors.....	26
Strategy 4: Prepare for Temperature Change	28
Strategy 5: Enhance Preparedness for Extreme Events	30
Strategy 6: Manage Farms and Fields as Part of a Larger Landscape	32
Strategy 7: Develop or Expand Co-Benefit Efforts to Mitigate Climate Change	33
RESOURCES	36
REFERENCES	47

A GUIDE FOR ADAPTATION PLANNING

Climate change and California agriculture

[California agriculture](#) is not immune to the risks of climate change^{1,2} and [specialty crops may be uniquely vulnerable](#) to the expected climatic changes. California's climate has changed since record-keeping began in the late 1800s. Warmer temperatures² and earlier snowmelt³ have been observed and in a warming world, we can expect changes in the frequency, duration, and intensity of droughts, floods, and heatwaves.^{2,4,5,6,7,8} Warming winter temperatures will reduce the accumulation of winter chill that is needed for fruit and nut crops,⁹ warmer springs may advance bloom in crops like almonds,¹⁰ and warmer summers may mean earlier harvest for tomatoes.¹¹ Warmer temperatures can also influence the timing and intensity of pest pressure,¹² while extreme heat events place additional stress on crops and the water resources that farmers rely on to manage the heat.⁷ All of these changes may have profound impacts on California's specialty crop production. In fact, some estimates suggest that climate change could reduce yields of specialty crops such as avocados, walnuts, and table grapes by as much as 40% by the end of the century.¹³ In light of the downward pressure climate change may place on California agriculture, adaptation will be critical for maintaining the state's specialty crop production in the decades to come.

What is this workbook and who is it for?

This workbook provides resources and guidance for California's agricultural producers and technical service providers to help them identify actions for increasing resilience to a changing climate. The workbook was developed with specialty crop production in mind, but it contains useful climate adaptation information for many agricultural production systems.

What does "climate adaptation" mean?

[Climate adaptation](#) for agriculture encompasses actions that help farmers and the agroecosystems they manage cope with the effects of climate change. Actions that help to reduce the damages from climate stressors like heatwaves, droughts, or floods can be considered climate adaptive. Actions that help farms more readily recover from these events are also climate adaptive. Different types of adaptation frameworks exist depending on land managers' short- and long-term adaptation goals. For example, some natural resource land managers use the [Resist-Accept-Direct \(RAD\) Decision Framework](#)¹⁴ to guide their management options for landscapes at risk of, or undergoing, ecological change. This framework helps managers decide whether to resist ecosystem changes and maintain historical conditions, or whether the most appropriate course of action is accepting the change as it occurs or actively directing the change toward preferred new conditions. The [Resistance-Resilience-Transformation \(RRT\)](#) framework¹⁵ (Figure 1) provides another way to conceptualize potential adaptation actions. Similar to RAD, the RRT framework offers three different approaches to adaptation but RRT can provide a spectrum of paths towards adaptation, ranging from *active resistance* to *accelerated transformation*. Adaptation frameworks like RAD and RRT can help growers develop strategies for responding to climate change on their farms.

Climate adaptation actions tend to focus on withstanding current and/or projected future climate conditions, but many adaptation actions also have benefits for [climate mitigation](#). These mitigation actions help to combat the effects of climate change by reducing greenhouse gas emissions or by sequestering (storing) carbon-based greenhouse gases from the atmosphere in plant matter and/or soils. In this workbook, you will mainly find climate adaptation actions, but we also include one section that is specifically devoted to climate mitigation, and we note when a given action provides both adaptation and mitigation benefits, sometimes referred to as co-benefits.



Figure 1. The Resistance–Resilience–Transformation typology provides one framework for understanding the range of climate adaptation actions available to producers. Different actions require or precipitate varying degrees of change from the status quo, but producers can choose actions from across the spectrum to meet their farm's unique needs and fit within their overall management plan.

How do I use this workbook?

This adaptation workbook draws on a [5-step process](#)¹⁶ (Figure 2) developed by the Northern Institute for Applied Climate Science (NIACS) for assessing the risks and vulnerabilities of land management goals to climate change, and it provides a path for developing a plan to address those challenges in California's specialty crop production systems. For each step, the workbook provides a brief description of that step's purpose and contextual content, followed by a corresponding worksheet that uses the concepts within the step and a series of guided questions to build a climate adaptation management plan. Collectively, these steps and worksheets guide users through the adaptation process, from defining one's goals and objectives through monitoring and evaluating the adaptive actions taken. We encourage users to read each step sequentially as the steps build on one another. The worksheets may be completed after each associated step or they may be completed together after reviewing all of the workbook steps. We suggest that you consider one goal or project at a time and complete each worksheet with that specific goal or project in mind.

Throughout the workbook, you will find references to resources that explore the relevant issues in greater detail, like Cooperative Extension guides, NRCS technical documents, and scientific literature. At the end of the workbook is a [resources section](#) with links to references that can further aid in adaptation planning. You can use this workbook and its 5-step framework to help guide you in developing a climate adaptation management plan that is specific for your farm's goals and needs.




Figure 2. The five steps of this adaptation workbook offer a guide for improving the adaptive capacity of your farm by providing a structured approach for incorporating climate change considerations into farm management planning. More detail on the purpose of each step is provided in the associated workbook section. (Figure adapted from USDA reports ^{16,17})

STEP 1 DEFINE GOALS AND OBJECTIVES

The idea of adapting a farm to cope with the effects of climate change can seem complicated and feel overwhelming. But by breaking the process down into a set of sequential steps the process will feel less onerous. The purpose of Step 1 is to **define your management goals and objectives**, providing a foundational context for developing the plan that will help you achieve your specific interests. These management goals and objectives answer questions like "What is the long-term vision for your farm?" and "How will you achieve your vision?" These are likely things you already know and understand well, but perhaps they haven't been expressed or considered explicitly. Creating a management plan with clearly defined long-term goals supported by short-term objectives provides a solid foundation for effective climate adaptation.

Goals, Objectives, and Baselines

In developing a farm management plan, it may be useful to create goals and outline objectives for the different categories of farm management. Long-term **goals** frame a farm's mission and vision, whereas short-term **objectives** may be quantifiable and tangible targets that producers can work towards, providing a roadmap for executing and accomplishing pre-determined goals. Once the management goals and objectives have been identified, you may wish to establish a **baseline** or measure of the current conditions for each one to serve as a starting point against which changes can be compared and implemented, and adaptation actions can be monitored and evaluated (please see Step 5 for more on monitoring and evaluation). Box 1 below shows one example of how you can use a long-term goal and subsequent baseline measurements to better determine and refine your short-term objectives.

Box 1. Linking goals, objectives, and baselines can provide organization in the farm management plan.

Farm Management Category: Plant Health
Long-Term Management Goal: Decrease occurrence or impact of salinization
Baseline: Total Dissolved Solids (TDS) in soil based on soil salinity test
Short-Term Management Objectives[*]:
1. Reduce salt in soils via leaching 2. Improve drainage to facilitate salt leaching

^{*}UCANR Publication 8550 Managing Salts by Leaching



Saline soils can negatively affect crop productivity. Adaptation actions such as leaching can help reduce the impacts of saline soils.

STEP 1 PUT IT INTO PRACTICE

Consider each of the guiding questions in the worksheets below. Remember, the collection of worksheets in this workbook provides a step-by-step pathway for developing your adaptation plan.

STEP 1 asks you to consider a specific management goal for your farm and identify the objectives for meeting that goal. Remember, **management goals** are broad, high-level outcomes, whereas **management objectives** are the approaches or actions that can help you meet your goals. Some examples of management goals and objectives can be found in Box 1 above.

Recall Box 1 on page 6. Now try filling out your own box below.

Project Name:

Farm Location:

Farm Management Category: *What would you like to focus on? Plant health, water storage, energy use, etc.?*

Management Goal: *What outcome(s) would you like to see for this category in the long-term?*


Baseline: *What measurable indicators can you use to assess your progress as you work towards your goal? It is usually very useful to take "baseline" measurements before starting new practices so the changes can be seen more clearly (see Step 5 worksheet).*

Management Objectives: *What are some actions you can take that will help you to meet your goal? Remember, your objectives should be quantifiable in the short-term and provide a roadmap towards meeting your long-term goal.*

STEP 2 ASSESS CLIMATE IMPACTS AND VULNERABILITIES

Global Changes, Local Impacts

Climate change is a global phenomenon, but the way it "looks" can vary widely from place to place. **Understanding the effects of climate variability and change** for a given location and production system is crucial for creating an adaptation plan that will provide operational resilience and improve adaptive capacity in the near- and long-term. Resources like [CalAgroClimate](#), [Cal-Adapt](#), [Climate Toolbox](#), and the [U.S. Climate Resilience Toolkit](#) provide location-specific current and future climate information that can help producers assess the potential impact of climate change for their farm. For example, Cal-Adapt hosts various tools (Figure 3) that show how climate indicators like temperature, precipitation, and extreme heat have changed in the past and how they are projected to change in the future under climate change. By exploring these tools, growers can see what climate changes to expect and then use that information to assess the vulnerability of their farm or crop.



The screenshot shows the Cal-Adapt website interface. At the top, a banner features a green and blue textured background with the text "Climate Tools" and "New to climate data? Start with the Local Climate Change Snapshot Tool". Below the banner is a navigation bar with buttons for "All", "Temperature", "Precipitation", "Snowpack", "Sea Level Rise", and "Wildfire". The main content area is divided into four sections: "Local Climate Change Snapshot" (with icons for sun, water drop, and leaf), "Annual Averages" (with icons for sun and water drop), "Sea Level Rise – Coastal Inundation Scenarios" (with an icon of water waves), and "Extreme Weather" (with an icon of a sun with a lightning bolt). Each section includes a brief description and an "EXPLORE" button.

Figure 3. Cal-Adapt has many useful tools for exploring California's climate at the local, county, or watershed scales. The website also hosts helpful how-to guides and informative documentation for first-time users and experts alike.


Determining Vulnerability

Determining how vulnerable a farm or crop system is to climate change is an important step in developing an adaptation plan. A farm or crop's **vulnerability** is defined by the degree to which it can cope with the adverse effects of climate change, including climate variability and extreme events.¹⁸ In addition to a location's climate conditions and expected climate changes, factors like water security, crop tolerance to unfavorable conditions, and site characteristics like topography and soils – not to mention social and economic factors – all influence how susceptible a region or farm is to climate-caused damages. A farm or crop system's **adaptive capacity**, that is, its ability to adjust to changes, take advantage of opportunities, and cope with disturbance, can also impact vulnerability.¹⁸

Understanding the specific vulnerabilities of your farm or ranch will help you to pinpoint the parts of your operation that are most at risk due to climatic changes, and where you should focus your attention as you create your adaptation plan. Region- or cropping system-specific **vulnerability assessments** can provide additional context that may be helpful in assessing a farm's vulnerability to current and future climate conditions. The State of California's [Climate Change Assessment](#) website and the [ResilientCA](#) website provide multiple resources for assessing climate impacts and vulnerability. The California Energy Commission's report on its [vulnerability index for California agriculture](#) (shown on the next page in Figure 4) offers another resource for assessing farm-specific climate vulnerabilities. While these tools and resources provide good general information, determining an individual farm's vulnerability may also require working with experts such as RCD, NRCS, or cooperative extension staff, farm advisors, or other trusted technical assistance providers.

Figure 4. The vulnerability index for California agriculture accounts for many factors, including climate exposure, crop sensitivity, land characteristics, and socioeconomic vulnerabilities. Much of the state's specialty crop production area falls into the moderately high to very high vulnerability categories. That being said, this map does not take into account any climate smart management practices growers may have implemented to better prepare their land for climatic changes.

This figure is available in the 2012 California Energy Commission Report *Vulnerability and Adaptation to Climate Change in California Agriculture*¹⁹ and is reprinted here under Creative Commons License Attribution 4.0.



STEP 2 PUT IT INTO PRACTICE

STEP 2 has you explore different dimensions of climate change and what they might mean for your farm. Climate change is globally dynamic, but the associated impacts and potential vulnerabilities are going to depend on your location and the crops you are farming, among other factors. To [assess climate impacts and vulnerabilities for your farm](#), check out the resources mentioned above and consider the guiding questions in the box below.

Key Climate Concerns: What are the three most important climate factors that have the greatest potential to influence your farming operation? Consider things like growing season length, seasonality, duration and intensity of heatwaves, drought, extreme precipitation and flooding, chill accumulation, frost damage, wildfire and smoke, damage from insects or diseases, etc.

Climate Projections: How will climate change influence these key concerns? For example, will climate change increase or decrease the incidence of extreme precipitation and flooding, or the frequency of heatwaves at your location? Tip: Use tools such as Cal-Adapt to help you determine the climate projections in your area.

Potential Impacts: What would change in terms of your on-farm operation with these projected changes? Would a longer growing season provide opportunities for new crop rotations? What about pest pressure? Consider both the potential challenges and opportunities that could come from the anticipated climate projections for your location.

Vulnerability: How vulnerable is your farm to these projected climate changes? Consider whether your farm expects few, some, or many impacts, and the capacity your farm has to cope. Would you describe your farm as having Low, Moderate, or High vulnerability?

Low – Your farm is ready and able to cope with the potential climate change impacts.

Moderate – Your farm will experience some challenging climate change impacts, but it will be able to cope with others.

High – Your farm's ability to cope with the impacts of climate change is limited and climate change will disrupt many important aspects of your farm's operations.

STEP 3 RE-EVALUATE GOALS AND OBJECTIVES GIVEN PROJECTED IMPACTS AND VULNERABILITIES

Understanding how climate change can affect the successful implementation of management goals and objectives allows producers to re-evaluate farm management goals and objectives in light of projected impacts and potential vulnerabilities.

By evaluating management objectives with climate change in mind, producers can revise their management plan to be better suited for a future of environmental change. For example, the management goal outlined in Box 1 was to decrease the occurrence or impact of salinization, with an objective of reducing the salts in soils via leaching. However, if a vulnerability assessment shows that this objective is not feasible due to reduced water availability, it would be necessary to revise the objectives in order to achieve the reduced salinization goal (Box 2).

Box 2. When the original management goals and objectives are considered in light of climate impacts, a revised objective can help you meet your management goals despite climate challenges.

Original Management Goal: Decrease occurrence or impact of salinization
Original Management Objective: Reduce salts in soils via leaching
Climate Change Impact: Reduced water availability due to drought may make leaching infeasible
Revised Management Objectives*:
1. Reduce application of fertilizers and/or salt-containing amendments like manure
2. Plant crops that are more salt tolerant

*UCANR Publication 8550 Managing Salts by Leaching

How will climate change make my goals more challenging?

Do I need to change any of my goals or management objectives given my farm's climate vulnerability?

Are there any opportunities that climate might present for my operation?



Figure 5. When re-evaluating management goals and objectives given climate change implications, it can be helpful to consider your farm's unique vulnerabilities to climate change, along with potential opportunities the future may bring.

Identifying areas of potential difficulty in light of the effects of climate change can help you determine the inputs or resources needed for climate adaptation (Figure 5). Inputs are anything put into the farm system to improve it, such as capital investments, infrastructure, labor, or soil amendments. Inputs also support the **actions** taken to achieve goal-driven **outcomes**. In the example listed in Box 2, selecting the fertilizer based on baseline salinization concerns is the action, fertilizer type is the input, and the resulting salinization levels after using the new fertilizer is the outcome. Once you have revised your goals and objectives in light of climate change, the next step in this workbook introduces a menu of adaptation strategies, approaches, and practices that provide a means to select your climate-adaptive actions for farm management.

STEP 3 PUT IT INTO PRACTICE

In the **STEP 3** worksheet [below](#) you have an opportunity to recall the management goals and objectives you initially outlined in Step 1. In this step, consider how your Step 1 goals and objectives may be affected by climate change.

First, refer back to your worksheets for Steps 1 and 2. What are some of the challenges and opportunities that climate change presents for your farm management goals and objectives?

Management Goals and Objectives: What management goals and objectives did you outline in Step 1?
Use these to guide your answers to the following questions.

Challenges: Consider your responses to your Step 2 worksheet. How will climate change make it more difficult to carry out your objectives and meet your goals?

Opportunities: Will climate change provide opportunities to further your goals and objectives?

After completing Steps 1 and 2, if you have determined that climate change will require a change in your management plan, use the boxes below to organize your [re-evaluated goals and objectives](#) with climate change vulnerabilities and challenges in mind.

Re-Evaluated Management Goal: What is your new or adjusted long-term management goal?

Re-Evaluated Management Objectives: What are your new or adjusted objectives? Do you need to reconsider your baseline for measuring progress?

STEP 4 IDENTIFY ADAPTATION STRATEGIES, APPROACHES, AND PRACTICES

Central to the adaptation process is identifying appropriate actions that can buffer a farm to the environmental shocks caused by climate change. In this step, an adaptation "menu" is presented. It offers science-based information for developing a farm management plan that incorporates climate adaptation. The menu consists of a collection of potential adaptation applications, organized by strategy, aimed at addressing the concerns related to climate change.

What is the Adaptation Menu?

In order to develop a successful adaptation plan, producers will need to identify a set of actions that can help them meet their climate-adjusted goals and objectives. The adaptation menu provides a starting point in selecting appropriate actions for adapting an operation to the current and future climate. The menu is organized by adaptation **strategies**, which are very high-level actions that may be applicable across landscapes and production systems. Within each strategy, **approaches** provide guidance on how a particular strategy might be implemented and more detailed information on the importance of the approach for agricultural adaptation. Each approach may be adopted by enacting one or more **practices** (Figure 6). The practices included in the menu do not represent an exhaustive list, but they are potential on-the-ground applications that accommodate the selected approach and strategy. The menu focuses on adaptation, but some approaches and practices can provide the secondary benefit of mitigating climate change by reducing greenhouse gas emissions or facilitating carbon sequestration.

*Note that Strategy 7 is entirely focused on mitigation efforts.



Figure 6. The adaptation menu offers a hierarchical organization that provides high-level strategies as a starting point, with more specific approaches and practices therein to guide the process of adaptive management planning from concept to action.

THIS MENU DOES

- Provide a starting point for exploring and identifying a range of potential actions to support on-farm adaptation.
- Help producers meet their management goals and objectives in light of climate change.
- Provide examples of practices that can be used to implement producer-prioritized climate adaptation strategies.

THIS MENU DOES NOT

- Make specific management recommendations.
- Prioritize or give preference to any particular strategy, approach, or practice. Many factors will determine what is best for your operation.
- Provide an exhaustive list of all possible practices for meeting strategic climate adaptation goals and objectives.

Box 3. Menu of Adaptation Strategies and Approaches for California Specialty Crops

Strategy 1: Support and Maintain Soil Health, Soil Biological Services, and Water Quality

Approach 1.1: Maintain and Improve Soil Structure

Approach 1.2: Enhance Soil Biology and Nutrient Availability with Organic Amendments

Approach 1.3: Manage for Soil and Water Salinization

Approach 1.4: Manage Fertilizer Application to Protect Groundwater Quality and Soil Health

Strategy 2: Cope with Uncertain Water Availability

Approach 2.1: Improve or Alter Water Systems to Meet Current and Expected Future Demands

Approach 2.2: Manage for the Effects of Drought

Approach 2.3: Prepare for Changing Patterns of Precipitation

Strategy 3: Manage Biological Crop Stressors

Approach 3.1: Reduce Pest Pressure

Approach 3.2: Reduce Disease Risk from Pathogens

Approach 3.3: Reduce Weed and Invasive Plant Pressure

Strategy 4: Prepare for Temperature Change

Approach 4.1: Adapt to Warmer Conditions

Approach 4.2: Manage for Changing Seasonality

Approach 4.3: Prepare for Longer-Term and Larger-Scale Temperature Changes

Strategy 5: Enhance Preparedness for Extreme Events

Approach 5.1: Prepare for Extreme Heat Events

Approach 5.2: Manage for Extreme Precipitation and Flooding

Approach 5.3: Enhance Resilience to Extreme Wind

Approach 5.4: Prepare for and Respond to Wildland Fire and Smoke Impacts

Strategy 6: Manage Farms and Fields as Part of a Larger Landscape

Approach 6.1: Integrate, Improve, or Maintain Natural Ecosystems within Agricultural Operations

Approach 6.2: Promote Biological Diversity Across the Landscape

Approach 6.3: Conserve Individual Farms

Strategy 7: Develop or Expand Co-Benefit Efforts to Mitigate Climate Change

Approach 7.1: Reduce On-Farm Greenhouse Gas Emissions

Approach 7.2: Increase On-Farm Soil Carbon Sequestration

STEP 4 PUT IT INTO PRACTICE

In **STEP 4** you have an opportunity to consider and identify the adaptation **strategies, approaches, and practices** that will help you realize your goals and objectives while accounting for the effects of climate change.

Use the **menu outline** in Box 3 above to help you narrow down your list of potential actions. Refer to the **detailed menu** provided after Step 5 to learn more about the importance of each strategy and approach, and see some example practices that can be put into place. Once you have identified a few adaptation options, you can use this template to organize and record your thoughts.

Management Goals and Objectives: Consider your revised goals and objectives from Step 3, if applicable.

Challenge or Vulnerability: What climate challenge or vulnerability are you addressing?

Current Strategies, Approaches, and Practices: Are there any strategies, approaches, and/or practices that you have previously implemented? What challenge/vulnerability were they addressing, and did they help you to meet your management goals and objectives?

New Strategy, Approach, and Practice: Which strategy, approach, and practice(s) have you selected to address the identified challenge/vulnerability and help you meet your management goals and objectives? Remember, the menu does not provide an exhaustive list of all possible adaptive actions that you can take!

Benefits: What benefits and/or opportunities could your selected approach and practice(s) provide? On what time scales?

Current Drawbacks and Barriers: If you have previously implemented climate-smart strategies, approaches, or practices, what drawbacks or barriers have you faced? How did you overcome those barriers?

New Drawbacks and Barriers: What drawbacks or barriers might you face in adopting the selected approach and practice(s)? How might those barriers be overcome?

STEP 5 MONITOR AND EVALUATE

After implementing one or more adaptation practices, it is important to **monitor and evaluate** the effectiveness of those actions in order to ensure long-term success. This step in the adaptation process is about identifying and reflecting on questions that will help determine whether the implemented practices are supporting the farm management goals and objectives, or if adjustments should be made. A monitoring plan can also help producers decide whether various approaches should be prioritized or additional practices executed as new climate observations are made, as new information or resources become available, or as long-term farm management goals change.

Some considerations when developing a monitoring plan include:

- What will be monitored to assess effectiveness? Depending on the practice enacted, this could be variables like soil characteristics, water use, or crop damages.
- How will you measure these indicators of effectiveness?
- When, where, and how often will you measure your indicators?
- What is your threshold for success? In other words, by how much do you want to increase soil health measures, curb water use, or reduce crop damages?
- Do you have the information and tools you need in order to carry out the monitoring and take your measurements?

Making sure that a monitoring plan is feasible – in terms of time, cost, and effort – is a key to its success.

When evaluating the results of your monitoring efforts, consider these questions:

- Have I met my threshold for success? Or am I at least trending in the right direction?
- Are the implemented practices helping me to achieve my management goals?
- Do my monitoring observations indicate that my management objectives should be re-evaluated?
- Am I still monitoring the most useful indicators, and do previously adopted practices still fit within my *current* farm management plan?

Regular and consistent evaluation of management plans and implemented adaptation practices is crucial in helping farms adapt to a changing climate.



STEP 5 PUT IT INTO PRACTICE

In **STEP 5** you are tasked with developing a **monitoring and evaluation (M&E)** plan to assess the effectiveness of the adaptation action(s) you selected in Step 4. Although you will not *carry out* your M&E plan until after you have implemented your action(s), thinking carefully about the M&E process or developing a plan prior to enacting on-farm changes may be useful for identifying the potential limitations and ensuring that your chosen action(s) are justifiable in the long term.

For developing your M&E plan, use the **MONITORING WORKSHEET** below.

Action(s) or Practice(s):

Measurable Indicator: Choose one. It can be general, such as farm water use, or specific, such as soil infiltration rates.

Tools and Methods: How will you monitor for change?

Schedule: When will you monitor?

Sites: Where will you monitor, and how will the sites be marked?

Name and Cost of Lab Analyses or Tools:

Baseline: Refer back to Step 1 if needed. If measurements have not been taken yet, is there time to do so before implementing your action(s) or practice(s)?

Interpretation: What outside information or help will you need to interpret the results?

Accuracy: In general, how accurate is the measurement method? How "noisy" or variable is the indicator, year-to-year or site-to-site? How many measurements are recommended in each round?

Target: What is your numeric threshold for success?

Time Scale: When do you expect to reach your numeric target?

Once you have successfully enacted your monitoring plan, use the **EVALUATION WORKSHEET** below to assess whether your actions have helped to achieve your goals and objectives. Farmers who have already been engaging in adaptation practices can also benefit from addressing these evaluation questions.

Progress Towards Adaptation Goals: In what ways are your actions helping you to achieve your management goals?

Updating Whole-Farm Adaptation Goals: How have the outcomes of this adaptation action changed your farm management goals and objectives?

Reaching Target(s): How have your indicators progressed in comparison to the thresholds for success?

Other Indicators or Methods of Measurement: Can other indicators be useful? How about different measurement methods for the established indicators? Are new measurement technologies or cheaper measurement methods available?

Improving Practice(s) Effectiveness: Are there any changes that you can make to improve the effectiveness or efficiency of your implemented action(s) or practice(s)?

Maintenance of Practice(s): How consistently, and how intensely, do your action(s) or practice(s) need to be implemented going forward in order to meet your goals?

Next Steps: How can you further advance your farm management plan and adaptation needs? Are there actions that should be expanded, or complementary practices that can be adopted?

A Menu of Adaptation Strategies and Approaches for California Specialty Crops

Strategy 1: Support and Maintain Soil Health, Soil Biological Services, and Water Quality

Healthy soils support the productivity and profitability of diverse agricultural enterprises, and they regulate water quality and quantity while providing for many agronomic functions and ecosystem services.^{20,21} These functions and services depend on the physical, chemical, and biological properties of the soil, some of which are dynamic and easily altered (e.g., soil biodiversity and structure), while others are inherent or more resistant to change (e.g., soil texture and color). Unfortunately, climate change creates multiple direct and indirect threats to soil health, such as disruption of soil structure from extreme weather events,²² decreased soil microbial diversity,²³ and soil and water salinization.²⁴ To buffer against these climatic impacts, producers may consider adaptive management approaches and practices such as those listed below.

Approach 1.1: Maintain and Improve Soil Structure

Soil structure results from the aggregation of soil particles, which is strongly dependent on particle size distribution ("texture"), charge-based attraction between clay particles, and organic "glues." Well-structured soils are those that have a better ability to draw in (infiltrate) and store water, to support microbial activity, and to support root health. They remain more stable under pressure, are easier to till, and can require less tillage. When the structure is promoted, the soil's functions can better withstand conditions like drought and extreme precipitation (see also Strategies 2 and 5).

Soil structure can be maintained and even improved through practices that increase soil organic matter; practices that protect soils from rain impacts, wind erosion, and heating; or practices that reduce tillage and the additional loss of carbon that comes with the break-up of aggregated soil particles.²⁵ For example, minimum tillage with the retention of residues reduces the amount of soil lost to storm runoff and also reduces evaporative water loss from the soil surface.²⁶ Alternatively, cover crop rotations may require some degree of tillage, but they can shield the soil from rain, improve drainage, and re-build degraded soils.²⁶

Examples of adaptation practices include:

- **Practice 1.1.1: Minimize or eliminate tillage** and other traffic to improve soil structure and fertility, ideally retaining residues as well.^{27,28,29}
- **Practice 1.1.2: Include selected cover crops in field rotations or in orchard alleys** to protect soil, break up compacted layers,³⁰ produce channels for water that maximize infiltration, water storage, and irrigation water use efficiency,³¹ and contribute carbon exudates for promoting microbial activity and soil aggregation.
- **Practice 1.1.3: Implement reduced-till cover cropping** by mowing, flaming and/or adding herbicide to avoid additional disking or disruption. Shallow disking or direct seeding may be adequate for establishment.³²
- **Practice 1.1.4: Promote healthy root growth** through practices such as optimizing irrigation and nutrient placement, or applying compost teas,³³ humic acids,³⁴ mycorrhizal inoculations when useful,³⁵ and gypsum amendments on saline soils.³⁶

Approach 1.2: Enhance Soil Biology and Nutrient Availability with Organic Amendments

Soil biological processes are considered to be the basis of health and functionality in any agricultural system,³⁷ so it is vital to maintain a suitable environment for soil microorganisms in all farming systems. At the same time, climate change is expected to have both direct and indirect negative impacts on certain soil biological processes,³⁸ soil biodiversity,³⁹ and soil health generally.^{40,41} As a result, some studies suggest that climate change will decrease surface soil organic matter,⁴² with impacts on fertility and biodiversity.³⁹

Organic amendments have been applied to cropped soils for millennia,⁴³ serving to restore some of the biological and nutrient cycles that are interrupted by the harvesting of crops. They contain nutrients and carbon substrates that microorganisms use for energy. The metabolic activity of soil microbes produces polysaccharides that act as organic soil "glues," which help soil particles to form stable aggregates and retain

nutrients through charge interactions. Organic amendments can thereby increase the rates of nutrient cycles, provide macro- and micro-nutrients, and increase soil biodiversity, and thus they can have benefits for pest and pathogen control,⁴⁴ and for enhanced crop production.⁴⁵

The selection of suitable organic amendments depends on soil type, cropping system, and irrigation practices. Growers may be well advised to undertake soil sampling in order to understand the specific composition and condition of their soil.

Examples of adaptation practices include:

- **Practice 1.2.1: Apply compost**⁴⁶ while paying attention to its feedstocks, nutrient contents, and pathogenic analyses where relevant, to build up the organic matter and its functions in the soil.
- **Practice 1.2.2: Apply a surface mulch** of wood chips, straw, or greenwaste to help reduce evaporation, suppress weeds, and build soil organic matter over time.^{47,48} Greenwaste mulches should always be checked against food safety requirements.
- **Practice 1.2.3: Apply manure (fresh, slurry, compressed, pelletized, or composted)** to supply soil with nitrogen, phosphorus, and micronutrients.⁴⁹ Check food safety requirements.
- **Practice 1.2.4: Retain crop residues, or allow animals to graze them**, which will return nutrients and organic matter to the soil through digestion and decomposition.
- **Practice 1.2.5: Consider undertaking Whole Orchard Recycling (WOR).** WOR incorporates large amounts of wood chips into the soil to improve soil functions including nutrient uptake, aggregation, and water retention.⁵⁰ Crop nitrogen needs may be moderately higher for a few years after incorporation, as the wood chip decomposition process creates a sink for nitrogen.⁵¹
- **Practice 1.2.6: Incorporate biochar into the soil** to retain nutrients⁵² and increase carbon, among other possible benefits. The specific feedstock and preparation strongly affect its performance as do the properties of the soil to be amended.
- **Practice 1.2.7: Incorporate cover crop rotations** into both annual and perennial systems⁵³ to boost productivity while providing organic matter, capturing nutrients that could otherwise leach out of the soil, and fixing nitrogen by using leguminous covers.

Whole Orchard Recycling at Rock'N Almonds

Whole Orchard Recycling (WOR) is the practice of grinding whole trees into chips once they have reached the end of their agronomic lifecycle. Incorporating these chips into the soil can **increase soil organic matter, improve water holding capacity, and increase yield**. Seeing the benefits of this practice can take anywhere from 2–5 years, but adopters say it is worth the wait.

Rock'N Almonds needed a practice that would work for them long term as they operate in the Central Valley, an area prone to sustained drought and extreme heat. So, in 2015 they started incorporating WOR on a block of their orchard. They first cut down and chipped up trees that had reached the end of their lifecycle and incorporated the chips into the soil. After letting the material break down for about 2 years, they planted new trees.

In comparison to their conventionally managed blocks, **they found the WOR block soil to have higher organic matter and carbon content, and enhanced water holding capacity**. Together, these benefits helped Rock'N Almonds increase yield and cut down on labor cost, which helped them reach their management goals of remaining profitable.

[See more info on Whole Orchard Recycling in California here!](#)

Approach 1.3: Manage for Soil and Water Salinization

As water supplies in California have constricted, irrigation water quality and soil salinity have worsened. When surface water is limited, the increased use of groundwater for irrigation can exacerbate salinity issues in some locations, depending on the salinity of the groundwater. Increases in soil salinity can create plant water stress, requiring more irrigation for crop needs and for leaching fractions.⁵⁴ Drainage management is also critical for managing salts in certain soils. This is seen in parts of the Delta and in the western San Joaquin Valley, where saline water tables that extend up to the root zone have aggravated the effects of naturally high soil salinity.⁵⁵ Increased soil salinization can have detrimental impacts on crop yield⁵⁶ and crop quality.⁵⁷ Certain specialty crops, including vegetables, are especially vulnerable to soil and water salinization.⁵⁸ This poses a particular problem in the vegetable-producing coastal areas where groundwater depletion has increased saltwater intrusion into groundwater

aquifers.⁵⁹ Under climate change and in the absence of adaptive action, California producers can expect increasing salinization challenges.²⁴ Fortunately, proper soil and irrigation management techniques at the field level, such as the practices listed below, can ameliorate some of the impacts of salinization. Prior to enacting these practices, consider soil and irrigation water testing by a reputable lab to guide your management decisions.

Examples of adaptation practices include:

- **Practice 1.3.1: Test irrigation water sources and agricultural soils** for specific salt species and levels. Many laboratories will make corresponding management recommendations.
- **Practice 1.3.2: Monitor plant symptoms and salt levels in annual tissues or leaves.** In specialty crops, leaf sampling for nutrient levels is a complementary tool, following UCANR critical thresholds⁶⁰ and sampling protocols.^{61,62}
- **Practice 1.3.3: Consider the salt contents in fertilizers before application**, making note of existing soil solution salinity levels and sodium absorption ratio testing.⁶³
- **Practice 1.3.4: Reposition driplines, reschedule irrigations, or shift irrigation methods** to address localized salinization.^{64,65,66}
- **Practice 1.3.5: In groundwater-dependent areas, consider annual maintenance leaching**, especially when growing vegetables.^{67,68} In California, farmers may be able to use floodwaters through field-based **"managed aquifer recharge" (Flood-MAR)**. **In other cases, timing a leaching fraction after major winter rains increases its effectiveness.**
- **Practice 1.3.6: Where practical, consider installing tile drains** to extract saline leaching water and prevent further salinization of groundwater.
- **Practice 1.3.7: Consider applying compost** to improve water uptake and nutrient availability in saline soils.⁶⁹ The salt contents of any applied compost should be assessed before use. For example, manure composts typically have higher salts, while greenwaste composts have less.



Approach 1.4: Manage Fertilizer Application to Protect Groundwater Quality and Soil Health

Groundwater is a vital component of California's water supply, with 40% of California's urban and agricultural areas relying on groundwater resources,⁷⁰ especially in times of drought⁷¹ (see [Strategy 2](#) for more on drought). Agricultural intensification has significantly increased food production since the mid-20th century. Unfortunately, the increased fertilizer application also increased the hazards for the state's water resources, including groundwater. Inefficiencies in nutrient cycling and crop use of applied nutrients can allow elements from the fertilizers and manure such as sodium, chloride, and nitrogen to reach groundwater, causing salinization, diminished soil health, and threats to human health.⁷² Improved nutrient and water management in recent decades has reduced agriculture's impacts on groundwater, and it is important to build upon those gains.

Focusing on nitrogen (N), which has the most complex cycle of any applied nutrient, excessive applications of N can allow bacteria to oxidize and consume soil organic matter.⁷³ On the other hand, if adequate N is not provided, crops may "mine the soil" by consuming N from soil organic matter, and thus degrading that organic matter over the long term.⁷⁴ Efforts to apply nitrogen accurately in California have found that irrigating with groundwater that has high nitrate levels can upset crop development, delay maturity, and reduce yield,^{75,76} so the background nitrate content of groundwater should be taken into account.

Fortunately, adaptive water-saving technologies also allow more precise applications of fertilizer to best match the 4 R's of nutrient management – the Right Amount, at the Right Time (coinciding with crop uptake), in the Right Form, and in the Right Place. These refinements in fertility management reduce the risk of moving nutrients below the rooting zone, where they are out of reach of crop roots and pose a nutrient loading hazard to groundwater.

Examples of adaptation practices include:

- **Practice 1.4.1: Develop a nitrogen (N) budget.** A table of N inputs including soil and water N, and exports including likely gaseous and leaching losses, can form the basis of efficient fertilization. Farmer-oriented computational tools are also available and they are becoming widely used for compliance and management.
- **Practice 1.4.2: Monitor soil and plant N levels during the growing season.** Soil nitrogen "quick tests" can be performed by farm staff without extensive training,⁷⁷ while monitoring leaf N concentrations is a useful tool in many specialty crops.⁶³
- **Practice 1.4.3: Manage N applications** to avoid spiking N in the soil beyond the crop's uptake capacity. Particularly on sandy soils, or soils with low organic matter, it may be important to change the N fertilizer formulation, change the frequency of N applications, consider controlled-release fertilizers or nitrification inhibitors, or even add biochar to the soil, which can retain N.⁷⁸
- **Practice 1.4.4: Optimize the use of fertigation** considering fertilizer formulation, high-frequency application, emitter placement, and the timing of fertilizer injection within the irrigation set, to place N at the root depth in the soil and avoid leaching.⁷⁹
- **Practice 1.4.5: Manage manure applications** by considering likely decomposition and N mineralization rates, as well as the contents of various nutrients. A wide range of salts and nutrients found in manure can affect groundwater quality, especially if their release does not correspond to crop needs.

Strategy 2: Cope with Uncertain Water Availability

Adequate water resources are imperative for agricultural production. While sensitive to the seasonal and year-to-year variability of the state's Mediterranean climate, California's extensive water infrastructure supports the state's expansive agricultural sector and has buffered it from in-season water shortages. However, even with the most elaborate water conveyance and storage system on earth, the state's water resources will be challenged to keep pace with the changing water needs driven by climate change. Production systems will need to endure increasing frequencies of both droughts and very wet years, as projected under future conditions.^{80,4} Such changes in water availability can have severe consequences for agricultural production if not properly managed. Outlined below are a series of approaches, each with examples of practices, that can aid in managing for and coping with uncertain water availability.

Approach 2.1: Improve or Alter Water Systems to Meet Current and Expected Future Demands

Approximately three quarters of California's cropland is irrigated, with more than one third of irrigated acreage producing high-value perennial fruits and nuts.⁸¹ Crop production in California uses more than 25 million acre feet of irrigation water per year, accounting for between 40%–60% of the state's total water supplies.⁸² Climate change is expected to increase agricultural water use as warmer temperatures increase crop transpiration rates and evaporation losses reduce available soil moisture. Across California's agricultural lands, climate change is projected to increase summer reference evapotranspiration by ~5–15% over historical averages by the end of the 21st century.⁸³ Likewise, warming temperatures will reduce snowpack and decrease runoff during the summer growing season, reducing surface water availability.² The warming-driven increases in water demand coupled with decreases in water availability will require the refinement of on-farm water management to optimize water use efficiency.

In recent decades, upgrades to irrigation systems have been shown to reduce irrigation water usage in some Mediterranean climates.⁸⁴ In order to be resilient against future conditions, irrigation systems will likely need to be altered with innovative methods that increase infiltration and enhance water use efficiency.⁸⁵ For example, when more efficient on-farm irrigation infrastructure is installed, irrigation water use typically decreases; however, under future conditions, California farmers may need to upgrade or enhance more than 30% of their irrigation infrastructure to maintain or reduce irrigation use below recent levels.⁸⁶ These alterations will provide a critical buffer, allowing irrigation to continue during future water-restricted years. The following practices provide examples for updating or enhancing irrigation systems to adapt to current and expected future water demands.

Examples of adaptation practices include:

- **Practice 2.1.1: Update existing irrigation infrastructure to improve irrigation efficiency.**^{87,88} For example, install irrigation blocks that follow soil type where possible and consider hybrid approaches such as combining subsurface drip and furrow.
- **Practice 2.1.2: Maintain irrigation systems for optimum performance and distribution uniformity.** Maintenance is also important for fertigation systems.
- **Practice 2.1.3: Consider adopting or expanding irrigation technology use.** Advances in soil moisture sensors, cloud computing, and estimates of field-scale ET via surface renewal and remote sensing approaches provide opportunities to improve water use efficiency and optimize irrigation scheduling.⁸⁹
- **Practice 2.1.4: Create a [water budget](#) to better assess water demands.** Using a [water budget tool](#) can help measure the efficiency and regional suitability for the amount of water applied to a landscape based on local climate data.⁹⁰

Minimum Till at Full Belly Farm

Full Belly Farm is a diverse, 400-acre farm in Yolo County's Capay Valley, where more than 80 different crops are produced. One of the projected impacts of climate change to Full Belly's operation is an increase in the frequency and severity of drought. The managers at Full Belly Farm have adopted **minimum till practices** to support their farm management objectives for maintaining soil health and their long-term goals of resilience and sustainability.

Minimum tillage avoids unnecessary disruption of the soil while allowing for crop-specific needs, such as shallow intrusions for planting. This practice **improves soil health** in the short term and increases the soil's ability to **hold onto water** in the longer term, which can enhance the farm's ability to **bounce back from events like drought**.

Full Belly Farm's minimum till practices include rolling down cover crops and moving residue to create space for planting vegetables with minimal soil disturbance. Despite the added labor in managing weed pressure, Full Belly believes **the benefits of minimum till outweigh the costs**, and the farm is undertaking other minimum till experiments with the goal of expanding this practice across the farm.

[Check out this University of California video on Minimum Tillage Systems!](#)

Approach 2.2: Manage for the Effects of Drought

Periodic drought is the norm rather than the exception in California. Drought can challenge a grower's ability to meet crop water demands. Increased water stress can slow plant development, reduce nutrient uptake, decrease crop yield, and in extreme cases kill much of the crop.² Climate change is expected to increase the risk of drought (including multi-year and multi-decadal droughts) as warmer temperatures drive higher evaporation rates and increase the atmospheric water holding capacity.^{91,92,93} These 'hot droughts' can be especially pernicious, as the decreased soil moisture serves to increase soil temperatures and raise surface-level air temperatures, creating a positive feedback loop that can deepen existing drought conditions.⁹⁴ Such changes make it imperative that producers carefully manage drought conditions.

In recent decades, growers have primarily used groundwater to lessen the effects of decreased surface water availability,⁹⁵ and in recent years, groundwater demand during drought has been growing due to the expanded acreage of perennial crops. Subsidence, overdraft, and changes in water flow patterns may challenge the capacity of groundwater to safeguard against future drought.⁹⁵ Management practices such as those that improve soil health and soil water holding capacity (Approaches [1.1](#) and [1.2](#)) or increase irrigation efficiency (Approach [2.1](#)) can provide a buffer against drought impacts. California producers may wish to consider multiple adaptive practices to enhance their resilience.

Examples of adaptation practices include:

- **Practice 2.2.1: Employ reduced tillage, cover cropping, and organic amendment practices to reduce soil water evaporation⁴⁷ and improve water infiltration and retention.⁹⁶***Note that reduced tillage may increase weed pressure (see [Approach 3.3](#) for information on weed pressure). Cover cropping opportunities may be limited in some systems (see [Approach 1.1](#) for more on BMPs for soil).
- **Practice 2.2.2: Utilize deficit irrigation strategies** to balance production goals with available water resources during drought. This practice works best with tree crops and vines.^{97,98,99}
- **Practice 2.2.3: Plant drought tolerant varieties¹⁰⁰ and/or shift to drought resistant rootstocks¹⁰¹** to maintain or enable crop production where water is scarce (see [Approach 4.3](#) for more information on incorporating new varieties).
- **Practice 2.2.4: Adopt conservation-oriented irrigation scheduling practices** such as soil moisture monitoring, evapotranspiration (ET) scheduling, and plant-based irrigation scheduling using temperature or water potential monitoring.¹⁰²

Approach 2.3: Prepare for Changing Patterns of Precipitation

California has historically received ~75% of its annual precipitation between November and March,¹⁰³ with more than 30% of the state's annual precipitation stemming from atmospheric river storms.¹⁰⁴ Observed daily precipitation patterns across the Western U.S. show increasing durations between precipitation events¹⁰⁵, and climate models project increasing year-to-year variability of atmospheric river storms¹⁰⁶ and a shortening of California's wet season.¹⁰⁷ Precipitation timing can strongly influence management decisions such as planning field work, weed and pest management, and spray regimes.¹⁰⁸ Challenges resulting from changing precipitation patterns can be ameliorated with proper management. For example, during unusually wet periods, growers may prepare for later needs using water catchment systems and management plans that aid aquifer recharge (e.g., flooding of orchard crops during the dormant season¹⁰⁹). As climate change alters the timing of the wet season and increases the likelihood of large swings in annual precipitation extremes,⁴ the adoption of on-farm recharge efforts, including optimizing infiltration and minimizing runoff, as well as the refinement of irrigation water use efficiency, will be essential. Additional practices for adapting to changing precipitation patterns can be found in [Approach 5.2](#).

Examples of adaptation practices include:

- **Practice 2.3.1: Incorporate swales into the landscape to slow the movement of water.**⁹⁶ Swales can reduce soil erosion ([Strategy 1](#)) and encourage water infiltration into soils, providing a buffer against drought.
- **Practice 2.3.2: Use retained stubble/residues or mulch** to protect the soil surface and prevent soil crusting, resulting in reduced runoff.²⁸ Residues can also support soil health and soil water retention (see [Approach 1.2](#)). However, they may not be appropriate in some vegetable systems.
- **Practice 2.3.3: Carry out annual cropping, cover cropping, or other plantings on temporarily fallowed land** to support soil health, which can increase soil water holding.
- **Practice 2.3.4: Implement agricultural managed aquifer recharge (AgMAR)** to increase groundwater recharge and counteract groundwater overdraft.^{110,111} Leave flood infrastructure intact in areas where surface water deliveries may allow groundwater recharge in the future.⁸⁷



Allowing fallowed fields or flood-tolerant crops to absorb excess rainwater can encourage groundwater recharge.

Strategy 3: Manage Biological Crop Stressors

Due to the direct impact of temperature on the growth and life cycles of various biological stressors of crops (e.g., insects and weeds), observed and projected increases in temperature will have significant implications for agricultural systems.² Agricultural insects and diseases account for a large percentage of crop yield damages every year,¹¹² while weeds can stress plants by competing for water and nutrient uptake. Under climate change, insect,¹¹³ disease,¹¹⁴ and weed pressures¹¹⁵ are projected to increase, requiring changes in pest and weed management systems. In order to alleviate some of the risks associated with biological stressors, it is critically important to consider related best management approaches in farm operations and planning.

Approach 3.1: Reduce Pest Pressure

Because insect pests are cold-blooded, their growth rates and lifecycles are largely regulated by weather conditions and can increase as temperatures rise.^{116,117} California's agricultural sector has already seen shifts in pest pressure, due in part to warmer average temperatures.² The continuation of warming temperatures driven by climate change are projected to alter insect migrations and developmental rates, and increase the number of pest generations born each year.^{113,118} Warmer temperatures can also increase insect outbreaks,¹¹⁹ and lead to the occurrence of new non-native pests affecting crops.² Reducing pest pressure will require the implementation of both preventive as well as control measures, and given the economic impacts of pest damages,^{113,120} producers should take an active role in implementing improved pest management practices.

Examples of adaptation practices include:

- **Practice 3.1.1: Follow the latest integrated pest management guidelines** using Prevention, Avoidance, Monitoring, and Suppression (PAMS) strategies to optimize pest management. Implement preventive and control measures, such as choosing varieties adapted to current conditions; optimizing nutrient and water management; removing diseased plant material or pest host material (e.g., sanitizing nut orchards during the winter); harvesting crops early to minimize pest damage risk; disrupting pest mating (e.g., by pheromones or sterile insect release); and biological control.^{121,122,123}
- **Practice 3.1.2: Create an environment that is inhospitable for the pests** by employing soil solarization, steam, and anaerobic soil disinfection.¹²⁴
- **Practice 3.1.3: Create habitat for beneficial insects that prey on agricultural pests and enhance natural enemy populations** through perennial hedgerows, wildflower areas, flowering islands, etc.^{125,126}
- **Practice 3.1.4: Ensure that monitoring and scouting efforts during the season match current conditions** to effectively time insecticide applications for reducing pest pressure.



Approach 3.2: Reduce Disease Risk from Pathogens

Plant pathogens are detrimental to crop health and pose a major threat to regional and global food supplies.¹²⁷ Globally, a warming-driven poleward expansion of crop pathogens has been observed.¹²⁸ In California, more than \$34 million in indemnity payments have been made to farmers due to pathogen losses since 1989, with an apparent positive trend in indemnity claims and the three most costly years occurring in the last decade according to [AgRisk Viewer](#)¹¹². Increasing pathogen risk is expected under climate change, as warmer weather can influence the spread and emergence of many pathogens.¹²⁹ For example, warmer weather during the typically cool winter season can increase the prevalence of Pierce's Disease in grapevines.¹³⁰ Climate change may also reduce the occurrence of frosts in the future,¹³¹ which could increase the risks and management challenges associated with pathogens and other biological crop stressors. While some of the impacts of pathogens associated with climate change might be inevitable, it is important to implement adaptive management practices to control and prevent pathogen spread as much as possible.

Examples of adaptation practices include:

- **Practice 3.2.1: Decrease pathogen spreading capability** by intercropping disease resistant and disease susceptible varieties.¹³²
- **Practice 3.2.2: Include other intercropping strategies such as strip intercropping, row intercropping, relay intercropping, and intercropping of genetic variants** to reduce disease.¹³²
- **Practice 3.2.3: Incorporate Biological Control Agents** that ecologically and environmentally reduce pathogen pressure.¹³³
- **Practice 3.2.4: When possible, utilize early warning systems** for diseases.^{134,135}
- **Practice 3.2.5: Use weather-based decision support tools** for monitoring and scheduling fungicide applications (e.g., Strawberry Advisory System; www.agroclimate.org).

Approach 3.3: Reduce Weed and Invasive Plant Pressure

Weather and climate have direct influences on weeds and invasive plants. Increasing temperatures, shifting precipitation patterns, and elevated CO₂ concentrations are expected to favor highly adaptive weeds and invasive plants.¹³⁶ In experimental trials, invasive plants in Mediterranean landscapes were shown to increase their reproductive output under projected climate change scenarios.¹³⁷ Similarly, weeds are expected to naturally adapt under a changing climate,¹³⁸ potentially allowing them to expand their ranges and compete with production agriculture for water and nutrient uptake.^{115,139} In the absence of careful management, these pressures can severely impact crop growth and yield.¹³⁸ While herbicide application has historically been a successful means of mitigating weeds in agricultural settings, climate change is expected to reduce herbicide efficacies, which can be impacted by various environmental conditions.¹⁴⁰ Looking forward, reducing weed pressure and invasive plant pressure in agricultural systems is crucial. Provided below are a few examples of high-priority practices that could be implemented.

Examples of adaptation practices include:

- **Practice 3.3.1: Include cover crops** to reduce weed pressure and enhance soil quality¹⁴¹ (see [Approaches 1.1](#) and [1.2](#) for more on cover crops).
- **Practice 3.3.2: Discourage weed growth by intercropping multiple varieties in one planting area, rotating crops after each lifecycle, and altering plant densities between seasons**¹³² (see [Approach 4.3](#) for more on intercropping and rotating).
- **Practice 3.3.3: Select crop cultivars that are highly adapted to climate variability** through better light interception or water and nutrient uptake, which can provide better competition and allow the crops to outperform weed growth¹¹⁵ (see [Approach 4.3](#) for more on cultivar selection).
- **Practice 3.3.4: Change seeding rates, row spacing, and row direction** to manipulate weed growth and control.¹³⁸
- **Practice 3.3.5: Survey farms in the fall and spring before the growing season to monitor weeds that escaped previous weed control efforts** and adjust the current weed management strategy to account for these weeds.¹⁴²
- **Practice 3.3.6: Consider weed management before planting.** It is easier and cheaper to control perennial weeds before planting a perennial crop because there is a wider selection of treatment options available when the ground is fallow.¹⁴²
- **Practice 3.3.7: Implement Integrated Weed Management**, a holistic approach to weed control (see the [Resources](#) section for crop-specific integrated weed management references).

Strategy 4: Prepare for Temperature Change

California's temperate climate, with warm summers and mild winters, has allowed high-value specialty crop production to thrive across much of the state. However, warming temperatures and shifting seasonality may challenge California specialty crop production by shifting planting patterns,² altering pollinator behavior,¹⁴³ lowering yields,¹³ and increasing water demand (see [Strategy 2](#)). As a result, producers may wish to employ adaptive approaches to help ameliorate the impacts associated with rising temperatures. The approaches below address the effects of warming average temperatures, changes in the timing of annual temperature patterns, and future temperature conditions that may be outside of the temperature regime seen today.

Approach 4.1: Adapt to Warmer Conditions

Though extreme temperature conditions are receiving a great deal of attention (see [Strategy 5](#)), changes in average or baseline temperatures can also influence crop geography, phenology, and quality. Both cool and warm season temperature increases are important considerations for California's specialty crop industry. California has experienced notable warming trends since 1895, with recent decades having greater rates of warming,² and models project that climate change will further increase temperatures over the coming decades.¹⁴⁴ These warming trends will have numerous implications for California's specialty crop production. Warmer winter temperatures will reduce chill accumulation and subsequently reduce both yields and the suitable cultivation areas for some high-chill crops like walnuts and pistachios.¹⁴⁵ Similarly, warmer summer temperatures can result in more rapid crop development in crops like processing tomatoes,¹⁴⁶ and reduced quality and reductions in suitable cultivation areas for crops like tomatoes, cantaloupe, and carrots.¹⁴⁷

The appropriate practices for addressing warming conditions will vary by geography, cropping system and planning time horizon. This Approach emphasizes practices that are useful for adapting crops in their current locations to warmer conditions, though additional efforts such as crop variety or species changes may be required over longer time horizons (see [Approach 4.3](#) below). Additionally, while warmer summer temperatures will increase water demand¹⁴⁸ and spur producers to employ practices that address irrigation (see [Strategy 2](#)), here we highlight non-irrigation management practices.

Examples of adaptation practices include:

- **Practice 4.1.1: Apply kaolin clay and/or dormant oil to nut crops to increase chill accumulation.**¹⁴⁸ The application of kaolin clay is most useful during sunny winters.¹⁴⁹
- **Practice 4.1.2: Where possible, use chill portions to calculate winter chill.** Chill portions have been shown to better reflect crop phenology in California.¹⁵⁰
- **Practice 4.1.3: Assess the need for earlier planting dates to avoid summer heat exposure** using localized, short-term summer warming projections.¹⁵¹
- **Practice 4.1.4: Prune the canopy or apply reflectants to reduce crop surface temperature.** Also consider incorporating plastic films, shade cloth, or netting to increase shade.^{152,153}
- **Practice 4.1.5: Include cover crops to reduce soil temperatures**¹⁵⁴ where appropriate. *Note that cover crops may not be suitable for all operations, they may compete with primary crops for water and nutrients, and in some circumstances, they may increase pest pressure.
- **Practice 4.1.6: Incorporate agrivoltaics into agricultural systems** to cool the vegetative understory and allow for longer retention times of irrigation water within the soil.¹⁵⁵ This practice has an added benefit of providing renewable energy for the operation (see [Strategy 7](#) for more information).



Agrivoltaics can provide dual benefits of shade for crops and on-farm renewable energy generation, offering opportunities for climate change adaptation and mitigation.

Approach 4.2: Manage for Changing Seasonality

Crop phenology, like other farm management decisions, is governed by the timing of the seasons. However, climate change threatens to alter the seasonality on which California's agricultural clock has been based. This Approach emphasizes changing springtime temperatures, though fall temperature changes have been observed² and projected,¹⁵⁶ and the implications of fall season changes are noted in Strategies [2](#), [3](#), and [5](#) while the changing seasonality of precipitation is addressed in Strategy [2](#).

Since the end of the 20th century, spring temperatures, as indicated by the date at which snowmelt is initiated, have arrived on average three weeks earlier than historic records.¹⁵⁷ Earlier and warmer springs have implications for water resources (see [Strategy 2](#)) and can result in earlier budbreak and bloom, which can increase the risk of frost damage¹⁵⁸ and may cause a mismatch between plant growth and pollinator activity.¹⁵⁹ Likewise, warming spring temperatures can compress the pollination windows in crops like almonds¹⁶⁰ and processing tomatoes,¹⁴⁶ and may decrease harvested fruit size in peaches.¹⁶¹ Although producers may see some benefits from warming spring temperatures in the long term, the potential negative outcomes seem to require multiple avenues of adaptation in order to improve the resilience of existing crop systems. Adaptive actions to changing seasonality may involve water resources and pest management (see Strategies [2](#) and [3](#), respectively), as well as management practices to address warming spring temperatures such as those in the following examples.

Examples of adaptation practices include:

- **Practice 4.2.1: Manage cover crops** to support and diversify wild pollinator populations.^{162,163}
- **Practice 4.2.2: Invest in infrastructure such as wind machines and sprinklers** to protect specialty crops from frost.¹⁵³ Consider that frost risk will ultimately decline under climate change projections¹³¹ but in some areas it may remain a sufficient risk to warrant mitigation investments. Also consider local water use ordinances/policies and drought status when using sprinklers for non-irrigation activities.
- **Practice 4.2.3: Adjust transplant dates** depending on local in-season¹⁴⁶ and long-term¹⁶⁴ temperature projections.

Approach 4.3: Prepare for Longer-Term and Larger-Scale Temperature Changes

Temperature is a key driver of crop geography, and the temperature regime is a primary determinant of which crops would be most suitable for a particular area. Climate change projections also suggest that crops like carrots, tomatoes, and cantaloupes¹⁴⁷ may see declining suitability within their current cultivation ranges and northward shifts in their growing regions. This Approach focuses on practices that provide adaptive capacity for changing temperature regimes that are increasingly outside the bounds of current conditions. These practices include crop diversification and, potentially, transition.

Diversification is a risk management method that reduces operational exposure to the particular vulnerabilities associated with any one cropping system. Additionally, agroecosystems thrive on diversification, especially given changing climate regimes. By diversifying, producers can improve their resilience and mitigate the risks associated with crop failure, pest outbreaks, and pathogen transmission.¹⁶⁵ Transition on agricultural lands may fall on a spectrum from shifting varieties of crops to moving toward non-agricultural land uses. Projected warming may provide opportunities for incorporating novel crops or varieties where they were previously less suitable, providing a strategy for both improved resilience and longer-term adaptation.^{166,167} Producers may begin with diversifying crop varieties or species as a means to incrementally transition to new crops that are better matched with the projected climate.

Examples of adaptation practices include:

- **Practice 4.3.1: Incorporate intercropping into annual systems** to achieve diversification benefits without compromising market value.^{168,169}
- **Practice 4.3.2: Consider adopting products or markets that can utilize imperfect produce.**⁹⁶ For example, creating value-added products such as juices or jams provides operational diversification.
- **Practice 4.3.3: Transition to lower-chill cultivars** for fruit and nut crops impacted by reduced chill hours, as available.¹⁷⁰
- **Practice 4.3.4: Transition to more heat-tolerant varieties** via repositories, genetic modification, or selective breeding.¹⁶⁵
- **Practice 4.3.5: Transition to or incorporate mixed or alternative land uses** (e.g., conservation or energy production) to better align land use with climate conditions. See [Strategy 6](#) for more information.

Strategy 5: Enhance Preparedness for Extreme Events

Extreme climate events can cause not only immense short-term damage but also long-term change,¹⁷¹ and they can cause more severe impacts than non-extreme events. Climate change will increase the frequency, duration, and/or intensity of extreme events in California, including extreme heat,⁷ extreme precipitation,^{4,6} extreme wind,¹⁷² and extreme wildfire events.¹⁵⁶ These events can have significant impacts on specialty crop production in California. For example, extreme heat can shift flowering times in perennials and create inhospitable growing conditions for many species,¹⁷³ and extreme precipitation can cause flooding, resulting in waterlogged soil and crop damage.¹⁷⁴ While producers may not be able to prevent these events outright, a number of approaches and practices can enhance preparedness and protect their operation.

Approach 5.1: Prepare for Extreme Heat Events

Extreme heat events can be a challenge for California agriculture, with impacts ranging from increased water demands and crop damage to loss of electrical power.² California's extreme heat events, which can be classified as either dry daytime or humid nighttime events,¹⁷⁵ increased at a rate of seven extreme heat days per year and 21 extreme heat nights per year over the 1987–2016 period.¹⁰³ However, climate change is projected to increase the frequency, intensity, and duration of extreme heat events across California, with regional variation in the degree of increase.^{175,176}

Extreme heat events can have varying impacts depending on the cropping system, the tolerance of a given crop to heat stress, and the timing of the heat event. Crops tend to experience heat stress during the warm season, but anomalously warm spring temperatures can also harm crops like almonds, cherries, grapes, and pistachios.⁷ Without proper preparation, extreme heat events can have deleterious consequences on specialty crop production including early flower drop and reduced pollen viability,¹⁷⁷ and decreased size, quality, and yield for multiple fruit and nut crops.⁷ Similar adaptive management practices are applicable across cropping systems.

Examples of adaptation practices include:

- **Practice 5.1.1: Update irrigation scheduling in anticipation of a heat event.** (See [Strategy 2](#) for more information on irrigation technology and scheduling.)
- **Practice 5.1.2: Add shade and/or shelter to modify the microclimate** with both natural (e.g., agroforestry) and/or man-made (e.g., tarps) methods.¹⁷⁸
- **Practice 5.1.3: Establish soil cover using crop residues in orchard systems** to insulate the soil and roots from extreme heat.¹⁷⁸
- **Practice 5.1.4: Shift to earlier planting dates to avoid summer heat stress.**¹⁷⁹ (See [Approach 4.1](#) for more on shifting harvest dates.)
- **Practice 5.1.5: Switch to heat-tolerant varieties.**



Shade netting can provide sun protection for crops sensitive to heat. Netting may also provide protection from birds and hail.

Approach 5.2: Manage for Extreme Precipitation and Flooding

Extreme precipitation in California is often produced by atmospheric rivers (ARs), which are long, narrow streams in the atmosphere that carry large amounts of water vapor and can produce heavy rainfall.¹⁸⁰ In fact, almost all of California's major historical floods can be attributed to ARs.¹⁰⁴ Observations of daily precipitation across the Western U.S. show increased precipitation intensity,¹⁰⁵ and the occurrences of extreme precipitation events are projected to double in the future compared to their historical frequency in California,⁸⁰ accompanied by a decreased frequency of lower-intensity precipitation events. In addition, because climate change is increasing not only the intensity of extreme precipitation events¹⁸¹ but also the duration and frequency of drought,^{91,92} extreme precipitation events can occur after extreme dry periods, compounding the impacts to agricultural lands by exacerbating soil erosion and flooding.¹⁸²

Agricultural impacts of extreme precipitation events can include crop damages and reduced crop quality,¹⁸³ increased soil erosion,¹⁸⁴ pollution from nutrient runoff, and pest infestation.^{2,185} Additionally, given the increased flood risk, growers may consider adopting on-farm management practices to mitigate erosion and the potential damages from waterlogging and root rot. Additional adaptation practices applicable to extreme precipitation and flood management are provided in Approach 2.3.

Examples of adaptation practices include:

- **Practice 5.2.1: Incorporate tailwater pond(s) to capture runoff.**¹⁸⁶ Tailwater ponds can be particularly effective at reducing the sediment loss and pollution of neighboring downstream waterways.
- **Practice 5.2.2: Include cover crops in planting rotations to increase soil organic matter** and thus reduce erosion from extreme precipitation, encourage water absorption, and reduce soil crusting.¹⁸⁷
- **Practice 5.2.3: Shift to flood-tolerant breeds and/or crop types.**^{188,96} Flood-tolerant crops can also allow for the adoption of AgMAR practices during flood events to capture excess rainfall and adapt to changing precipitation patterns (see [Approach 2.3](#)).

Approach 5.3: Enhance Resilience to Extreme Wind

California's coastline is prone to strong and gusty winds due in part to the state's unique coastal topography and regional climate conditions, but the spatial and temporal distribution of the state's wind speed is expected to change with a warming climate.¹⁷² While wind is a part of the climate in these coastal regions, an uptick in non-Santa Ana wind strength has been observed in the past 60 years.¹⁸⁹ Likewise, extreme Santa Ana events have also increased in intensity over recent decades.¹⁹⁰ Looking forward, summertime winds throughout the state are projected to increase in both frequency and strength, particularly in those coastal regions;¹⁹¹ however, Santa Ana winds may decrease in frequency, if not intensity.¹⁹⁰ Current and future extreme wind events may impact agricultural systems in a number of ways. For example, extreme wind can cause or exacerbate erosion,¹⁹² damage orchard trees,¹⁹³ and increase ignition risk and the spread of wildfires.¹⁵⁶ To protect against wind damages, growers may consider adopting practices that prevent erosion and reduce soil loss.

Examples of adaptation practices include:

- **Practice 5.3.1: Incorporate wind breaks in the form of tall, live barriers** such as hedgerows, fruit trees, or natural vegetation to provide shelter from wind and reduce erosion.¹⁹⁴
- **Practice 5.3.2: Plant short-season crops or cover crops for high wind periods** to protect soil from wind.¹⁹⁵
- **Practice 5.3.3: Consider measures such as speed control or polymer application** to control wind erosion and dust from unpaved roads.¹⁹⁶

Approach 5.4: Prepare for and Respond to Wildland Fire and Smoke Impacts

Wildfires are a natural phenomenon in California and provide ecosystem services to parts of the state's forested land.¹⁹⁷ Multiple factors have led to an increase in wildfire frequency and severity in recent decades, with the state experiencing a fivefold increase in annual burned area from 1972–2018.¹⁹⁸ Climate models project that rising temperatures and related climatic factors will lead to a 36–74% increase in burned area by 2085,¹⁹⁹ and that extreme wildfire events (>10,000 hectares burned¹⁹⁹) and extreme fire weather¹⁵⁶ will occur more frequently in the future. In addition, other climate-driven changes like shifts toward a later-onset wet season²⁰⁰ and increasingly frequent hot and dry conditions^{5,83} will increase wildfire risk.

Both wildfire flames and smoke impact California agriculture in unique ways. Wildfires can burn crops, destroy power lines and other vital infrastructure, and cause health problems for farm workers. For crops like winegrapes, smoke exposure can negatively impact fruit quality by causing smoke taint.^{201,202} While there are limited field-level actions that can prevent wildfires, producers can make management changes to protect agricultural land and energy systems from the flames. This Approach emphasizes practices that prevent the spread of wildfire onto agricultural land and encourage energy security.

Examples of adaptation practices include:

- **Practice 5.4.1: Apply artificial grape cuticle to wine grapes one week before anticipated smoke exposure** to reduce the impact of smoke taint.²⁰³
- **Practice 5.4.2: Incorporate firebreaks into the agricultural landscape** to help protect land and crops from wildfire and reduce the possibility of fire spreading.²⁰⁴
- **Practice 5.4.3: Incorporate on-farm renewable energy into farm operations** to sustain production during power loss. See [Approach 7.2](#) for more information regarding on-farm renewable energy.
- **Practice 5.4.4: Transition to crops that are less impacted by smoke.**

Strategy 6: Manage Farms and Fields as Part of a Larger Landscape

Individual farms influence and are influenced by the larger surrounding landscape and ecosystems within which they reside. While climate change impacts both farm and landscape-level systems, with purposeful management decisions that support ecosystem services, biological diversity, and the conservation of natural and working lands, farms can increase the resilience of the broader landscape and ameliorate the effects of climate change.²⁰⁵ Encouraging management approaches and practices that focus on landscape-scale, agroecosystem-focused dynamics, such as the examples shown below, may help California's farms and fields adapt to climate change on a longer time horizon.

Approach 6.1: Integrate, Improve, or Maintain Natural Ecosystems within Agricultural Operations

The overall health and wellbeing of agricultural land is heavily influenced by the ecosystem of the surrounding landscape. As such, integrating, improving, and maintaining that ecosystem can help shield the lands that are in agricultural production from some of the negative effects associated with climate change.²⁰⁶ Efforts that promote landscape function, connectivity, and diversity can support field-level adaptive capacity. For example, integrating and maintaining natural systems on farm field boundaries can support soil health and water quality.²⁰⁷ Similarly, improving or maintaining vegetative strips, riparian buffers, or hedgerows can help to reduce water pollution²⁰⁸ and support biological diversity, and they may increase the potential for soil carbon sequestration²⁰⁹ (see [Approach 6.2](#)).

Examples of adaptation practices include:

- **Practice 6.1.1: Maintain or restore riparian areas, wetlands, bottomlands, and floodplains.**^{210,211}
- **Practice 6.1.2:** Plant a smaller portion of the land to **give natural ecosystem features such as streams the ability to move and adapt.**
- **Practice 6.1.3: Match landscapes and ecosystems with appropriate cropping patterns** to stay within the physical and productive potential of the land.²¹²
- **Practice 6.1.4: Consider cultural cool season prescribed burns** of native vegetation to reduce the risk of catastrophic fire.²¹³



Developing and/or maintaining pollinator habitat through incorporating natural or semi-natural landcover or increasing crop diversity can provide benefits for the farm and the broader ecosystem. Here, pollinator habitat is maintained alongside solar panels, offering opportunities to manage the farm as a part of the larger landscape and contribute to climate change mitigation efforts by generating renewable energy.

Approach 6.2: Promote Biological Diversity Across the Landscape

Ecosystems are expected to experience biodiversity loss in the coming decades as a consequence of climate change,²¹⁴ meaning that promoting and maintaining biological diversity will be crucial to maintaining ecosystem balance.¹⁸⁶ In agricultural systems, diversity is often limited at the field scale. As such, supporting biological diversity at the landscape scale and improving diversity along the bounds of land in production can serve to support broader ecosystem function. For example, diversity-promoting management practices can support habitat for beneficial insects and wildlife. Moreover, many of the practices that producers can employ to increase biological diversity on the landscape may have similar benefits as those promoted in other Approaches (e.g., reduced runoff and erosion, improved soil health).²¹⁵ Unlike Strategy 4, Approach 3, which refers to diversity within agricultural systems, this Approach underscores some of the benefits of biological diversity on the land surrounding agricultural systems and highlights some on-farm practices that can help to increase diversity at the landscape level.

Examples of adaptation practices include:

- **Practice 6.2.1: Create and maintain pollinator habitat for wild bees** within agroecosystems, such as with hedgerows or wildflowers (also see [Approach 3.1](#)), or on-field cover crops (also see [Approaches 1.1, 1.2, and 3.3](#)).
- **Practice 6.2.2: Promote diverse non-crop vegetation types** that can support arthropod predators and insect-eating birds and bats²¹⁶ (also see [Approach 3.1](#)).

- **Practice 6.2.3: Maintain wildlife corridors** that can allow wildlife populations to travel across multiple properties, impeded by roads and other human activity as little as possible.²¹⁷
- **Practice 6.2.4: Increase crop diversity in plantings when planting across multiple fields.** Diversifying crop types across a landscape may benefit the biological diversity of pollinators and birds as effectively as incorporating natural or semi-natural landcover.²¹⁸

Approach 6.3: Conserve Individual Farms

California's farms play an important role in adapting to and mitigating some of the effects of climate change.²¹⁹ Perennial agriculture, tap-rooted cover crops, and minimally-disturbed soils can all serve to sequester carbon, and California's irrigated agricultural land emits far less greenhouse gases (GHGs) per acre than urban lands.²⁰⁵ Farms can also provide additional services to the surrounding area, including limiting heat effects²²⁰ and providing opportunities for water supply improvement that may benefit the broader community. Although the growing population and associated land use changes consumed more than 1 million acres of the state's agricultural lands between 1984 and 2014,²²¹ by conserving farms as working landscapes, California can benefit from their unique role in combating climate change. Some steps that may preserve California farmland viability are noted here.

Examples of adaptation practices include:

- **Practice 6.3.1: Encumber property with a conservation easement**, possibly with an affirmative farming covenant.
- **Practice 6.3.2: Participate in the [Williamson Act](#)**. Participation allows landowners to voluntarily conserve farmland by restricting land conversion away from agricultural use during a defined contract period. This stabilizes the landscape in the short term (10-20 years) and provides tax benefits for landowners.
- **Practice 6.3.3: Create a farm succession plan** to ensure that the land will remain agricultural or working. Organizations such as [Farmlink](#) can help facilitate this action.

Strategy 7: Develop or Expand Co-Benefit Efforts to Mitigate Climate Change

California is integral to the national food system. As such, it is notable that since the United States signed the UN Framework on Climate Change in 1992, annual GHG emissions in the U.S. have decreased by 7% (1990-2020),²²² while California's emissions have declined by 3% (1990-2019), and the California agricultural sector has performed similarly (down 3% in 2000-2019).²²³ Many programs have been implemented to lower California agriculture's GHG footprint, with effects that can take years to become apparent, but there is still great potential for action.

Many climate adaptation practices have co-benefits, and those co-benefits can reach above the farm level. One such co-benefit is climate change mitigation, such as when carbon is sequestered in the soil or greenhouse gas emissions are reduced. Carbon dioxide (CO₂) and nitrous oxide (N₂O) are the principal greenhouse gases emitted by California specialty crop farm management practices. On-farm emissions and carbon sequestration can be measured in a variety of ways. For example, on-farm CO₂ emissions can be quantifiably reduced by improving the energy efficiency of farm equipment switching to electrical power, and/or sourcing electricity from utilities that are less dependent on fossil fuels. Carbon (C) sequestration can be quantified by widely-available soil analyses, measuring changes in organic carbon levels within the soil, although the challenges and timescale required should be understood ahead of time. Nitrous oxide emissions can be reduced by lowering nitrogen (N) fertilizer applications, or through changes in irrigation or fertigation, which have been studied extensively in California.²²⁴ The approaches and examples of practices listed below may be useful in sequestering carbon and mitigating other GHG emissions.

Approach 7.1: Reduce On-Farm Greenhouse Gas Emissions

Carbon dioxide is not generally emitted in large quantities by soils that have been under long-term cultivation in California. However, equipment such as tractors and diesel pumps are responsible for on-farm CO₂ emissions and also make up a large part of California agriculture's GHG footprint.²²⁵ Also, it is estimated that in California specialty crop production, on-farm nitrous oxide emissions contribute even more than CO₂ to total on-farm emissions.²²³ Nitrous oxide is primarily emitted under conditions of excess nitrogen, high moisture, and adequate temperatures in the soil.²²⁶ Management of nitrogen-bearing fertilizers is the key to reducing N₂O emissions. Fortunately, nitrogen management goals such as reducing cost, improving N productivity,²²⁷ and improving soil health,²²⁸ generally help to lower N₂O, so that the reduced emissions are a co-benefit.

Examples of mitigation practices include:

- **Practice 7.1.1: Pursue "incremental progress"** such as energy audits, variable frequency drives on electric pumps, upgrades of diesel equipment to engines with the best available emissions technology, etc.
- **Practice 7.1.2: Invest in electric-powered equipment** such as electric irrigation pumps or electric tractors (SWEEP, FARMER, and Carl Moyer funds may assist California growers).
- **Practice 7.1.3: Consider using rooftops and residual areas for solar or wind power generation.** Mixed crop-and-solar-panel "agrivoltaic" systems now being developed may optimize land use overall,²²⁹ especially where water supplies are lower than demand.¹⁵⁵
- **Practice 7.1.4: Reduce soil nitrous oxide emissions** through efficient and timely N management (e.g., fertigation,²²⁴ controlled release N fertilizers,²³⁰ and/or urease or nitrification inhibitors²³¹).



Moving to electric farm equipment such as electric tractors is one way to reduce on-farm greenhouse gas emissions, thereby expanding on-farm efforts to mitigate climate change.

Approach 7.2: Increase On-Farm Soil Carbon Sequestration

Since 1970, atmospheric carbon has increased from 325 parts per million (ppm) to 412 ppm,²³² and that increase is the main driver of the climatic shifts discussed in earlier sections of this workbook. Research suggests that California farms can provide opportunities to capture carbon and store it in the soil as organic matter.^{233,234} Because agricultural soils typically have 25–75% less organic carbon than their undisturbed counterparts,²³⁵ carbon capture into cultivated soils provides "regenerative" soil health benefits (see also [Strategy 1](#)). Moreover, capturing carbon from the atmosphere into the soil supports all of the Strategies in this workbook, ultimately serving to increase resilience to climate change, input efficiency, and food security.²³⁶

On-farm soil carbon sequestration can occur in a number of ways. It is limited by soil type and climate as well as whether the land is irrigated, but will also depend on three principal management decisions. First is the quantity of

Reducing On-Farm Greenhouse Gas Emissions at Jackson Family Wines

Like any other industry, agricultural operations emit carbon dioxide into the atmosphere. Reducing on-farm greenhouse gas (GHG) emissions is one way to decrease an operation's carbon footprint. [Jackson Family Wines](#) is an excellent example of an operation that is actively reducing their GHG emissions by making [incremental changes](#).

As a family-owned company, multigenerational sustainability is integral to the Jackson Family Wines mission. After seeing the effects of climate disasters on their operations, the company set a goal of [cutting their carbon footprint in half by 2030 and becoming climate positive by 2050](#) without purchasing carbon offsets.

Using their 2015 emissions as a baseline, Jackson Family Wines calculated that they need to reduce their carbon footprint by 4% each year in order to reach their goals. To achieve this, they are focusing on efforts such as [transitioning to electric vehicles, reinvesting renewable energy](#), shifting to lighter weight glass wine bottles, and going "zero-waste."

Jackson Family Wines is pursuing these changes to act as a leader in their industry and show that companies can do a lot to be agents of change.

[Read more about what Jackson Family Wines is doing to reduce their GHG emissions here!](#)

organic amendments, including crop residues. Second is the quality of those amendments. Long-lasting, "resistant" soil organic matter is largely made up of deceased microorganisms, which have a carbon-to-nitrogen ratio of 10:1, so organic amendments ideally have a C:N ratio of 24:1 or lower (understanding that a good deal of carbon will be used for microbial energy, while most nitrogen will go into microbial or plant biomass). Some studies indicate that large additions of organic amendments above a 24:1 ratio will cause nitrogen hunger for the crop and will fail to produce resistant soil organic matter unless accompanied by a nitrogen supplement,^{237,238,239} but the field results for soil carbon have been mixed. Thirdly, the conflict between reduced tillage systems with residue retention, and cover-cropping or amendment practices requiring more tillage, must be assessed for each case. Tillage causes losses of organic matter. With additional tillage, seasonal cover cropping will still improve soil health, but in Californian conditions it has often failed to measurably increase soil carbon.^{28,32,233}

Examples of mitigation practices include:

- **Practice 7.2.1: Implement the Practices from Approaches 1.1 and 1.2.** The practices likely to produce the greatest soil carbon increases are the retention of residues and elimination of tillage. With organic amendments, the effects depend on type of amendment, amount, soil type, and tillage.
- **Practice 7.2.2: Plan ahead for soil carbon sampling, using established protocols.** [CDFA's Healthy Soils Program](#) includes recommendations for when, where, and how to prepare. See Resources for more information.
- **Practice 7.2.3: Consider using biochar for future use.** Biochar has soil benefits beyond carbon sequestration, but as a newer technology, various forms of biochar are still being studied, and how long it will remain in the soil and how durable its C sequestration benefits may be are still not known.²⁴⁰
- **Practice 7.2.4: Plant woody perennials or cover crops with taproots** for deeper soil C deposition with higher retention.^{241,242} With perennials, carbon will also be stored in the plant for as long as it stands.
- **Practice 7.2.5: Monitor soil N and consider N supplementation for high-carbon organic amendments.** Both crop health and resistant soil carbon buildup are the aims of this Practice.

RESOURCES

Many web-based resources are linked above in this workbook. When using a print version of the workbook, the website addresses (URLs) below correspond to the sites linked in the text above. Should you wish to visit these websites, they are organized by workbook Section and are listed in the order in which they appear.

A Guide for Adaptation Planning

Resource Name	Resource Owner	Resource Type and Description	URL
Climate Risks to California Agriculture	USDA California Climate Hub	This fact sheet provides a 30,000-foot view of some climate risks affecting California Agriculture and explains some ways that the California Climate Hub can assist producers impacted by these changes.	https://www.climatehubs.usda.gov/sites/default/files/Climate%20Risks%20for%20Ag%20FS%20050220.pdf
California Crops Under Climate Change	USDA California Climate Hub	This story map provides a broad overview of California as "America's Fruit and Nut Basket", the challenges specialty crop production will face, and the potential climate adaptive opportunities that can support California agriculture.	https://www.climatehubs.usda.gov/hubs/california/california-crops-under-climate-change
Resist-Accept-Direct (RAD) – A Framework for the 21 st -century Natural Resource Manager	National Parks Service, U.S. Department of the Interior	This white paper breaks down the Resist-Accept-Direct (RAD) decision framework in the context of natural resource management, and explains how it can be used by natural resource managers collaborating at larger scales across jurisdictions.	https://irma.nps.gov/DataStore/Download-File/654543
R–R–T (resistance–resilience–transformation) typology reveals differential conservation approaches across ecosystems and time	Communications Biology	This peer reviewed paper explains the proposed 6-point continuous interval scale representing different stages of action (or in-action) from active resistance to accelerated transformation.	https://www.nature.com/articles/s42003-020-01556-2
Climate Change Response Framework and Adaptation Workbook	USDA Climate Hubs, USFS, American Forests, Northern Institute of Applied Climate Science	This landing page walks viewers through an explanation of what their Adaptation Workbook for Forests is, and provides a 'quick guide', training, and demonstrations.	https://forestadaptation.org/adapt/adaptation-workbook

Resources Mentioned in Step 1: Defining Goals and Objectives			
Resource Name	Resource Owner	Resource Type and Description	URL
Managing Salts by Leaching	University of California, Agriculture and Natural Resources	This fact sheet provides information on how growers can manage various types of crops impacted by salinity through leaching.	https://anrcatalog.ucanr.edu/pdf/8550.pdf
Resources Mentioned in Step 2: Assessing Climate Impacts and Vulnerabilities			
Resource Name	Resource Owner	Resource Type and Description	URL
California's Fourth Climate Change Assessment	California Governor's Office of Planning and Research	This statewide climate report provides information for building resilience to climate impacts throughout the state.	https://climateassessment.ca.gov/
Resilient California	California's Governor's Office of Planning and Research	This clearinghouse site includes several tools, data, scientific studies, and more to help Californians adapt to climate change.	https://resilientca.org/tools/
Vulnerability and Adaptation to Climate Change in California Agriculture	California Water Library, California Energy Commission, UC Davis	This white paper/vulnerability assessment lays out California's agricultural vulnerability, taking into account 22 climate, crop, land use, and socioeconomic variables.	https://cawaterlibrary.net/document/vulnerability-and-adaptation-to-climate-change-in-california-agriculture/
CalAgroClimate	University of California Agriculture and Natural Resources and the USDA California Climate Hub	This decision support tools site hosts interactive web tools and related resources to help California farmers make on-farm management decisions.	https://calagroclimate.org/
Cal-Adapt	California Energy Commission, State of California	This decision support tools site includes interactive web tools for visualizing past and projected climate metrics across California.	https://cal-adapt.org/
The Climate Toolbox	University of California, Merced	This clearinghouse site includes a collection of tools and resources for visualizing past and projected climate and hydrology of the contiguous United States.	https://climatetoolbox.org/
U.S. Climate Resilience Toolkit	U.S. Global Change Research Program, U.S. National Oceanic and Atmospheric Administration	This clearinghouse site includes a collection of tools, case studies, and additional information and resources to improve resilience through improving understanding of climate-related risks and opportunities.	https://toolkit.climate.gov/

*Refer to the quick start guides and/or tutorials for the web tools above for more information on how they can be used to assess the climate conditions on your farm.

A Menu of Adaptation Strategies and Approaches for California Specialty Crops

Provided below are more resources that can aid in your adaptation planning, in addition to those linked in the document above. These resources are organized by the menu Strategy and Approach that they principally correspond to, although some resources can be used for multiple Strategies, Approaches, and Practices. For those referencing a print version of this workbook, complete web addresses are also provided should you wish to visit these websites.

STRATEGY 1: Support and Maintain Soil Health, Soil Biological Services, and Water Quality

Approach 1.1: Maintain and Improve Soil Structure			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 340: Cover Crops	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and environmental considerations land managers may consider when implementing cover crops.	https://www.nrcs.usda.gov/resources/guides-and-instructions/cover-crop-ac-340-conservation-practice-standard
USDA NRCS Conservation Practice 327: Conservation Cover	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and environmental considerations land managers may consider when implementing conservation cover.	https://www.nrcs.usda.gov/resources/guides-and-instructions/conservation-cover-ac-327-conservation-practice-standard
USDA NRCS Plant Materials Technical Documents	USDA NRCS	This clearinghouse page provides various publications published by the Lockeford Plant Materials Center to help land managers implement conservation practices.	https://www.nrcs.usda.gov/plant-materials/publications/search

Approach 1.2: Enhance Soil Biology and Nutrient Availability with Organic Amendments			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Interim Conservation Practice 808: Soil Carbon Amendment. NOTE: When adopted as a permanent standard, expected in late 2022, this will become Practice 336.	USDA NRCS	This fact sheet is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when adding soil carbon amendments.	https://efotg.sc.egov.usda.gov/api/CPS-File/10954/808_PI_ICPS_Soil_Carbon_Amendment_2020
NRDC Composting 101	Natural Resources Defense Council	This web page explains the basics of composting, from the various types to the do's and don'ts of the practice.	https://www.nrdc.org/stories/composting-101#howto
CDFA On-Farm Compost Resources	California Department of Food and Agriculture	This web page provides a series of resources for on-farm composting, including regulatory compliance, best practices, nutrient management creation, and more.	https://www.cdfa.ca.gov/healthysoils/ofcwg.html
UC Davis Grower's Guide to Whole Orchard Recycling	UC Davis	This web page provides FAQs, cost considerations, information on incentives, and lists of the multiple benefits related to Whole Orchard Recycling (WOR).	https://orchardrecycling.ucdavis.edu/growers-guide-wor

Approach 1.3: Manage for Soil and Water Salinization			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 610: Saline and Sodic Soil Management	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and environmental considerations land managers may consider when managing for salts and sodium in soil.	https://www.nrcs.usda.gov/resources/guides-and-instructions/saline-and-sodic-soil-management-ac-610-conservation-practice
Salinity Management Practices	UC ANR	This web page provides a host of information on the benefits of proper salinity management practices and how to implement them.	https://ucanr.edu/sites/Salinity/Salinity_Management/Salinity_management_practices/

Approach 1.4: Manage Fertilizer Application to Protect Groundwater Quality and Soil Health			
Resource Name	Resource Owner	Resource Type and Description	URL
NRCS Conservation Practice 590: Nutrient Management	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and environmental considerations land managers may consider when implementing nutrient management practices.	https://www.nrcs.usda.gov/resources/guides-and-instructions/nutrient-management-ac-590-conservation-practice-standard

STRATEGY 2: Cope with Uncertain Water Availability

Approach 2.1: Improve or Alter Water Systems to Meet Current and Expected Future Demands			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 441: Irrigation System, Microirrigation	USDA NRCS	This fact sheet PDF is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for implementing microirrigation systems.	https://www.nrcs.usda.gov/sites/default/files/2022-11/IRRIGATION.pdf
USDA NRCS Conservation Practice 442: Sprinkler System	USDA NRCS	This fact sheet PDF is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for implementing sprinkler systems.	https://www.nrcs.usda.gov/wps/PA_NRCSConsumption/download?cid=nrcse-prd1822439&ext=pdf
USDA NRCS Conservation Practice 449: Irrigation Water Management	USDA NRCS	This fact sheet PDF is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for implementing irrigation water management practices.	https://www.nrcs.usda.gov/sites/default/files/2022-10/FY23_DIA%20163_Irrigation%20Water%20Management%20Design.pdf

Approach 2.2: Manage for the Effects of Drought			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 340: Cover Crops	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and environmental considerations land managers may consider when implementing cover crops.	https://www.nrcs.usda.gov/resources/guides-and-instructions/cover-crop-ac-340-conservation-practice-standard
Deficit Irrigation Practices	Food and Agriculture Organization (FAO)	This is a compilation of scholarly research presented by the Food and Agriculture Organization in 2002, aiming to provide information on the ways crops react to drought stress	http://large.stanford.edu/courses/2016/ph241/lsm2/docs/wr22e.pdf
Drought Resistant Crops and Varieties List	UC ANR	This is a substantial, but not exhaustive, list of crops and varieties that can be drought tolerant or drought resistant.	https://ucanr.edu/sites/scmg/files/183771.pdf

Approach 2.3: Prepare for Changing Patterns of Precipitation			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 412: Grassed Waterway	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and ideal conditions for adding grassed waterways to a managed landscape.	https://www.nrcs.usda.gov/resources/guides-and-instructions/grassed-waterway-ac-412-conservation-practice-standard
USDA NRCS Conservation Practice 447: Irrigation and Drainage Tailwater Recovery	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that explain the definition, purpose, and ideal conditions for implementing irrigation and drainage tailwater recovery.	https://www.nrcs.usda.gov/resources/guides-and-instructions/irrigation-and-drainage-tailwater-recovery-no-447-conservation
USDA NRCS Interim Conservation Practice 817: On-Farm Recharge	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and ideal conditions for implementing on-farm recharge.	https://efotg.sc.egov.usda.gov/api/CPS-File/30313/817_CA_ICPS_On_Farm_Recharge_2020

STRATEGY 3: Manage Biological Stressors

Approach 3.1: Reduce Pest Pressure			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 595: Pest Management Conservation Systems	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It hosts resources that describe the definition, purpose, and ideal conditions for implementing pest management conservation systems.	https://www.nrcs.usda.gov/resources/guides-and-instructions/pest-management-conservation-system-ac-595-conservation-practice
USDA NRCS Conservation Practice 420: Wildlife Habitat Planning	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for establishing wildlife habitat by planting herbaceous vegetation or shrubs.	https://www.nrcs.usda.gov/resources/guides-and-instructions/wildlife-habitat-planting-ac-420-conservation-practice-standard
USDA NRCS Conservation Practice 645: Upland Wildlife Habitat Management	USDA NRCS	This fact sheet PDF is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for providing and managing upland habitats and connectivity within the landscape for wildlife.	https://efotg.sc.egov.usda.gov/api/CPS-File/1281/645_MN_CPS_Upland_Wildlife_Habitat_Management_2016
USDA NRCS Conservation Practice 422: Hedgerow Planting	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for implementing hedgerows into one's landscape.	https://www.nrcs.usda.gov/resources/guides-and-instructions/hedgerow-planting-ft-422-conservation-practice-standard

Approach 3.3: Reduce Weed and Invasive Plant Pressure			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 328: Conservation Crop Rotation	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for implementing conservation crop rotation onto one's landscape.	https://www.nrcs.usda.gov/resources/guides-and-instructions/conservation-crop-rotation-ac-328-conservation-practice-standard
USDA NRCS Conservation Practice 315: Herbaceous Weed Treatment	USDA NRCS	This fact sheet PDF is part of a series of conservation practices highlighted by the USDA NRCS. It explains the definition, purpose, and ideal conditions for removing or controlling herbaceous weeds, including invasive, noxious, prohibited, or undesirable plants from one's landscape.	https://www.nrcs.usda.gov/sites/default/files/2022-10/CO%20Herbaceous%20weed%20treatment%20E315A.pdf
UC ANR Integrated Pest Management Guidelines	UC ANR	This webpage details information about managing pests, broken down by crop type and then specific crop varieties. Information includes the University of California's official guidelines for monitoring pests and using pesticides and non-pesticide alternatives for managing insect, mite, nematode, weed, and disease pests.	https://www2.ipm.ucanr.edu/agriculture/

STRATEGY 4: Prepare for Temperature Change

Approach 4.1: Adapt to Warmer Conditions			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 340: Cover Crops	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when implementing cover crops.	https://www.nrcs.usda.gov/resources/guides-and-instructions/cover-crop-ac-340-conservation-practice-standard
Chill Calculator	UC Davis Fruit & Nut Information Center	This virtual chill calculator tool shows both historic and projected chill hours per year at specific CIMIS Stations.	https://fruitsand-nuts.ucdavis.edu/chill-calculator

Approach 4.2: Manage for Changing Seasonality			
Resource Name	Resource Owner	Resource Type and Description	URL
Map of Projected Change	CalAdapt	This interactive webtool shows maps of projected changes in long term (30 years) annual averages of maximum temperature, minimum temperature, and precipitation in California.	https://cal-adapt.org/tools/maps-of-projected-change

Approach 4.3: Prepare for Longer-Term and Larger-Scale Temperature Changes			
Resource Name	Resource Owner	Resource Type and Description	URL
Guidelines for Intercropping	Sustainable Agriculture Research and Education (SARE)	This planning manual can be used as a guide or reference for growers integrating crop rotations on their fields.	https://www.sare.org/publications/crop-rotation-on-organic-farms/guidelines-for-intercropping/ (introduction to the manual) https://www.sare.org/wp-content/uploads/Crop-Rotation-on-Organic-Farms.pdf (manual)

STRATEGY 5: Enhance Preparedness for Extreme Events

Approach 5.1: Prepare for Extreme Heat Events			
Resource Name	Resource Owner	Resource Type and Description	URL
Irrigation Scheduling Tables	Washington State University	This webpage includes Excel templates that allow users to schedule irrigation based on crop water use and soil moisture monitoring.	http://irrigation.wsu.edu/Content/Resources/Irrigation-Schedule-Tables.php

Approach 5.2: Manage for Extreme Precipitation and Flooding			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 340: Cover Crops	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when implementing cover crops.	https://www.nrcs.usda.gov/resources/guides-and-instructions/cover-crop-ac-340-conservation-practice-standard
Flood Managed Aquifer Recharge Information Page	California Department of Water Resources (CDWR)	This government webpage explains what Flood-Managed-Aquifer Recharge is and its benefits, and provides additional outside resources for implementing the practice.	https://water.ca.gov/programs/all-programs/flood-mar

Approach 5.3: Enhance Resilience to Extreme Wind			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 589: Cross Wind Trap Strips	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when incorporating cross wind trap strips.	https://www.nrcs.usda.gov/resources/guides-and-instructions/cross-wind-trap-strips-ac-589c-conservation-practice-standard
USDA NRCS Conservation Practice 380: Windbreak Shelterbelt Establishment	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when establishing, enhancing, or renovating windbreaks (aka, shelterbelts).	https://www.nrcs.usda.gov/resources/guides-and-instructions/windbreakshelterbelt-establishment-ft-380-conservation-practice

Approach 5.4: Prepare for and Respond to Wildland Fire and Smoke Impacts			
Resource Name	Resource Owner	Resource Type and Description	URL
USDA NRCS Conservation Practice 394: Firebreaks	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when integrating firebreaks into their landscape.	https://www.nrcs.usda.gov/resources/guides-and-instructions/fire-break-ft-394-conservation-practice-standard
USDA NRCS Conservation Practice 383: Fuel Breaks	USDA NRCS	This website is part of a series of conservation practices highlighted by the USDA NRCS. It describes the definition, purpose, and environmental considerations land managers may consider when adding fuel breaks into their landscape.	https://www.nrcs.usda.gov/resources/guides-and-instructions/fuel-break-ac-383-conservation-practice-standard

REFERENCES

¹ Medellín-Azuara, J., Howitt, R. E., MacEwan, D. J., and Lund, J. R. (2011). Economic impacts of climate-related changes to California agriculture. *Climatic Change*, 109(1), 387–405.

² Pathak, T. B., Maskey, M. L., Dahlberg, J. A., Kearns, F., Bali, K. M., and Zaccaria, D. (2018). Climate change trends and impacts on California agriculture: a detailed review. *Agronomy*, 8(3), 25.

³ Kapnick, S., and Hall, A. (2010). Observed climate–snowpack relationships in California and their implications for the future. *Journal of Climate*, 23(13), 3446–3456.

⁴ Swain, D. L., Langenbrunner, B., Neelin, J. D., and Hall, A. (2018). Increasing precipitation volatility in twenty-first-century California. *Nature Climate Change*, 8(5), 427–433.

⁵ Diffenbaugh, N. S., Swain, D. L., and Touma, D. (2015). Anthropogenic warming has increased drought risk in California. *Proceedings of the National Academy of Sciences of the United States of America*, 112(13), 3931–3936.

⁶ Polade, S. D., Gershunov, A., Cayan, D. R., Dettinger, M. D., and Pierce, D. W. (2017). Precipitation in a warming world: Assessing projected hydro-climate changes in California and other Mediterranean climate regions. *Scientific Reports*, 7(1), 1–10.

⁷ Parker, L. E., McElrone, A. J., Ostoja, S. M., and Forrestel, E. J. (2020). Extreme heat effects on perennial crops and strategies for sustaining future production. *Plant Science*, 295, 110397.

⁸ Walsh, M. K., Backlund, P. W., Buja, L., DeGaetano, A., Melnick, R., Prokopy, L., Takle, E., Todey, D., and Ziska, L. (2020). Climate indicators for agriculture. USDA Report No. 1488-2022-961.

⁹ Luedeling, E., Zhang, M., and Girvetz, E. H. (2009). Climatic changes lead to declining winter chill for fruit and nut trees in California during 1950–2099. *PLoS One*, 4(7), e6166.

¹⁰ Parker, L. E., and Abatzoglou, J. T. (2018). Shifts in the thermal niche of almond under climate change. *Climatic Change*, 147(1), 211–224.

¹¹ Pathak, T. B., and Stoddard, C. S. (2018). Climate change effects on the processing tomato growing season in California using growing degree day model. *Modeling Earth Systems and Environment*, 4(2), 765–775.

¹² Trumble, J., and Butler, C. (2009). Climate change will exacerbate California's insect pest problems. *California Agriculture*, 63(2):73–78.

¹³ Lobell, D. B., Field, C. B., Cahill, K. N., and Bonfils, C. (2006). Impacts of future climate change on California perennial crop yields: Model projections with climate and crop uncertainties. *Agricultural and Forest Management*, 141(2–4), 208–218.

¹⁴ Schuurman, G. W., Hawkins Hoffman, C., Cole, D. N., Lawrence, D. J., Morton, J. M., Magness, D. R., Cravens, A. E., Covington, S., O'Malley, R. and Fisichelli, N. A. (2020). Resist–accept–direct (RAD)—a framework for the 21st-century natural resource manager. *Natural Resource Report NPS/NRSS/CCRP/NRR–2020/2213*. National Park Service, Fort Collins, Colorado. Available at: <https://doi.org/10.36967/nrr-2283597>.

¹⁵ Peterson St-Laurent, G., Oakes, L. E., Cross, M., and Hagerman, S. (2021). R–R–T (resistance–resilience–transformation) typology reveals differential conservation approaches across ecosystems and time. *Communications Biology*, 4(1), 1–9.

¹⁶ Swanson, C. W., Janowiak, M. K., Brandt, L. A., Butler, P. R., Handler, S. D., Shannon, P. D., Lewis, A. D., and Others. (2016). *Forest adaptation resources: climate change tools and approaches for land managers*. General Technical Report NRS-GTR-87-2. Newtown Square, PA: US Department of Agriculture, Forest Service, Northern Research Station. 161 p.

¹⁷ Janowiak, M. K., Dostie, D. N., Wilson, M. A., Kucera, M. J., Skinner, R. H., Hatfield, J. L., Hollinger, D., and Swanson, C. W. (2016). *Adaptation Resources for Agriculture: Responding to Climate Variability and Change in the Midwest and Northeast*. Technical Bulletin 1944. Washington, DC: United States Department of Agriculture.

¹⁸ IPCC. (2001) *Climate Change 2001: Impacts, Adaptation and Vulnerability*, WMO.

¹⁹ Jackson, L., Haden, V. R., Wheeler, S. M., Hollander, A. D., Perlman, J., O'Geen, T., Mehta, V. K., Clark, V., Williams, J., and Thrupp, A. (2012). *Vulnerability and Adaptation to Climate Change in California Agriculture*. California Energy Commission. Publication number: CEC-500-2012-031. Available at: <https://cawaterlibrary.net/wp-content/uploads/2017/06/CEC-500-2012-031.pdf>

²⁰ Comerford, N. B., Franzluebbers, A. J., Stromberger, M. E., Morris, L., Markewitz, D., and Moore, R. (2013). Assessment and evaluation of soil ecosystem services. *Soil Horizons*, 54(3), 1–14.

²¹ Paul, C., Kuhn, K., Steinhoff-Knopp, B., Weißhuhn, P., and Helming, K. (2021). Towards a standardization of soil related ecosystem service assessments. *European Journal of Soil Science*, 72(4), 1543–1558.

²² Todorovska, M. I., and Al Rjoub, Y. (2006). Effects of rainfall on soil–structure system frequency: Examples based on poroelasticity and a comparison with full-scale measurements. *Soil Dynamics and Earthquake Engineering*, 26(6), 708–717.

²³ Morrow, J. G., Huggins, D. R., and Reganold, J. P. (2017). Climate change predicted to negatively influence surface soil organic matter of dryland cropping systems in the inland Pacific Northwest, USA. *Frontiers in Ecology and Evolution*, 5, 10.

²⁴ Corwin, D. L. (2021). Climate change impacts on soil salinity in agricultural areas. *European Journal of Soil Science*, 72(2), 842–862.

²⁵ Calderón, F. J., and Jackson, L. E. (2002). Rototillage, disking, and subsequent irrigation: effects on soil nitrogen dynamics, microbial biomass, and carbon dioxide efflux. *Journal of Environmental Quality*, 31(3), 752–758.

²⁶ Kassam, A., Friedrich, T., Derpsch, R., Lahmar, R., Mrabet, R., Basch, G., González-Sánchez, E. J., and Serraj, R. (2012). Conservation agriculture in the dry Mediterranean climate. *Field Crops Research*, 132, 7–17.

²⁷ Canali, S., Campanelli, G., Ciaccia, C., Leteo, F., Testani, E., and Montemurro, F. (2013). Conservation tillage strategy based on the roller crimper technology for weed control in Mediterranean vegetable organic cropping systems. *European Journal of Agronomy*, 50, 11–18.

²⁸ Mitchell, J. P., Shrestha, A., Mathesius, K., and Scow, K. M. (2017). Cover cropping and no-tillage improve soil health in an arid irrigated cropping system in California's San Joaquin Valley, USA. *Soil and Tillage Research*, 165, 325–335.

²⁹ Morugán-Coronado, A., Linares, C., Gómez-López, M. D., Faz, Á., and Zornoza, R. (2020). The impact of intercropping, tillage and fertilizer type on soil and crop yield in fruit orchards under Mediterranean conditions: A meta-analysis of field studies. *Agricultural Systems*, 178, 102736.

³⁰ Chen, G., and Weil, R. R. (2010). Penetration of cover crop roots through compacted soils. *Plant and Soil*, 331(1-2), 31–43.

³¹ Carof, M., De Tourdonnet, S., Coquet, Y., Hallaire, V., and Roger-Estrade, J. (2007). Hydraulic conductivity and porosity under conventional and no-tillage and the effect of three species of cover crop in northern France. *Soil Use and Management*, 23(3), 230–237.

³² Wolff, M. W., Alsina, M. M., Stockert, C. M., Khalsa, S. D. S., and Smart, D. R. (2018). Minimum tillage of a cover crop lowers net GWP and sequesters soil carbon in a California vineyard. *Soil and Tillage Research*, 175, 244–254.

³³ Kim, M. J., Shim, C. K., Kim, Y. K., Hong, S. J., Park, J. H., Han, E. J., Kim, J. H., and Kim, S. C. (2015). Effect of aerated compost tea on the growth promotion of lettuce, soybean, and sweet corn in organic cultivation. *The Plant Pathology Journal / the Korean Society of Plant Pathology*, 31(3), 259–268.

³⁴ Canellas, L. P., Olivares, F. L., Okorokova-Façanha, A. L., and Façanha, A. R. (2002). Humic acids isolated from earthworm compost enhance root elongation, lateral root emergence, and plasma membrane H⁺-ATPase activity in maize roots. *Plant Physiology*, 130(4), 1951–1957.

³⁵ Van Geel, M., De Beenhouwer, M., Lievens, B., and Honnay, O. (2016). Crop-specific and single-species mycorrhizal inoculation is the best approach to improve crop growth in controlled environments. *Agronomy for Sustainable Development*, 36(2), 37.

³⁶ Mitchell, J. P., Shennan, C., Singer, M. J., Peters, D. W., Miller, R. O., Prichard, T., Grattan, S. R., Rhoades, J. D., May, D. M., and Munk, D. S. (2000). Impacts of gypsum and winter cover crops on soil physical properties and crop productivity when irrigated with saline water. *Agricultural Water Management*, 45(1), 55–71.

³⁷ Doran, J. W., and Zeiss, M. R. (2000). Soil health and sustainability: managing the biotic component of soil quality. *Applied Soil Ecology: A Section of Agriculture, Ecosystems and Environment*, 15(1), 3–11.

³⁸ Pritchard, S. G. (2011). Soil organisms and global climate change. *Plant Pathology*, 60(1), 82–99.

³⁹ Weil, R., and Magdoff, F. (2004). Significance of soil organic matter to soil quality and health. In F. Magdoff, and R. R. Weil (eds.) *Soil Organic Matter in Sustainable Agriculture*. Boca Raton, FL: CRC Press, pp. 1–44.

⁴⁰ Allen, D. E., Singh, B. P., and Dalal, R. C. (2011). Soil health indicators under climate change: a review of current knowledge. In B. P. Singh, A. L. Cowie, and K. Y. Chan (eds.) *Soil Health and Climate Change*. Berlin/Heidelberg, Germany: Springer, pp. 25–45.

⁴¹ Morrow, J. G., Huggins, D. R., and Reganold, J. P. (2017). Climate change predicted to negatively influence surface soil organic matter of dryland cropping systems in the inland Pacific Northwest, USA. *Frontiers in Ecology and Evolution*, 5, 10.

⁴² Jones, C., McConnell, C., Coleman, K., Cox, P., Falloon, P., Jenkinson, D., and Powlson, D. (2005). Global climate change and soil carbon stocks: predictions from two contrasting models for the turnover of organic carbon in soil. *Global Change Biology*, 11(1), 154–166.

⁴³ van Zwieten, L. (2018). The long-term role of organic amendments in addressing soil constraints to production. *Nutrient Cycling in Agroecosystems*, 111(2), 99–102.

⁴⁴ Wall, D. H., Nielsen, U. N., and Six, J. (2015). Soil biodiversity and human health. *Nature*, 528(7580), 69–76.

⁴⁵ Lucas, S. T., D'Angelo, E. M., and Williams, M. A. (2014). Improving soil structure by promoting fungal abundance with organic soil amendments. *Applied Soil Ecology: A Section of Agriculture, Ecosystems and Environment*, 75, 13–23.

⁴⁶ White, K. E., Brennan, E. B., Cavigelli, M. A., and Smith, R. F. (2020). Winter cover crops increase readily decomposable soil carbon, but compost drives total soil carbon during eight years of intensive, organic vegetable production in California. *PLoS One*, 15(2), e0228677.

⁴⁷ Nielsen, G. H., Hogue, E. J., Forge, T., and Nielsen, D. (2003). Surface application of mulches and biosolids affect orchard soil properties after 7 years. *Canadian Journal of Soil Science*, 83(1), 131–137.

⁴⁸ TerAvest, D., Smith, J. L., Carpenter-Boggs, L., Granatstein, D., Hoagland, L., and Reganold, J. P. (2011). Soil carbon pools, nitrogen supply, and tree performance under several groundcovers and compost rates in a newly planted apple orchard. *HortScience: A Publication of the American Society for Horticultural Science*, 46(12), 1687–1694.

⁴⁹ Gale, E. S., Sullivan, D. M., Cogger, C. G., Bary, A. I., Hemphill, D. D., and Myhre, E. A. (2006). Estimating plant available nitrogen release from manures, composts, and specialty products. *Journal of Environmental Quality*, 35(6), 2321–2332.

⁵⁰ Jahanzad, E., Holtz, B. A., Zuber, C. A., Doll, D., Brewer, K. M., Hogan, S., and Gaudin, A. C. M. (2020). Orchard recycling improves climate change adaptation and mitigation potential of almond production systems. *PLoS One*, 15(3), e0229588.

⁵¹ Holtz, B., and Culumber, M. (2019). 2019 Nitrogen Considerations. *West Coast Nut* 2019(Feb), 14–19. https://www.wcngg.com/wp-content/uploads/2019/02/WCN_Feb19.pdf

⁵² Jeffery, S., Abalos, D., Spokas, K. A., and Verheijen, F. G. A. (2015). Biochar effects on crop yield. *Biochar for Environmental Management: Science, Technology and Implementation*, 2, 301–325.

⁵³ Shackelford, G. E., Kelsey, R., and Dicks, L. V. (2019). Effects of cover crops on multiple ecosystem services: Ten meta-analyses of data from arable farmland in California and the Mediterranean. *Land Use Policy*, 88, 104204.

⁵⁴ Pang, X. P., and Letey, J. (1998). Development and evaluation of ENVIRO-GRO, an integrated water, salinity, and nitrogen model. *Soil Science Society of America Journal*, 62(5), 1418–1427.

⁵⁵ University of California Agricultural Issues Center (Ed.). (2009). *The Measure of California Agriculture*. Oakland, CA: Regents of the University of California. Available at: <https://aic.ucdavis.edu/publications/moca/moca09.pdf> (Accessed: 1 August 2022).

⁵⁶ Bañuelos, G. S. (2015). Coping with naturally-high levels of soil salinity and boron in the westside of central California. *Perspectives in Science*, 3(1), 4–6.

⁵⁷ Zribi, L., Fatma, G., Fatma, R., Salwa, R., Hassan, N., and Néjib, R. M. (2009). Application of chlorophyll fluorescence for the diagnosis of salt stress in tomato *Solanum lycopersicum* (variety Rio Grande). *Scientia Horticulturae*, 120(3), 367–372.

⁵⁸ Machado, R. M. A., and Serralheiro, R. P. (2017). Soil salinity: effect on vegetable crop growth. management practices to prevent and mitigate soil salinization. *Horticulturae*, 3(2), 30.

⁵⁹ Loáiciga, H. A., Pingel, T. J., and Garcia, E. S. (2009). *Assessment of Seawater Intrusion Potential from Sea-level Rise in Coastal Aquifers of California*. Berkeley, CA: University of California Berkeley, University of California Water Resources Center. Available at: <https://escholarship.org/uc/item/38w0p9st>.

⁶⁰ Beede, R. H., Brown, H. P., and Kallsen, C. (2011). *Understanding and Correcting Nutrients in Nut Crops*. Davis, California: University of California Cooperative Extension.

⁶¹ Gradziel, T. M. (Ed.). (2017). *Almonds: Botany, Production and Uses*. Wallingford, England: CABI.

⁶² Siddiqui, M. I., and Brown, P. (2013). Pistachio early-season sampling and in-season nitrogen application maximizes productivity, minimizes loss. Protocol for early-season sampling and in-season nitrogen budgeting. Davis, CA: University of California, Division of Agriculture and Natural Resources. Available at: <http://fruitsandnuts.ucdavis.edu/files/165545.pdf>.

⁶³ O'Geen, A. (2015). *Drought Tip: Reclaiming Saline, Sodic, and Saline-Sodic Soils*. UCANR Publication 8519. Davis, CA: University of California, Division of Agriculture and Natural Resources.

⁶⁴ Burt, C. M., and Isbell, B. (2005). Leaching of accumulated soil salinity under drip irrigation. *Transactions of the ASAE. American Society of Agricultural Engineers*, 48(6), 2115–2121.

⁶⁵ Friedman, S. P., and Gamlie, A. (2021). A modelling approach and module for salinity management in drip irrigation. *Biosystems Engineering*, 207, 141–161.

⁶⁶ Yang, T., Šimůnek, J., Mo, M., McCullough-Sanden, B., Shahrokhnia, H., Cherchian, S., and Wu, L. (2019). Assessing salinity leaching efficiency in three soils by the HYDRUS-1D and -2D simulations. *Soil and Tillage Research*, 194, 104342.

⁶⁷ Hanson, B., and May, D. (2011). *Drip Irrigation Salinity Management for Row Crops*. UCANR Publication 8447. Davis, CA: University of California, Division of Agriculture and Natural Resources.

⁶⁸ Machado, R. M. A., and Serralheiro, R. P. (2017). Soil salinity: effect on vegetable crop growth. management practices to prevent and mitigate soil salinization. *Horticulturae*, 3(2), 30.

⁶⁹ Lakhdar, A., Rabhi, M., Ghnaya, T., Montemurro, F., Jedidi, N., and Abdelly, C. (2009). Effectiveness of compost use in salt-affected soil. *Journal of Hazardous Materials*, 171(1–3), 29–37.

⁷⁰ Chappelle, C., Hanak, E., Harter, T. (2017). *Groundwater in California [Fact sheet]*. San Francisco, CA: Public Policy Institute of California Water Policy Center.

⁷¹ Christian-Smith, J., Cooley, H., and Gleick, P. H. (2012). Potential water savings associated with agricultural water efficiency improvements: a case study of California, USA. *Water Policy*, 14(2), 194–213. Available at: <https://iwaponline.com/wp/article-abstract/14/2/194/20007>.

⁷² Lockhard, K. M., King, A. M., and Harter, T. (2013). Identifying sources of groundwater nitrate contamination in a large alluvial groundwater basin with highly diversified intensive agricultural production. *Journal of Contaminant Hydrology*, 151, 140–154.

⁷³ Khan, S. A., Mulvaney, R. L., Ellsworth, T. R., Boast, C. W. 2007. The myth of nitrogen fertilization for soil carbon sequestration. *Journal of Environmental Quality* 36, 1821–1832.

⁷⁴ Ladha, J. K., Reddy, C. K., Padre, A. T., and van Kessel, C. (2011). Role of nitrogen fertilization in sustaining organic matter in cultivated soils. *Journal of Environmental Quality*, 40(6), 1756–1766.

⁷⁵ Ayers, R. S., and Westcot, D. W. (1985). *Water Quality for Agriculture*. FAO Irrigation and Drainage Paper 29 Rev. 1. Rome: Food and Agricultural Organization.

⁷⁶ Bauder, T. A., Waskom, R. M., Sutherland, P. L., Davis, J. G., Follett, R. H., and Soltanpour, P. N. (2011). Irrigation water quality criteria. Service in Action, Fact Sheet No. 0. 506. Fort Collins, CO: Colorado State University Extension. Available at: <https://itc.tamu.edu/files/2018/05/00506.pdf>.

⁷⁷ Hartz, T. K. (1994). A quick test procedure for soil nitrate–nitrogen. *Communications in Soil Science and Plant Analysis*, 25(5-6), 511–515.

⁷⁸ Bai, S. H., Reverchon, F., Xu, C.-Y., Xu, Z., Blumfield, T. J., Zhao, H., Van Zwieten, L., and Wallace, H. M. (2015). Wood biochar increases nitrogen retention in field settings mainly through abiotic processes. *Soil Biology and Biochemistry*, 90, 232–240.

⁷⁹ Azad, N., Behmanesh, J., Rezaverdinejad, V., Abbasi, F., and Navabian, M. (2018). Developing an optimization model in drip fertigation management to consider environmental issues and supply plant requirements. *Agricultural Water Management*, 208, 344–356.

⁸⁰ Berg, N., and Hall, A. (2015). Increased interannual precipitation extremes over California under climate change. *Journal of Climate*, 28(16), 6324–6334.

⁸¹ Morris, K. S., and Bucini, G. (2016). California's drought as opportunity: Redesigning U.S. agriculture for a changing climate. *Elementa* (Washington, D.C.), 4, 000142.

⁸² Johnson, R., and Cody, B. A. (2015). California agricultural production and irrigated water use. Available at: <https://nationalaglawcenter.org/wp-content/uploads/assets/crs/R44093.pdf>.

⁸³ McEvoy, D. J., Pierce, D. W., Kalansky, J. F., Cayan, D. R., and Abatzoglou, J. T. (2020). Projected changes in reference evapotranspiration in California and Nevada: Implications for drought and wildland fire danger. *Earth's Future*, 8(11), e2020EF001736.

⁸⁴ García, I. F., Díaz, J. R., Poyato, E. C., Montesinos, P., and Berbel, J. (2014). Effects of modernization and medium term perspectives on water and energy use in irrigation districts. *Agricultural Systems*, 131, 56–63.

⁸⁵ Stubbs, M. (2016). Irrigation in U.S. agriculture: On-farm technologies and best management practices. Available at: <http://nationalaglawcenter.org/wp-content/uploads/assets/crs/R44158.pdf>.

⁸⁶ Zhang, T., Lin, X., Rogers, D. H., and Lamm, F. R. (2015). Adaptation of irrigation infrastructure on irrigation demands under future drought in the United States. *Earth Interactions*, 19(7), 1–16.

⁸⁷ Joyce, B. A., Mehta, V. K., Purkey, D. R., Dale, L. L., and Hanemann, M. (2011). Modifying agricultural water management to adapt to climate change in California's central valley. *Climatic Change*, 109(1), 299–316.

⁸⁸ Frisvold, G. B., and Deva, S. (2013). Climate and choice of irrigation technology: implications for climate adaptation. *Journal of Natural Resources Policy Research*, 5(2-3), 107–127.

⁸⁹ Cahn, M. D., and Johnson, L. F. (2017). New approaches to irrigation scheduling of vegetables. *Horticulturae*, 3(2), 28.

⁹⁰ Richter, B. D., Brown, J. D., DiBenedetto, R., Gorsky, A., Keenan, E., Madray, C., Morris, M., Rowell, D., and Ryu, S. (2017). Opportunities for saving and reallocating agricultural water to alleviate water scarcity. *Water Policy*, 19(5), 886–907.

⁹¹ Cook, B. I., Ault, T. R., and Smerdon, J. E. (2015). Unprecedented 21st century drought risk in the American Southwest and Central Plains. *Science Advances*, 1(1), e1400082.

⁹² Mann, M. E., and Gleick, P. H. (2015). Climate change and California drought in the 21st century [Review of Climate change and California drought in the 21st century]. *Proceedings of the National Academy of Sciences of the United States of America*, 112(13), 3858–3859.

⁹³ Overpeck, J. T. (2013). The challenge of hot drought. *Nature*, 503(7476), 350–351.

⁹⁴ Benson, D. O., and Dirmeyer, P. A. (2021). Characterizing the relationship between temperature and soil moisture extremes and their role in the exacerbation of heat waves over the contiguous United States. *Journal of Climate*, 34(6), 2175–2187.

⁹⁵ Tortajada, C., Kastner, M. J., Buurman, J., and Biswas, A. K. (2017). The California drought: Coping responses and resilience building. *Environmental Science and Policy*, 78, 97–113.

⁹⁶ Ames, G. K., and Dufour, R. (2014). Climate change and perennial fruit and nut production: Investing in resilience in uncertain times. ATTRA Sustainable Agriculture National Center for Appropriate Technology. Available at: <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.670.5416&rep=rep1&type=pdf> Accessed: 10 May 2022.

⁹⁷ Fereres, E., and Soriano, M. A. (2007). Deficit irrigation for reducing agricultural water use. *Journal of Experimental Botany*, 58(2), 147–159.

⁹⁸ Fulton, A., Buchner, R., and Niederholzer, F. (2015). Drought strategies for California prune production. UCANR Publication 8520. Davis, CA: University of California, Division of Agriculture and Natural Resources.

⁹⁹ Sanden, B. (2007). Fall irrigation management in a drought year for almonds, pistachios, and citrus. Bakersfield, CA: University of California Cooperative Extension Kern Soil and Water. University of California Agriculture and Natural Resources. Available at: <https://ucanr.edu/sites/Kern22/files/98822.pdf> (Accessed: 29 July 2022).

¹⁰⁰ da Silva Dias, J. C. (2015). Plant breeding for harmony between modern agriculture production and the environment. *Agricultural Sciences*, 6(01), 87.

¹⁰¹ Heinitz, C. C., Fort, K., and Walker, M. A. (2014, July). Developing drought and salt resistant grape rootstocks. In Li, S., Archbold, D., and London, J. (eds.) *XI International Conference on Grapevine Breeding and Genetics. Acta Horticultae 1082*. Leuven, Belgium: International Society for Horticultural Science, pp. 305-312.

¹⁰² UC Drought Management. (2022). Irrigation Scheduling. Davis, CA: University of California, Division of Agriculture and Natural Resources. Available at: https://ucmanagedrought.ucdavis.edu/Agriculture/Irrigation_Scheduling/. (Accessed: 10 May 2022).

¹⁰³ Milanes, C., Kadir, T., Lock, B., Monserrat, L., Pham, N., and Randles, K. (2018). Indicators of Climate Change in California. Sacramento, CA: Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. Available at: <https://oehha.ca.gov/media/downloads/climate-change/report/2018ccindicatorsreportmay2018.pdf>. (Accessed: 10 May 2022).

¹⁰⁴ Dettinger, M. D., Ralph, F. M., Das, T., Neiman, P. J., and Cayan, D. R. (2011). Atmospheric rivers, floods and the water resources of California. *WATER*, 3(2), 445-478.

¹⁰⁵ Zhang, F., Biederman, J. A., Dannenberg, M. P., Yan, D., Reed, S. C., and Smith, W. K. (2021). Five decades of observed daily precipitation reveal longer and more variable drought events across much of the western United States. *Geophysical Research Letters*, 48(7), e2020GL092293.

¹⁰⁶ Gershunov, A., Shulgina, T., Clemesha, R. E. S., Guirguis, K., Pierce, D. W., Dettinger, M. D., Lavers, D. A., Cayan, D. R., Polade, S. D., Kalansky, J., and Ralph, F. M. (2019). Precipitation regime change in Western North America: The role of Atmospheric Rivers. *Scientific Reports*, 9(1), 9944.

¹⁰⁷ Mahoney, K., Scott, J. D., Alexander, M., McCrary, R., Hughes, M., Swales, D., and Bukovsky, M. (2021). Cool season precipitation projections for California and the Western United States in NA-CORDEX models. *Climate Dynamics*, 56(9), 3081-3102.

¹⁰⁸ Kistner, E., Kellner, O., Andresen, J., Todey, D., and Morton, L. W. (2018). Vulnerability of specialty crops to short-term climatic variability and adaptation strategies in the Midwestern USA. *Climatic Change*, 146(1), 145-158.

¹⁰⁹ O'Geen, A. T., Saal, M. B. B., Dahlke, H. E., Doll, D. A., Elkins, R. B., Fulton, A., Fogg, G. E., Harter, T., Hopmans, J. W., Ingels, C., Niederholzer, F. J., Solis, S. S., Verdegaaal, P. S., and Walkinshaw, M. (2015). Soil suitability index identifies potential areas for groundwater banking on agricultural lands. *California Agriculture*, 69(2), 75-84.

¹¹⁰ Dahlke, H. E., LaHue, G. T., Mautner, M. R., Murphy, N. P., Patterson, N. K., Waterhouse, H., Yang, F., and Foglia, L. (2018). Managed aquifer recharge as a tool to enhance sustainable groundwater management in California: examples from field and modeling studies. In Friesen, J., and Rodríguez-Sinobas, L. (eds.) *Advances in Chemical Pollution, Environmental Management and Protection*. Volume. 3. Amsterdam, Netherlands: Elsevier, pp. 215-275.

¹¹¹ Waterhouse, H., Bachand, S., Mountjoy, D., Choperena, J., Bachand, P., Dahlke, H., and Horwath, W. (2020). Agricultural managed aquifer recharge – water quality factors to consider. *California Agriculture*, 74(3), 144-154.

¹¹² Reyes, J. J., and Elias, E. (2019). Spatio-temporal variation of crop loss in the United States from 2001 to 2016. *Environmental Research Letters*, 14(7), 074017.

¹¹³ Pathak, T. B., Maskey, M. L., and Rijal, J. P. (2021). Impact of climate change on navel orangeworm, a major pest of tree nuts in California. *The Science of the Total Environment*, 755(1), 142657.

¹¹⁴ Delgado-Baquerizo, M., Guerra, C. A., Cano-Díaz, C., Egidi, E., Wang, J. T., Eisenhauer, N., Singh, B. K. and Maestre, F. T. (2020). The proportion of soil-borne pathogens increases with warming at the global scale. *Nature Climate Change*, 10(6), 550-554.

¹¹⁵ Korres, N. E., Norsworthy, J. K., Tehranchian, P., Gitsopoulos, T. K., Loka, D. A., Oosterhuis, D. M., Gealy, D. R., Moss, S. R., Burgos, N. R., Miller, M. R., and Palhano, M. (2016). Cultivars to face climate change effects on crops and weeds: a review. *Agronomy for Sustainable Development*, 36(1), 12.

¹¹⁶ Cammell, M. E., and Knight, J. D. (1992). Effects of climatic change on the population dynamics of crop pests. In Begon, M., Fitter, A. H., and A. Macfadyen, A. (Eds.) *Advances in Ecological Research*, Volume 22. Cambridge, MA: Academic Press, pp. 117-162.

¹¹⁷ Porter, J. H., Parry, M. L., and Carter, T. R. (1991). The potential effects of climatic change on agricultural insect pests. *Agricultural and Forest Meteorology*, 57(1), 221-240.

¹¹⁸ Skendžić, S., Zovko, M., Živković, I. P., Lešić, V., and Lemić, D. (2021). The impact of climate change on agricultural insect pests. *Insects*, 12(5), 440.

¹¹⁹ Trumble, J., and Butler, C. (2009). Climate change will exacerbate California's insect pest problems. *California Agriculture*, 63(2):73-78.

¹²⁰ Deutsch, C. A., Tewksbury, J. J., Tigchelaar, M., Battisti, D. S., Merrill, S. C., Huey, R. B., and Naylor, R. L. (2018). Increase in crop losses to insect pests in a warming climate. *Science*, 361(6405), 916-919.

¹²¹ Haviland, D. R., Symmes, E. J., Adaskaveg, J. E., Duncan, R. A., Roncoroni, J. A., Gubler, W. D., Hanson, B., Hembree, K. J., Holtz, B. A., Stapleton, J. J., and Others. (2020). *UC IPM Pest Management Guidelines: Almond*. UC ANR Publication 3431. Davis, CA: University of California, Division of Agriculture and Natural Resources.

¹²² Haviland, D. R., Bentley, W. J., Beede, R. H., Daane, K. M., Michailides, T. J., Westerdahl, B. B., and Hembree, K. J. (2015). *UC IPM Pest Management Guidelines: Pistachio*. UC ANR Publication 3461. Davis, CA: University of California, Division of Agriculture and Natural Resources.

¹²³ Grant, J. A., Symmes, E. J., Baldwin, R. A., Fichtner, E. J., Roncoroni, J. A., Westerdahl, B. B., Adaskaveg, J. E., Bostock, R. M., Browne, G. T., and Buchner, R. P. (2020). *UC IPM Pest Management Guidelines: Walnut*. UC ANR Publication 3471. Davis, CA: University of California, Division of Agriculture and Natural Resources.

¹²⁴ Hodson, A., and Lewis, E. (2016). Managing for soil health can suppress pests. *California Agriculture*, 70(3):137–141.

¹²⁵ Altieri, M. A., Ponti, L., and Nicholls, C. I. (2005). Manipulating vineyard biodiversity for improved insect pest management: case studies from northern California. *International Journal of Biodiversity Science and Management*, 1(4), 191–203.

¹²⁶ Morandin, L. A., Long, R. F., and Kremen, C. (2014). Hedgerows enhance beneficial insects on adjacent tomato fields in an intensive agricultural landscape. *Agriculture, Ecosystems and Environment*, 189, 164–170.

¹²⁷ Avila-Quezada, G. D., Esquivel, J. F., Silva-Rojas, H. V., Leyva-Mir, S. G., de Jesus Garcia-Avila, C., Noriega-Orozco, L., Rivas-Valencia, P., Ojeda-Barrios, D., and Castillo, A. M. (2018). Emerging plant diseases under a changing climate scenario: Threats to our global food supply. *Emirates Journal of Food and Agriculture*, 30(6), 443–450.

¹²⁸ Bebber, D. P., Ramotowski, M. A. T., and Gurr, S. J. (2013). Crop pests and pathogens move polewards in a warming world. *Nature Climate Change*, 3(11), 985–988.

¹²⁹ Chaloner, T. M., Gurr, S. J., and Bebber, D. P. (2021). Plant pathogen infection risk tracks global crop yields under climate change. *Nature Climate Change*, 11(8), 710–715.

¹³⁰ Daugherty, M. P., Cooper, M., Smith, R., Varela, L., and Almeida, R. (2019, December). Has climate contributed to a Pierce's disease resurgence in North Coast Vineyards? *Wine Business Monthly*. Galley December 2019. pp. 2–7. Available at: <https://faculty.ucr.edu/~mattd/assets/files/PD%20climate%20-%20PWV.pdf> (Accessed: 10 May 2022).

¹³¹ Parker, L., Pathak, T., and Ostoja, S. (2021). Climate change reduces frost exposure for high-value California orchard crops. *The Science of the Total Environment*, 762, 143971.

¹³² Kremen, C., and Miles, A. (2012). Ecosystem services in biologically diversified versus conventional farming systems: benefits, externalities, and trade-offs. *Ecology and Society*, 17(4), 40. Available at: <https://www.jstor.org/stable/26269237>.

¹³³ Priyanka, A. K. M., Varma, S., Kumar, V., and Sharma, R. S. (2020). Impact of climate change on plant diseases and management strategies: A review. *IJCS*, 8(2), 2968–2973.

¹³⁴ Thomas, C. S., Gubler, W. D., and Leavitt, G. (1994). Field testing of a powdery mildew disease forecast model on grapes in California. *Phytopathology* 84, 1070 (abstract).

¹³⁵ Weber E., Gubler, D., and Derr, A. (1996). Powdery mildew controlled with fewer fungicide applications. *Practical Winery and Vineyard*, January/February 1996.

¹³⁶ Clements, D. R., and Ditomaso, A. (2011). Climate change and weed adaptation: can evolution of invasive plants lead to greater range expansion than forecasted? *Weed Research*, 51(3), 227–240.

¹³⁷ Gianoli, E., and Molina-Montenegro, M. A. (2021). Evolution of physiological performance in invasive plants under climate change. *Evolution; International Journal of Organic Evolution*, 75(12), 3181–3190.

¹³⁸ Ruttledge, A., and Chauhan, B. S. (2020). Climate change and weeds of cropping systems. In Jabran, K., Florentine, S., and Chauhan, B. S. (eds.), *Crop Protection Under Changing Climate*. Berlin, Germany: Springer International Publishing, pp. 57–84.

¹³⁹ Ziska, L. H., and Bunce, J. A. (2007). Predicting the impact of changing CO₂ on crop yields: some thoughts on food. *New Phytologist*, 175(4), 607–618.

¹⁴⁰ Matzraf, M., Seiwert, B., Reemtsma, T., Rubin, B., and Peleg, Z. (2016). Climate change increases the risk of herbicide-resistant weeds due to enhanced detoxification. *Planta*, 244(6), 1217–1227.

¹⁴¹ Sarrantonio, M., and Gallandt, E. (2003). The role of cover crops in North American cropping systems. *Journal of Crop Production*, 8(1–2), 53–74.

¹⁴² Roncoroni, J. A., and Wright, S. D. (2017). Weeds. In UC Statewide IPM Program (Ed.), *Pest Management Guidelines for Agriculture: Nectarine* UCANR Publication 3451 Davis, CA: University of California, Division of Agriculture and Natural Resources, pp. 90–109.

¹⁴³ Visser, M. E., and Gienapp, P. (2019). Evolutionary and demographic consequences of phenological mismatches. *Nature Ecology and Evolution*, 3(6), 879–885.

¹⁴⁴ Pierce, D. W., Das, T., Cayan, D. R., Maurer, E. P., Miller, N. L., Bao, Y., Kanamitsu, M., Yoshimura, K., Snyder, M. A., Sloan, L. C., Franco, G., and Tyree, M. (2013). Probabilistic estimates of future changes in California temperature and precipitation using statistical and dynamical downscaling. *Climate Dynamics*, 40(3), 839–856.

¹⁴⁵ Luedeling, E., Zhang, M., and Girvetz, E. H. (2009). Climatic changes lead to declining winter chill for fruit and nut trees in California during 1950–2099. *PLoS One*, 4(7), e6166.

¹⁴⁶ Pathak, T. B., and Stoddard, C. S. (2018). Climate change effects on the processing tomato growing season in California using growing degree day model. *Modeling Earth Systems and Environment*, 4(2), 765–775.

¹⁴⁷ Marklein, A., Elias, E., Nico, P., and Steenwerth, K. (2020). Projected temperature increases may require shifts in the growing season of cool-season crops and the growing locations of warm-season crops. *The Science of the Total Environment*, 746, 140918.

¹⁴⁸ Doll, D., Coelho, J., Zwieniecki, M., Zhang, L., Ferguson, L., and Culumber, M. (2017). The evaluation of winter kaolin clay and dormant oil applications on chill accumulation and yield in the California pistachio (*Pistacia vera*) cultivars 'Kerman' and 'Peters'. In Wirthensohn, M. G. (ed.) *VII International Symposium on Almonds and Pistachios, Acta Horticulturae 1219*. Leuven, Belgium: International Society for Horticultural Science, pp. 111–118.

¹⁴⁹ Doll, D., Beck, C., and Coelho, J. (2015). Kaolin clay may be useful in increasing chill accumulation in pistachios. Davis, CA: University of California Cooperative Extension. Available at: <Http://thealmonddoctor.com/2015/10/26/kaolin-Clay-May-Be-Useful-in-Increasing-Chill-Accumulation-in-Pistachios> (Accessed: 1 March 2022).

¹⁵⁰ Pope, K. S., da Silva, D., Brown, P. H., and DeJong, T. M. (2014). A biologically based approach to modeling spring phenology in temperate deciduous trees. *Agricultural and Forest Meteorology*, 198–199, 15–23.

¹⁵¹ Elias, E., Reyes, J., Steele, C., and Rango, A. (2018). Diverse landscapes, diverse risks: synthesis of the special issue on climate change and adaptive capacity in a hotter, drier Southwestern United States. *Climatic Change*, 148(3), 339–353.

¹⁵² Lamont, W. J. (2005). Plastics: Modifying the microclimate for the production of vegetable crops. *HortTechnology*, 15(3), 477–481.

¹⁵³ Houston, L., Capalbo, S., Seavert, C., Dalton, M., Bryla, D., and Sagili, R. (2018). Specialty fruit production in the Pacific Northwest: adaptation strategies for a changing climate. *Climatic Change*, 146(1), 159–171.

¹⁵⁴ Haruna, S. I., Anderson, S. H., Nkongolo, N. V., Reinbott, T., and Zaibon, S. (2017). Soil thermal properties influenced by perennial biofuel and cover crop management. *Soil Science Society of America Journal*. *Soil Science Society of America*, 81(5), 1147–1156. (Also see Easing the soil's temperature: Cover crops shield soil from extreme temps <https://www.sciencedaily.com/releases/2017/11/171108092434.htm>)

¹⁵⁵ Barron-Gafford, G. A., Pavao-Zuckerman, M. A., Minor, R. L., Sutter, L. F., Barnett-Moreno, I., Blackett, D. T., Thompson, M., Dimond, K., Gerlak, A. K., Nabhan, G. P., and Macknick, J. E. (2019). Agrivoltaics provide mutual benefits across the food–energy–water nexus in drylands. *Nature Sustainability*, 2(9), 848–855.

¹⁵⁶ Goss, M., Swain, D. L., Abatzoglou, J. T., Sarhadi, A., Kolden, C. A., Williams, A. P., and Diffenbaugh, N. S. (2020). Climate change is increasing the likelihood of extreme autumn wildfire conditions across California. *Environmental Research Letters*, 15(9), 094016.

¹⁵⁷ Arnold, C., Ghezzehei, T. A., and Berhe, A. A. (2014). Early spring, severe frost events, and drought induce rapid carbon loss in high elevation meadows. *PLoS One*, 9(9), e106058.

¹⁵⁸ Lamichhane, J. R. (2021). Rising risks of late-spring frosts in a changing climate. *Nature Climate Change*, 11(7), 554–555.

¹⁵⁹ Petanidou, T., Kallimanis, A. S., Sgardelis, S. P., Mazaris, A. D., Pantis, J. D., and Waser, N. M. (2014). Variable flowering phenology and pollinator use in a community suggest future phenological mismatch. *Acta Oecologica*, 59, 104–111.

¹⁶⁰ Kodad, O., and Socias i Company, R. (2014). Erratic fruit set in almond under warm climates. *International Journal of Horticultural Science*, 20(1–2), 59–64.

¹⁶¹ Lopez, G., Johnson, R. S., and DeJong, T. M. (2007). High spring temperatures decrease peach fruit size. *California Agriculture*, 61(1), 31–34.

¹⁶² Klein, A.-M., Brittain, C., Hendrix, S. D., Thorp, R., Williams, N., and Kremen, C. (2012). Wild pollination services to California almond rely on semi-natural habitat. *The Journal of Applied Ecology*, 49(3), 723–732.

¹⁶³ Vaughan, M., and Mader, E. (2009). *Pollinator Biology and Habitat in California*. https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/16/nrcs143_022094.pdf

¹⁶⁴ Anwar, M. R., Liu, D. L., Macadam, I., and Kelly, G. (2013). Adapting agriculture to climate change: a review. *Theoretical and Applied Climatology*, 113(1), 225–245.

¹⁶⁵ Lin, B. B. (2011). Resilience in agriculture through crop diversification: adaptive management for environmental change. *Bioscience*, 61(3), 183–193.

¹⁶⁶ Ahmed, S., and Stepp, J. R. (2016). Beyond yields: Climate change effects on specialty crop quality and agroecological management. *Elementa* (Washington, D.C.), 4, 000092.

¹⁶⁷ Jackson, L. E., Wheeler, S. M., Hollander, A. D., and O'Geen, A. T. (2011). Case study on potential agricultural responses to climate change in a California landscape. *Climatic Change*, 109, 407–427.

¹⁶⁸ Alcon, F., Marín-Miñano, C., Zabala, J. A., de-Miguel, M.-D., and Martínez-Paz, J. M. (2020). Valuing diversification benefits through intercropping in Mediterranean agroecosystems: A choice experiment approach. *Ecological Economics: The Journal of the International Society for Ecological Economics*, 171, 106593.

¹⁶⁹ Li, C., He, X., Zhu, S., Zhou, H., Wang, Y., Li, Y., Yang, J., Fan, J., Yang, J., Wang, G., Long, Y., Xu, J., Tang, Y., Zhao, G., Yang, J., Liu, L., Sun, Y., Xie, Y., Wang, H., and Zhu, Y. (2009). Crop diversity for yield increase. *PLoS One*, 4(11), e8049.

¹⁷⁰ Luedeling, E. (2012). Climate change impacts on winter chill for temperate fruit and nut production: A review. *Scientia Horticulturae*, 144, 218–229.

¹⁷¹ Katz, R. W., and Brown, B. G. (1992). Extreme events in a changing climate: Variability is more important than averages. *Climatic Change*, 21(3), 289–302.

¹⁷² Rasmussen, D. J., Holloway, T., and Nemet, G. F. (2011). Opportunities and challenges in assessing climate change impacts on wind energy—a critical comparison of wind speed projections in California. *Environmental Research Letters: ERL*, 6(2), 024008.

¹⁷³ Hatfield, J. L., and Prueger, J. H. (2015). Temperature extremes: Effect on plant growth and development. *Weather and Climate Extremes*, 10, 4–10.

¹⁷⁴ Rosenzweig, C., Tubiello, F. N., Goldberg, R., Mills, E., and Bloomfield, J. (2002). Increased crop damage in the US from excess precipitation under climate change. *Global Environmental Change: Human and Policy Dimensions*, 12(3), 197–202.

¹⁷⁵ Gershunov, A., and Guirguis, K. (2012). California heat waves in the present and future. *Geophysical Research Letters*, 39(18). L18710.

¹⁷⁶ Sheridan, S. C., and Lee, C. C. (2018). Temporal trends in absolute and relative extreme temperature events across north America. *Journal of Geophysical Research*, 123(21), 11889–11898.

¹⁷⁷ Greer, D. H., and Weston, C. (2010). Heat stress affects flowering, berry growth, sugar accumulation and photosynthesis of *Vitis vinifera* cv. Semillon grapevines grown in a controlled environment. *Functional Plant Biology: FPB*, 37(3), 206–214.

¹⁷⁸ Smith, P., and Olesen, J. E. (2010). Synergies between the mitigation of, and adaptation to, climate change in agriculture. *The Journal of Agricultural Science*, 148(5), 543–552.

¹⁷⁹ Moriondo, M., and Bindi, M. (2007). Impact of climate change on the phenology of typical Mediterranean crops. *Italian Journal of Agrometeorology*, 3, 5–12.

¹⁸⁰ What are atmospheric Rivers? National Oceanic and Atmospheric Administration. Available at: <https://www.noaa.gov/stories/what-are-atmospheric-rivers>

¹⁸¹ Espinoza, V., Waliser, D. E., Guan, B., Lavers, D. A., and Ralph, F. M. (2018). Global analysis of climate change projection effects on atmospheric rivers. *Geophysical Research Letters*, 45(9), 4299–4308.

¹⁸² Wang, S.-Y. S., Yoon, J.-H., Becker, E., and Gillies, R. (2017). California from drought to deluge. *Nature Climate Change*, 7(7), 465–468.

¹⁸³ Southwick, S., and Uyemoto, J. (1999). Cherry crinkle-leaf and deep suture disorders. Davis, CA: University of California, Department of Agriculture and Natural Resources.

¹⁸⁴ Murphy, S. F., Writer, J. H., Blaine McCleskey, R., and Martin, D. A. (2015). The role of precipitation type, intensity, and spatial distribution in source water quality after wildfire. *Environmental Research Letters*, 10(8), 084007.

¹⁸⁵ Ficklin, D. L., Luo, Y., Luedeling, E., Gatzke, S. E., and Zhang, M. (2010). Sensitivity of agricultural runoff loads to rising levels of CO₂ and climate change in the San Joaquin Valley watershed of California. *Environmental Pollution*, 158(1), 223–234.

¹⁸⁶ Smukler, S. M., Sánchez-Moreno, S., Fonte, S. J., Ferris, H., Klonsky, K., O'Geen, A. T., Scow, K. M., Steenwerth, K. L., and Jackson, L. E. (2010). Biodiversity and multiple ecosystem functions in an organic farmscape. *Agriculture, Ecosystems and Environment*, 139(1), 80–97.

¹⁸⁷ Kaye, J. P., and Quemada, M. (2017). Using cover crops to mitigate and adapt to climate change. A review. *Agronomy for Sustainable Development*, 37(1), 4.

¹⁸⁸ Steinwand, M. A., and Ronald, P. C. (2020). Crop biotechnology and the future of food. *Nature Food*, 1(5), 273–283.

¹⁸⁹ Sydeman, W. J., García-Reyes, M., Schoeman, D. S., Rykaczewski, R. R., Thompson, S. A., Black, B. A., and Bograd, S. J. (2014). Climate change. Climate change and wind intensification in coastal upwelling ecosystems. *Science*, 345(6192), 77–80.

¹⁹⁰ Guzman-Morales, J., and Gershunov, A. (2019). Climate change suppresses Santa Ana winds of southern California and sharpens their seasonality. *Geophysical Research Letters*, 46(5), 2772–2780.

¹⁹¹ Wang, M., Ullrich, P., and Millstein, D. (2020). Future projections of wind patterns in California with the variable-resolution CESM: a clustering analysis approach. *Climate Dynamics*, 54(3), 2511–2531.

¹⁹² Zamani, S., and Mahmoodabadi, M. (2013). Effect of particle-size distribution on wind erosion rate and soil erodibility. *Archives of Agronomy and Soil Science*, 59(12), 1743–1753.

¹⁹³ Merhav, D. J., Corkidi, L., Faber, B., Newman, J., and Webb, S. (2012). Best management practices in avocado orchard management. *California Avocado Society Yearbook 95: 70-88*. Available at: http://www.avocadosource.com/cas_yearbooks/cas_95_2012/cas_2012_v95_pg_070-088.pdf.

¹⁹⁴ Brandle, J. R., Hodges, L., and Zhou, X. H. (2004). Windbreaks in North American agricultural systems. In Nair, P. K. R., Rao, M. R., and Buck, L.E. (Eds.), *New Vistas in Agroforestry: A Compendium for 1st World Congress of Agroforestry, 2004*. Dordrecht, Netherlands: Springer Netherlands, pp. 65–78.

¹⁹⁵ Blanco-Canqui, H., Shaver, T. M., Lindquist, J. L., Shapiro, C. A., Elmore, R. W., Francis, C. A., and Hergert, G. W. (2015). Cover crops and ecosystem services: Insights from studies in temperate soils. *Agronomy Journal*, 107(6), 2449–2474.

¹⁹⁶ Smarik, S., S. Aney, A. Boes, D. Brown, D., Dubois, B. Edwards, E. Elias, M. Eve, R. Steele, N. Webb, M. Wilson, G. Zwicke, G. 2019. *Dust Mitigation Handbook*. Las Cruces, NM: U.S. Department of Agriculture.

¹⁹⁷ Pausas, J. G., and Keeley, J. E. (2019). Wildfires as an ecosystem service. *Frontiers in Ecology and the Environment*, 17(5), 289–295.

¹⁹⁸ Williams, A. P., Abatzoglou, J. T., and Gershunov, A. (2019). Observed impacts of anthropogenic climate change on wildfire in California. *Earth's Future* 7(8), 892–910. Available at: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019EF001210>

¹⁹⁹ Westerling, A. L., Bryant, B. P., Preisler, H. K., Holmes, T. P., Hidalgo, H. G., Das, T., and Shrestha, S. R. (2011). Climate change and growth scenarios for California wildfire. *Climatic Change*, 109(1), 445–463.

²⁰⁰ Swain, D. L. (2021). A shorter, sharper rainy season amplifies California wildfire risk. *Geophysical Research Letters*, 48(5), e2021GL092843.

²⁰¹ Fryer, J. A., Collins, T. S., and Tomasino, E. (2021). Evaluation of different interstimulus rinse protocols on smoke attribute perception in wildfire-affected wines. *Molecules*, 26(18), 5444. Available at: <https://doi.org/10.3390/molecules26185444>

²⁰² Summerson, V., Gonzalez Viejo, C., Pang, A., Torrico, D. D., and Fuentes, S. (2021). Review of the effects of grapevine smoke exposure and technologies to assess smoke contamination and taint in grapes and wine. *Beverages*, 7(1), 7.

²⁰³ Favell, J. W., Noestheden, M., Lyons, S. M., and Zandberg, W. F. (2019). Development and evaluation of a vineyard-based strategy to mitigate smoke-taint in wine grapes. *Journal of Agricultural and Food Chemistry*, 67(51), 14137–14142.

²⁰⁴ Suffling, R., Grant, A., and Feick, R. (2008). Modeling prescribed burns to serve as regional firebreaks to allow wildfire activity in protected areas. *Forest Ecology and Management*, 256(11), 1815–1824.

²⁰⁵ O'Connell, D., and Livingston, A. (2017). Ecosystem services and California's working landscapes. Davis, CA: University of California, Division of Agriculture and Natural Resources. Available at: <https://ucanr.edu/files/272736.pdf>

²⁰⁶ Bryant, B. P., Kelsey, T. R., Vogl, A. L., Wolny, S. A., MacEwan, D., Selmants, P. C., Biswas, T., and Butterfield, H. S. (2020). Shaping land use change and ecosystem restoration in a water-stressed agricultural landscape to achieve multiple benefits. *Frontiers in Sustainable Food Systems*, 4, 138.

²⁰⁷ Brodt, S., Klonsky, K., Jackson, L., Brush, S. B., and Smukler, S. (2009). Factors affecting adoption of hedgerows and other biodiversity-enhancing features on farms in California, USA. *Agroforestry Systems*, 76(1), 195–206.

²⁰⁸ Power, A. G. (2010). Ecosystem services and agriculture: tradeoffs and synergies. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 365(1554), 2959–2971.

²⁰⁹ Drexler, S., Gensior, A., and Don, A. (2021). Carbon sequestration in hedgerow biomass and soil in the temperate climate zone. *Regional Environmental Change*, 21(3), 74.

²¹⁰ Cooper, C. M., and Moore, M. T. (2003). Wetlands and Agriculture. In Holland, M., Blood, E. R. and Shaffer, L. R. (eds.) *Achieving Sustainable Freshwater Systems: A Web of Connections*. Washington, D.C.: Island Press, pp. 221–236.

²¹¹ Moore, M. T., Denton, D. L., Cooper, C. M., Wrysinski, J., Miller, J. L., Reece, K., Crane, D., and Robins, P. (2008). Mitigation assessment of vegetated drainage ditches for collecting irrigation runoff in California. *Journal of Environmental Quality*, 37(2), 486–493.

²¹² Gliessman, S. R. (2014). *Agroecology: the Ecology of Sustainable Food Systems*. Boca Raton, FL: CRC Press.

²¹³ York, R. A., Levine, J., Russell, K., and Restaino, J. (2021). Opportunities for winter prescribed burning in mixed conifer plantations of the Sierra Nevada. *Fire Ecology*, 17(1), 33.

²¹⁴ Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., and Courchamp, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, 15(4), 365–377.

²¹⁵ Kremen, C., and Merenlender, A. M. (2018). Landscapes that work for biodiversity and people. *Science*, 362(6412), eaau6020.

²¹⁶ Power, A. G. (2010). Ecosystem services and agriculture: tradeoffs and synergies. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 365(1554), 2959–2971.

²¹⁷ Hilty, J. A., and Merenlender, A. M. (2004). Use of riparian corridors and vineyards by mammalian predators in northern California. *Conservation Biology: The Journal of the Society for Conservation Biology*, 18(1), 126–135.

²¹⁸ Sirami, C., Gross, N., Baillod, A. B., Bertrand, C., Carrié, R., Hass, A., Henckel, L., Miguët, P., Vuillot, C., Alignier, A., Girard, J., Batáry, P., Clough, Y., Violette, C., Giralt, D., Bota, G., Badenhausser, I., Lefebvre, G., Gauffre, B., Fahrig, L. (2019). Increasing crop heterogeneity enhances multitrophic diversity across agricultural regions. *Proceedings of the National Academy of Sciences of the United States of America*, 116(33), 16442–16447.

²¹⁹ Vinge, H. (2018). Farmland conversion to fight climate change? Resource hierarchies, discursive power and ulterior motives in land use politics. *Journal of Rural Studies*, 64, 20–27.

²²⁰ Vahmani, P., Sun, F., Hall, A., and Ban-Weiss, G. (2016). Investigating the climate impacts of urbanization and the potential for cool roofs to counter future climate change in Southern California. *Environmental Research Letters*, 11(12), 124027.

²²¹ California Department of Conservation Farmland Mapping and Monitoring Program (n.d.). Net important farmland conversion 1984–2014. Available at: https://www.conservation.ca.gov/dlfp/fmmp/Documents/fmmp_1984_2014_summary.xlsx. (Accessed: 29 July 2022).

²²² United States Environmental Protection Agency. Fast Facts 1990–2020 National-level U.S. Greenhouse Gas Inventory. Available at: <https://www.epa.gov/system/files/documents/2022-04/fastfacts-1990-2020.pdf>. (Accessed: 1 July 2022).

²²³ California Air Resources Board. Greenhouse Gas Emission Inventory Data. Available at: <https://ww2.arb.ca.gov/ghg-inventory-data>. (Accessed: 1 July 2022).

²²⁴ Deng, J., Guo, L., Salas, W., Ingraham, P., Charrier-Klobas, J. G., Frolking, S., Li, C. (2018). Changes in irrigation practices likely mitigate nitrous oxide emissions from California cropland. *Global Biogeochemical Cycles*, 32, 1514–1527.

²²⁵ California Air Resources Board. (2021). California Greenhouse Gas Emissions for 2000 to 2019: Trends of Emissions and Other Indicators. Available at: https://ww2.arb.ca.gov/sites/default/files/classic/cc/inventory/2000_2019_ghg_inventory_trends_20220516.pdf

²²⁶ Zhu-Barker, X., Easter, M., Swan, A., Carlson, M., Thompson, L., Horwath, W. R., Paustian, K., and Steenwerth, K. L. (2019). Soil management practices to mitigate nitrous oxide emissions and inform emission factors in arid irrigated specialty crop systems. *Soil Systems*, 3(4), 76.

²²⁷ Kennedy, T. L., Suddick, E. C., and Six, J. (2013). Reduced nitrous oxide emissions and increased yields in California tomato cropping systems under drip irrigation and fertigation. *Agriculture, Ecosystems and Environment*, 170, 16–27.

²²⁸ Suddick, E. C., Scow, K. M., Horwath, W. R., Jackson, L. E., Smart, D. R., Mitchell, J., and Six, J. (2010). Chapter Four - The potential for California agricultural crop soils to reduce greenhouse gas emissions: A holistic evaluation. In Sparks, D. L. (Ed.), *Advances in Agronomy*, Volume 107. Cambridge, MA: Academic Press, pp. 123–162.

²²⁹ Dupraz, C., Marrou, H., Talbot, G., Dufour, L., Nogier, A., and Ferard, Y. (2011). Combining solar photovoltaic panels and food crops for optimising land use: Towards new agrivoltaic schemes. *Renewable Energy*, 36(10), 2725–2732.

²³⁰ Wu, Y., Li, Y., Fu, X., Shen, J., Chen, D., Wang, Y., Liu, X., Xiao, R., Wei, W., and Wu, J. (2018). Effect of controlled-release fertilizer on N₂O emissions and tea yield from a tea field in subtropical central China. *Environmental Science and Pollution Research International*, 25(25), 25580–25590.

²³¹ Cayuela, M. L., Aguilera, E., Sanz-Cobena, A., Adams, D. C., Abalos, D., Barton, L., Ryals, R., Silver, W. L., Alfaro, M. A., Pappa, V. A., Smith, P., Garnier, J., Billen, G., Bouwman, L., Bondeau, A., and Lassaletta, L. (2017). Direct nitrous oxide emissions in Mediterranean climate cropping systems: Emission factors based on a meta-analysis of available measurement data. *Agriculture, Ecosystems and Environment*, 238, 25–35.

²³² Ziska, L. H. (2021). Crop adaptation: Weedy and crop wild relatives as an untapped resource to utilize recent increases in atmospheric CO₂. *Plants*, 10(1), 88.

²³³ Tautges, N. E., Chiartas, J. L., Gaudin, A. C. M., O'Geen, A. T., Herrera, I., and Scow, K. M. (2019). Deep soil inventories reveal that impacts of cover crops and compost on soil carbon sequestration differ in surface and subsurface soils. *Global Change Biology*, 25(11), 3753–3766.

²³⁴ Williams, J. N., Hollander, A. D., O'Geen, A. T., Thrupp, L. A., Hanifin, R., Steenwerth, K., McGourty, G., and Jackson, L. E. (2011). Assessment of carbon in woody plants and soil across a vineyard-woodland landscape. *Carbon Balance and Management*, 6(1), 11.

²³⁵ Lal, R. (2004). Soil carbon sequestration impacts on global climate change and food security. *Science*, 304(5677), 1623–1627.

²³⁶ Lal, R., Negassa, W., and Lorenz, K. (2015). Carbon sequestration in soil. *Current Opinion in Environmental Sustainability*, 15, 79–86.

²³⁷ Kirkby, C. A., Richardson, A. E., Wade, L. J., Batten, G. D., Blanchard, C., and Kirkegaard, J. A. (2013). Carbon-nutrient stoichiometry to increase soil carbon sequestration. *Soil Biology and Biochemistry*, 60, 77–86.

²³⁸ Moran, K. K., Six, J., Horwath, W. R., and van Kessel, C. (2005). Role of mineral-nitrogen in residue decomposition and stable soil organic matter formation. *Soil Science Society of America Journal. Soil Science Society of America*, 69(6), 1730–1736.

²³⁹ USDA Natural Resources Conservation Service. (2011). Carbon to Nitrogen Ratios in Cropping Systems. Available at: https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcseprd331820.pdf

²⁴⁰ de la Rosa, J. M., Rosado, M., Paneque, M., Miller, A. Z., and Knicker, H. (2018). Effects of aging under field conditions on biochar structure and composition: Implications for biochar stability in soils. *Science of the Total Environment*, 613, 969–976.

²⁴¹ De Stefano, A., and Jacobson, M. G. (2017). Soil carbon sequestration in agroforestry systems: a meta-analysis. *Agroforestry Systems*, 92, 285–299.

²⁴² Gioacchini, P., Cattaneo, F., Barbanti, L., Montecchio, D., Ciavatta, C., and Marzadori, C. (2016). Carbon sequestration and distribution in soil aggregate fractions under Miscanthus and giant reed in the Mediterranean area. *Soil and Tillage Research*, 163, 235–242.

