



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

AGRO PRODUCTIVIDAD

**Water supply
in artificial
troughs:**
a strategy to mitigate
the impacts of climate
change in the Maya forest

pág. 69

Año 17 • Volumen 17 • Número 4 • abril, 2024

The Mexican Beekeeping Agri-food System: A descriptive analysis 3

Productive response of dual-purpose cows supplemented with chelate minerals and calcium salts 11

Materials and methods for the microencapsulation of substances of food and agricultural interest 19

Blood disorders caused by hypophosphatemia in dairy cows 27

Efficacy of chemical fungicides against the anthracnose disease caused by *Colletotrichum gloeosporioides* in *Carica papaya* fruits 35

Cardinal temperatures of populations developed from native maize (*Zea mays* L.) from central and southern, Tamaulipas, Mexico 47

y más artículos de interés...



Colegio de
Postgraduados

In vitro gas production and digestibility of oat and triticale forage mixtures ensiled with fibrolytic enzymes and inoculants

González Reyes, Mónica¹; López-Rodríguez, Fredy¹; Gayosso-Barragán, Odilón²; Miranda-Romero, Luis Alberto³; Tirado-González, Deli Nazmín^{4*}; Tirado-Estrada, Gustavo^{4*}

¹ Tecnológico Nacional de México/Tecnológico El Llano Aguascalientes, División de Estudios de Posgrado e Investigación, Carretera Aguascalientes-S.L.P. Km. 18, El Llano, Aguascalientes, México, C. P. 20330.

² Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias (INIFAP)/Centro Nacional de Investigación Disciplinaria Agricultura Familiar (CENID AF), Carretera Ojuelos-Lagos de Moreno Km 8.5, Ojuelos, Jalisco, México, C. P. 47540.

³ Universidad Autónoma Chapingo, Posgrado en Producción Animal, Departamento de Zootecnia, Carretera Federal México-Texcoco Km 38.5, Texcoco, Estado de México, México, C. P. 56230.

⁴ Tecnológico Nacional de México/Tecnológico El Llano Aguascalientes, Departamento de Ingenierías, Carretera Aguascalientes-S.L.P. Km. 18, El Llano, Aguascalientes. México. C.P. 20330.

* Correspondence: deli.tg@llano.tecnm.mx; gustavo.te@llano.tecnm.mx

ABSTRACT

Objective: To assess the effects of adding fibrolytic enzymes (FE) or lactic acid bacteria (LAB) inoculants to 40 d silages with oat and triticale (O:T) mixtures on the ratio and composition of neutral detergent fiber (NDF) and its subsequent *in vitro* gas production (GP) and *in vitro* dry matter digestibility (IVDMD) at 24 h.

Design/Methodology/Approach: Silages elaborated with two O:T ratios (60:40 and 80:20) treated with low (LD), medium (MD), and high (HD) doses of FE (0.75, 1, and 1.25 g/kg forage in wet basis (WB), respectively), and LAB (0.188, 0.25, and 0.31 g/kg WB, respectively). In both cases (FE and LAB), the control had a value of 0. Subsequently, pH, NDF, acid detergent fiber (ADF), acid detergent lignin (ADL), hemicellulose (HEM), cellulose (CEL), dry matter (DM), crude protein (CP), GP parameters, and IVDMD were assessed. GP parameters included maximum velocity (V_{max}), fractional rate (S), and lag. Experiments were planned in complete randomized designs (CRD), including factorial and split-plot arrangements. Variance analysis (ANOVA) models included fixed (doses, additives, and FR) and random (place/moment of sampling) effects.

Results: LAB improved the IVDMD at 24 h of 60:40 and 80:20 O:T silages. FE did not reduce the NDF of 60:40 silages, but LD and MD increased the HEM and CP, and reduced the ADF, ADL, and CEL; these results are correlated (r) with the improvement of pH pattern, GP, and IVDMD.

Study Limitations/Implications: The differences in the NDF of FR mixtures could affect the effectiveness of FE and LAB.

Findings/Conclusions: Although FE and LAB did not reduce the NDF, they changed the ratios of ADF, ADL, HEM, CEL, and CP of silages, potentially improving the GP and IVDMD.

Keywords: Oats and triticale, ensiling, fibrolytic enzymes, lactic acid bacteria, gas production, ruminal degradability.

Citation: González-Reyes, M., López-Rodríguez, F., Gayosso-Barragán, O., Miranda-Romero, L. A. Tirado-González, D.J., & Tirado-Estrada, G. (2024). *In vitro* gas production and digestibility of oat and triticale forage mixtures ensiled with fibrolytic enzymes and inoculants. *Agro Productividad*. <https://doi.org/10.32854/agrop.v17i4.2711>

Academic Editors: Jorge Cadena Iniguez and Lucero del Mar Ruiz Posadas

Guest Editor: Daniel Alejandro Cadena Zamudio

Received: October 24, 2023.

Accepted: March 18, 2024.

Published on-line: May 06, 2024.

Agro Productividad, 17(4). April. 2024. pp: 137-149.

This work is licensed under a Creative Commons Attribution-Non-Commercial 4.0 International license.



INTRODUCTION

In semi-arid regions, the high frequency of droughts has a negative effect on the availability and nutritional value of forages (Acosta *et al.*, 2003), limiting the ability of

producers and ranchers from mainly extensive and family systems to maintain the production of bovine milk and sheep, goat, and bovine meat. Therefore, maximizing the ratio of fibrous forages in the diet of ruminants, without reducing the quality and quantity of the products (Tirado-González *et al.*, 2018; Tirado-Estrada *et al.*, 2020), can help to reduce production costs (Oba and Allen, 1999, 2000a, b; Tirado-Estrada *et al.*, 2015), improve animal health (Saleem *et al.*, 2012; Petri *et al.*, 2013), and diminish the environmental impact of deforestation, consequently maintaining intensive grain production (McGinn *et al.*, 2004; Knapp *et al.*, 2014; Mora de Alba *et al.*, 2018).

The ensiling process contributes to the conservation and modification of the degradability of the cell walls of the forages, improving the nutritional characteristics of the forage through the formation of lactic acid and alcohols and the increase in crude protein (CP) (Tiwari *et al.*, 2008; Ajila *et al.*, 2015; Chen *et al.*, 2016). Meanwhile, the use of lactic acid bacteria (LAB) inoculants can improve the carbohydrate fermentation period, while the volatile fatty acid (VFA) profile and the formation of lactic acid increase the aerobic stability and quality of silages (Tabacco *et al.*, 2011; Skládanka *et al.*, 2012; Guo *et al.*, 2013; Schroeder, 2013). Fibrolytic enzymes (FE) can help to break the bonds of the cellulose and hemicellulose components (Arriola *et al.*, 2017; Kholif *et al.*, 2017; Tirado-González *et al.*, 2015, 2018), favoring the fermentation process through the increase of the sugars available for the microorganisms (Gado *et al.*, 2013; Salem *et al.*, 2015; Kholif *et al.*, 2017). However, the efficient action of LAB and FE depends on the interaction of various factors, such as: type and ratios of forages (Dehghani *et al.*, 2012), LAB strain or FE mixture (Dean *et al.*, 2005; Lynch *et al.*, 2012), dose (Del Valle *et al.*, 2019), and ensiling time (Lynch *et al.*, 2012). This phenomenon is caused by the high specificity of the enzymatic components, which record small differences in their neutral detergent fiber (NDF), acid detergent fiber (ADF), and acid detergent lignin (ADL) composition (Tirado-González *et al.*, 2016).

The objective of this work is to assess the effect of the use of EF and LAB in silages with oat and triticale mixtures on the contents and compositions of NDF and their relationship with GP and IVDMD.

MATERIALS AND METHODS

Experiment location and biological material

Representative samples of 60:40 and 80:20 ratios of oat (*Avena sativa* L.) and triticale (*Triticosecale* Wittmack. Ex. A. Camus) were taken in two locations in the central-northern region of Mexico:

- 1) El Llano, Aguascalientes (21° 55' 00 N, 101° 58' 00 W; 2,021 m.a.s.l.). The climate is semi-arid (BS1kw, Köppen), with an average temperature of 17.4 °C and an average annual precipitation of 540 mm. The soil mainly consists of planosol (66%) and phaeozem (23%).
- 2) Lagos de Moreno, Jalisco (21° 21' 23 N, 101° 55' 45 W; 1,942 m.a.s.l.). The climate is subtropical (Csa, Köppen), with an average temperature of 18.4 °C and an average annual precipitation of 670 mm. The soil is mainly composed of lithosol, planosol, and phaeozem (31%, 29% and 25%, respectively).

The samples were taken 120 hours after the cultivation began: 10 samples (20 kg of complete plants/sample) were selected at random in each of the blocks (DM=33.1±2.7%).

Preparation of microsilages

The microsilages and chemical analyzes were carried out at TecNM/ITEL (21° 55' N, 101° 58' W; 1,840 m.a.s.l.), in El Llano, Aguascalientes (average annual temperature of 17 °C, average annual precipitation of 455 mm).

The additives applied to the microsilages were an FE preparation —with different ratios of cellulases and xylanases (Fibrozyme, Alltech Inc., Nicholasville, KY, USA)— and a LAB inoculant —with *Lactobacillus plantarum* and *Pediococcus pentosaceus* (EnziBiolac, Enzimas y Productos Químicos, S. A. de C. V., SAGARPA Reg. A-9912-001, Mexico). Samples of 1±0.1 kg (WB) of the 60:40 and 80:20 O:T mixtures collected in the two locations (DM=30±2.5%) were chopped into 2 cm particles and treated with: 1) 0 (control), 0.75 (LD), 1 (MD), and 1.25 g (LD) doses of FE/kg WB; and 2) 0 (control), 0.188 (LD), 0.25 (MD), and 0.31 g (HD) doses of LAB/kg WB.

The FE and LAB mixtures were dissolved in distilled water and sprayed uniformly on the forage. The treatments were placed in 5.08 cm wide×30 cm long polyvinyl chloride (PVC) pipes (with reinforced PVC end caps installed at both ends of the pipe). The treatments were compacted with a metal piston (to eliminate as much oxygen as possible inside the microsilages) and stored in a closed room with an average temperature of 20 °C. They were then allowed to ferment for 40 days.

Chemical analysis

At the time of ensiling and every 0, 10, 20, 30, and 40 d after the start of the ensiling process, 300 g WB/microsilage fractions were sampled from different parts of each microsilage. These samples were placed in a Felisa[®] AR-290 forced air oven at 60 °C, until a constant weight (initial DM; 0 d) was reached. The dried samples were ground with a 1 mm sieve.

In the chemical analysis, the pH of the samples was measured using an Orion Star[™] A2110 pH-meter (Thermo Scientific). The initial and final DM, crude protein (CP), and ash (ASH) were determined using the 10.136, 990.03 and 942.05 methods, respectively (AOAC, 2005). The NDF, ADF, and ADL of the samples collected from the microsilages at 40 d were determined (Van Soest *et al.*, 1991). For this purpose, the samples were adapted for the reagents of the F57 filters (2016, Ankom Technol Technology, Macedon, NY, USA) and an ANKOM²⁰⁰ Fiber Analyzer (ANKOM Technology,). Hemicellulose (HEM) and cellulose (CEL) were calculated by difference (HEM=NDF-ADF and CEL=ADF-ADL-ASH). Each sample was analyzed in duplicate.

In vitro fermentation

The gas production of the silage samples taken at 40 d was analyzed using the gas production technique published by Menke and Steingass (1998). Ruminant fluid was obtained from two cannulated Dorper sheep (live weight: 60±5 kg) fed with a 76:24 forage:concentrate ratio (forage: barley (*Hordeum vulgare*) straw; concentrate: mixture of

corn and soybean grain). The chemical composition of the forage was: 58% NDF, 10% CP, and 44.1% crude fiber (CF). Meanwhile, the chemical composition of the concentrate was: 22.4% NDF, 16% CP, and 5.4% CF.

The ruminal fluid was filtered with an 8-layer gauze and mixed with the mineral solutions reported by Cobos and Yokoyama (1995). Once the ruminal inoculum was prepared, it was placed in amber bottles with 0.5 g of DM and 90 mL of inoculum from the samples taken from the microsilages at 0 and 40 d (each sample was analyzed in duplicate). Incubation was carried out in a water bath (39 °C) under conditions of continuous flow of carbon dioxide (CO₂). The excess CO₂ from each bottle was extracted with the 63100 analog manometer (Metron[®]). The pressure of the fermentation gas (0 to 1 kg/cm²) was measured with the manometer at 2, 4, 6, 8, 12, 16, 20, 24, 30, 36, 42, 48, 60, and 72 h of incubation and it was converted to mL.

At the end of the incubation period, the residue from each bottle was filtered through a previously weighed filter paper (Whatman[®] qualitative filter paper, Grade 4: 1004-110, pore size: 20-25 μm); the filter papers with residue were dried at 65 °C for 48 h and subsequently weighed. After 24 h, the fermentation of half of the flasks was stopped and the IVDMD was determined as follows:

$$IVDMD_{24} = 100 - \left[\frac{(initialDM - finalDM)}{initialDM} * 100 \right]$$

The maximum volume, fractional rate, and lag phase (V_{max}, S, and Lag) parameters of gas production were optimized in Equation (1) of Schofield *et al.* (1994).

$$V_0 = \frac{V_{max}}{1 + e^{2-4S(t-Lag)}} \quad (1)$$

Where: V₀=accumulated GP volume; V_{max}=maximum volume of GP; S=fractional rate; Lag=lag phase.

Statistical analysis

Data analysis was performed with the SAS [9.2] statistical software (Statistical Analysis System). An analysis of variance (ANOVA) took into consideration the DCA, with factorial and split-plot arrangements with 4 repetitions per treatment, as well as 2 sub-repetitions for the fixed and random effects of Models 1 and 2. The significances, coefficients of determination (R²), and coefficients of variation (CV) were obtained using the General Linear Procedure (Proc GLM); the LsMeans instruction was used for the adjusted means; and the standard errors (SE) were determined with the Mixed Models Procedure (Proc Mixed). The DMS were calculated using the SE values (P=0.05).

Model 1

$$Y = \mu + Rep(Loc)_{ij} + Tra_k + T_l + (Tra * T)_{kl} + \varepsilon_{ijkl}$$

Where: $Y = \text{pH}$; μ is the overall mean; $Rep(Loc)_{ij}$ is the random effect of the i -th repetition within the j -th sowing location; Tra_k is the effect of the k -th treatment; T_l is the effect of the l -th time; $(Tra*T)_{kl}$ is the interaction between fixed factors; ε_{ijkl} is the random error.

Model 2

$$Y = \mu + Subrep(Rep)_{ij} + D_k + A_l + PAT_m + (D*A)_{kl} + (D*PAT)_{km} + (A*PAT)_{lm} + (D*A*PAT)_{ijklm} + \varepsilon_{ijklm}$$

Where: $Y = \text{initial DM, final DM, Vmax, S, Lag, pH, NDF, ADF, ADL, CP, ASH, and IVDMD24}$; $Subrep(Rep)_{ij}$ is the random effect of the i -th subrepetition within the j -th locality or analysis time; D_k is the effect of the k -th dose; A_l is the effect of the l -th additive; PAT_m is the effect of the m -th ratio of O:T; $(D*A)_{kl}$, $(D*PAT)_{km}$, $(A*PAT)_{lm}$, and $(D*A*PAT)_{ijklm}$ are the interactions of the fixed factors; ε_{ijklm} is the random error.

Correlation analysis. Simple Pearson correlation analyzes were performed to analyze the relationship between the Vmax, S, Lag, NDF, ADF, ADL, CP, ASH, and IVDMD24 variables.

RESULTS AND DISCUSSION

Changes in chemical composition and pH

Chemical composition. Table 1 shows the chemical composition of the silages. Overall, the use of LD and MD of FE and LAB in the 60:40 O:T silages did not affect the ratio of NDF. However, the HD of both additives increased the ratio of NDF in the 60:40 O:T silages. The LD of FE reduced the ratio of NDF ($P < 0.0009$) in 80:20 O:T silages. Furthermore, the ratio of ADF did not diminish in any of the FE and LAB treatments in both types of silage; even treatments with HD of LAB showed the highest ADF values compared to the control ($P < 0.02$). Although the additives did not consistently reduce NDF and ADF, their inclusion did affect the ratios of HEM and CEL: the 60:40 O:T silages had more HEM and less CEL than the 80:20 silages ($P < 0.008$), while silages with FE had a lower CEL content than those treated with LAB ($P < 0.01$). The ratios of ADL and ASH were lower in 60:40 O:T silages than in 80:20 silages ($P < 0.0002$); likewise, ASH were higher in silages treated with FE ($P < 0.03$).

The 60:40 O:T silages had better CP content than 80:20 silages ($P < 0.0001$). In the 60:40 O:T silages, the use of FE did not improve the CP in relation to the Control. In the 80:20 O:T silages, the use of LD, MD, and HD of FE increased the CP. Similarly, LD, MD, and HD of LAB improved the CP ratios of the 60:40 and 80:20 O:T silages ($P < 0.05$).

Higher initial and final DM were observed in the 60:40 O:T silages than in the 80:20 silages ($P < 0.0001$), as well as in those treated with FE with regard to those treated with LAB ($P < 0.01$).

Table 1. Nutritional quality of silages of oat and triticale (O:T) forage mixtures (60:40 and 80:20 ratios), supplemented with various doses of fibrolytic enzymes (FE) and lactic acid bacteria (LAB).

Silages		NDF (%)	HEM (%)	ADF (%)	CEL (%)	ADL (%)	CP (%)	Ashes (%)	PreE-DM (%)	PostE-DM (%)
Oats: triticale 60:40%										
EFE	Control	61.38c	28.28b	33.51de	19.25cd	4.15c	7.25c	9.69cd	36.88cd	41.88cd
	LD	61.48c	28.15ab	33.34e	19.32cd	4.17c	7.80bc	9.86bc	39.65a	44.81a
	MD	62.28c	28.78ab	33.5de	20.18cd	4.27c	7.76bc	9.04d	38.15b	43.16ab
	HD	64.61a	29.23a	35.38c	21.70bc	4.53bc	7.37c	9.15d	38.59ab	43.71b
ALB	Control	62.69bc	28.37ab	34.20d	20.36cd	4.23c	6.92d	9.17d	37.13c	42.13c
	LD	62.29c	29.24a	34.05d	20.85bc	4.26c	9.06a	8.94d	37.11c	42.08cd
	MD	62.54bc	27.95b	34.59cd	21.54bc	4.31c	8.06b	8.74d	37.73bc	42.89bc
	HD	63.68ab	28.39ab	35.29c	21.55bc	4.29c	7.35c	9.46cd	35.83d	40.98d
Oats: triticale 80:20%										
EFE	Control	63.01ab	28.86ab	35.16cd	19.38cd	4.49bc	5.09h	9.28cd	28.43f	33.53f
	LD	61.89c	27.29bc	34.6cd	19.33d	4.57b	6.38e	10.69ab	28.28f	33.44f
	MD	63.27b	27.45bc	35.81ab	20.49c	4.57b	5.92f	10.75a	29.48ef	34.87e
	HD	62.66bc	27.30c	34.36d	20.53c	4.65b	5.80f	10.18b	28.26f	33.36f
ALB	Control	63.74a	28.13b	35.61c	20.56c	4.62b	4.83h	9.27cd	29.14ef	34.24ef
	LD	63.71ab	27.25c	36.46b	22.05ab	4.75ab	5.47g	9.66c	27.68f	32.58fg
	MD	64.41a	26.88c	37.53a	23.23a	5.02a	5.44gh	9.28cd	28.22f	33.16g
	HD	64.25a	27.07c	37.17a	21.85b	4.89ab	5.41gh	9.44cd	29.05ef	34.01ef
P-values										
	Doses	0.006	0.84	<0.0001	0.006	0.12	0.0004	0.36	0.56	0.89
	Additive	0.004	0.01	<0.0001	0.004	0.24	0.04	0.003	0.03	0.05
	OT	0.37	<0.0001	<0.0001	0.008	0.0002	<.0001	<0.0001	<0.0001	<0.0001
	D*A	0.009	0.03	0.17	0.06	0.83	0.003	0.23	<0.0001	<0.0001
	D*OT	0.11	0.26	0.05	0.02	0.67	0.21	0.22	<0.0001	<0.0001
	A*OT	0.10	0.17	0.02	0.99	0.11	0.002	0.10	0.13	0.01
	D*A*OT	0.47	0.38	0.66	0.93	0.26	0.05	0.40	0.004	0.05
	R ²	0.65	0.59	0.8	0.57	0.44	0.87	0.48	0.98	0.97
	V.C. (%)	1.77	3.20	2.19	4.91	11.77	7.92	6.00	2.99	4.01
	LSD (0.05) =	1.05	1.04	0.72	1.20	0.34	0.46	0.50	0.95	1.01
	S.E.	0.62	0.63	0.42	0.73	0.20	0.27	0.29	0.583	0.571

Different letters represent statistical media differences; FE, fibrolytic enzymes; LAB, lactic acid bacteria; LD, low dose; MD, medium dose; HD, high dose; Initial DM, initial dry matter (0 d, prior to the ensiling process); Final DM, final dry matter (DM of the silage at 40 d); NDF, neutral detergent fiber; HEM, hemicellulose; ADF, acid detergent fiber; CEL, cellulose; ADL, acid detergent lignin; CP, crude protein; P-values, probability values; ASH, ashes; D, doses; A, additive (FE or LAB); ROT, oat:triticale ratio; R², determination coefficient; V.C., variation coefficient; S.E., standard error; LSD, least significant difference (P<0.05).

pH modifications during the ensiling process. Figure 1 shows the pH changes of the treatments over the course of the ensiling period. The pH of the 60:40 O:T silages decreased from 6.1 to 4.5 (average) in the first 10 d. The use of FE had no significant effect on the reduction of pH, but there were significant effects resulting from the use of LD and MD of LAB ($P < 0.05$). On the one hand, the use of HD of LAB increased pH (on day 20 after the fermentation began) with regard to the LD and MD of LAB ($P < 0.0001$). On the other hand, in the 80:20 O:T silages, the pH was reduced from 6.1 to 4.2 in the first 30 d. However, in average, the pH increased after 30 d of fermentation ($P < 0.01$) in all treatments.

***In vitro* fermentation of DM**

Table 2 shows the V_{max} , S , and Lag phase of GP of the treatments assessed *in vitro*, as well as the IVDMD24. Overall, differences in V_{max} and S were observed in the two types of silage ($P < 0.0001$). In 80:20 O:T silages, the use of FE and LAB did not increase V_{max} and S with regard to the control, but, in the 60:40 O:T silages, the use of LD and MD of FE and LAB resulted in higher V_{max} and S than in the control treatments ($P < 0.003$). Similarly, the MD of FE, and the LD, MD, and HD of LAB reduced the Lag phase time of the 60:40 O:T silages (5.73 h *vs.* 6.55 h, MD of FE *vs.* Control; 5.92 h *vs.* 6.97 h, average LD, MD, and HD of LAB *vs.* Control), but did not affect the Lag time of the 80:20 O:T silages ($P < 0.03$). The use of LD of FE and MD of LAB increased the IVDMD24 of the




Figure 1. Effects on pH during the ensiling process of 60:40 and 80:20 oat:triticale silages supplemented with additives: a) fibrolytic enzymes (FE) and b) lactic acid bacteria (LAB).

Table 2. *In vitro* gas production (GP) and *in vitro* dry matter digestibility (IVDMD) at 24 h of silages with 60:40 and 80:20 oat:triticale (O:T) ratios supplemented with different doses of fibrolytic enzymes (FE) and lactic acid bacteria (LAB).

Silages		Vmax mL/g	S mL gas/h	Lag h	IVDMD %
Oats: triticale 60:40%					
EFE	Control	302.53cd	0.032b	6.55b	66.41b
	LD	314.24bc	0.033a	6.52b	67.70a
	MD	333.28ab	0.032b	5.73c	65.21c
	HD	322.75b	0.031c	6.60b	65.81bc
ALB	Control	308.33c	0.032b	6.97ab	65.68bc
	LD	330.55ab	0.033a	6.02c	65.21c
	MD	331.58b	0.032b	6.01c	66.27ab
	HD	346.18a	0.033a	5.73c	62.66d
Oats: triticale 80:20%					
EFE	Control	283.83de	0.032b	6.50ab	60.95e
	LD	295.58cd	0.031c	6.61b	62.14de
	MD	294.28d	0.031c	6.79ab	59.96ef
	HD	302.01cd	0.031c	6.52b	59.95ef
ALB	Control	288.96de	0.032b	7.18a	59.76ef
	LD	300.79cd	0.031c	6.97ab	60.62ef
	MD	292.93de	0.031c	6.73ab	59.71f
	HD	277.10e	0.030d	7.01ab	59.95ef
P-values					
	Doses	0.25	0.05	0.42	0.0002
	Additives	0.05	0.84	0.83	<0.0001
	OT	<0.0001	<0.0001	<0.0001	<0.0001
	D*A	0.11	0.002	0.78	<0.0001
	D*OT	0.24	0.28	0.64	0.25
	A*OT	0.003	0.003	0.0005	0.0002
	D*A*OT	0.10	0.11	0.03	<0.0001
	R ²	0.80	0.79	0.94	0.91
	C.V. (%)	6.67	4.27	9.67	3.14
	LSD (0.05)=	17.00	0.001	0.52	1.20
	S.E.	10.33	0.0007	0.31	0.72

FE, fibrolytic enzyme; LAB, lactic acid bacteria; LD, low dose; MD, medium dose; HD, high dose; Vmax, maximum volume of GP; S, fractional gas production rate; Lag, lag phase; IVDMD, in vitro dry matter digestibility at 24 h; P-values, probability values; D, doses; A, additive (FE or LAB); OT, oat:triticale ratio; R², determination coefficient; V.C., variation coefficient; S.E., standard error; LSD, last significant difference (P<0.05).

60:40 O:T silages (67.70% vs. 66.41%, LD of FE vs. Control; 66.27% vs. 65.68%, MD of LAB vs. Control) (P<0.0001); however, using FE and LAB did not have a positive effect on the IVDMD of 80:20 O:T silages.

Pearson correlations

The IVDMD24 did not correlate with NDF (Table 3); however, it showed a positive correlation with the HEM and CP ratio, but negative correlation with ADF, CEL, and ADL ($P < 0.0001$). Consequently, the IVDMD24 had a positive correlation with Vmax and S and a negative correlation with the Lag phase ($P < 0.01$).

FE has been used to improve the digestibility of forages intended for ruminant feed. These products can combine different ratios of forms and isoforms of cellulases and xylanases, depending on the extraction time of the FE and the substrate from which it was obtained (Tirado-González *et al.*, 2015, 2016, 2018; Carrillo-Díaz *et al.*, 2022). Using FE can have positive effects on the production and kinetics of GP and IVDMD (Phakachoe *et al.*, 2013; Salem *et al.*, 2015; Li *et al.*, 2018).

The variation in the oat and triticale ratios in this work affected the action of the FE and LAB added to the silages, confirming the high specificity of the enzymes extracted from fungi and the potential synergisms and negative interactions between enzymes and bacteria reported in previous research (Tirado-González *et al.*, 2016). Furthermore, the inconsistent fermentation and in vitro degradability results arising from the use of enzymes between silages with different O:T ratios are explained by the interaction between the types of cell walls with the exogenous enzyme preparations and the activity of bacteria and endogenous FE (Beauchemin *et al.*, 2003; Tirado-González *et al.*, 2015). The components and structures of the cell walls of forages depend on their type, maturity, and stress (Jung and Casler, 2006a, b).

Therefore, this research shows how the optimal activity of commercial FE depends more on changes in the structure and composition of NDF and its interaction with endogenous and exogenous bacteria than on the total ratio of NDF (Tirado-González *et al.*, 2018; 2021). Likewise, changes in the ratios of ADF, HEM, CEL, and ADL can improve the activity of FE. The relationships between cellulases and xylanases in commercial FEs depend on the fungus from which they were extracted, the type of substrate, and the extraction time. In addition, their action, combined with or independent from LAB, will be reflected in the pH —indicating greater production of acids during the fermentation of the silage (Skládanka *et al.*, 2012; Mora de Alba *et al.*, 2018; Li *et al.*, 2018)—, in the chemical composition of the silages at the end of fermentation (Gado *et al.*, 2013; Kholif *et al.*, 2017), in the subsequent fermentation (Tirado-Estrada *et al.*, 2015), and in the rumen degradability of NDF (Carrillo-Díaz *et al.*, 2022; Khan *et al.*, 2015).

This study shows how LD and MD of FE improved IVDMD24 in 60:40 O:T silages. However, their addition did not have positive effects in 80:20 O:T silages, partially showing the effects of interactions between the ratio and composition of NDF and the exogenous and endogenous enzymes and bacteria. This phenomenon is relevant because NDF degradability has previously been related to DM intake, as well as to the potential production and quality of ruminant milk and meat (Oba and Allen, 1999, 2000a, b; Arriola *et al.*, 2017; Tirado-González *et al.*, 2018, 2020). Likewise, changes in the productive behavior of ruminants are caused by changes in rumen kinetics —which depend on the type of successive populations of highly specific microorganisms that

Table 3. Pearson correlations among the following variables: composition, gas production (GP), and *in vitro* dry matter digestibility (IVDMD) at 24 h.

	S	Lag	IVDMD	PreE-DM	Ash	NDF	HEM	ADF	CEL	ADL	pH	CP	PostE-DM
Vmax	0.32**	-0.74***	0.66***	0.68***	-0.43***	-0.03	0.46***	-0.41***	-0.03	-0.44**	0.03	0.56***	0.68***
S		-0.50***	0.64***	0.43**	-0.09	-0.23§	0.24§	-0.45**	-0.30*	-0.31*	-0.17	0.61***	0.43**
Lag			-0.76***	-0.60***	0.21§	0.16	-0.38**	0.50***	0.22§	0.46***	-0.08	-0.62***	-0.60***
IVDMD				0.74***	-0.18	-0.21	0.48***	-0.62***	-0.37**	-0.47***	0.03	0.76***	0.74***
PreE-DM					-0.40**	-0.13	0.57***	-0.61***	-0.25*	-0.46***	0.09	0.77***	0.87***
Ash						0.06	-0.12	0.17	-0.40**	0.15	0.11	-0.33§	-0.40§
NDF							0.50***	0.70***	0.64***	0.22	0.01	-0.22	-0.13
HEM								-0.27*	-0.05	-0.38**	0.15	0.52***	0.57***
ADF									0.75***	0.56***	-0.12	-0.68***	-0.61***
CEL										0.07	-0.11	-0.36**	-0.25*
ADL											-0.16	-0.45***	-0.46***
pH												-0.14	0.11
CP													0.77***

* , ** , *** , statistical significances at P<0.001, P<0.01, P<0.05, respectively; Vmax, maximum volume of GP; S, fractional rate; Lag, lag phase; Initial DM, initial dry matter (0 d, prior to the ensiling process); ASH, ashes; NDF, neutral detergent fiber; HEM, hemicellulose; ADF, acid detergent fiber; CEL, cellulose; ADL, acid detergent lignin; CP, crude protein; IVDMD24, *in vitro* dry matter digestibility at 24 h; Final DM, final dry matter (DM of the silage at 40 d).

degrade the components of the NDF and non-fibrous carbohydrates of the diets, and on their efficient use to provide net production energy (Knapp *et al.*, 2014).

The use of LAB can contribute to the stability and quality of silages by promoting an increase in the lactic acid:acetic acid ratio (Guo *et al.*, 2013; Chen *et al.*, 2016; Mora de Alba *et al.*, 2018) and by limiting the growth of yeast in the silage (Tabacco *et al.*, 2011).

In this study, LD and MD of LAB improved the quality of the silages, based on the CP contents of the 60:40 O:T silages; in addition, they improved the IVDMD₂₄, V_{max}, S, and Lag of the 80:20 O:T silages. Lynch *et al.* (2012) and other authors have found that some strains and/or doses of *Lactobacillus* may not improve or that they may even worsen the quality of silage, as a consequence of the reduction in the concentration of lactic acid that can be used as a substrate by some lactic acid bacteria.

Finally, this study proves the relationship between IVDMD₂₄ and GP during *in vitro* fermentation with ruminal fluid. This relation is deeply connected with the changes in the ratios of hemicellulose and cellulose caused by the use of additives during the ensiling process. However, they are not always related to the total NDF content, which may not even be affected by the use of FE or LAB.

CONCLUSIONS

This work shows the degree of specificity with which enzyme preparations and lactic acid bacteria act on the different types of cell walls of different oat and triticale ratios in a silage. The contents of cell walls (NDF) and CP, *in vitro* degradability and fermentation were better in mixtures containing a lower ratio of oat. Although the additives did not consistently reduce NDF and ADF, their inclusion did affect the hemicellulose and cellulose ratios: the 60:40 O:T silages had more hemicellulose and less cellulose, ADL, and ASH than 80:20 silages, while silages with FE had a lower cellulose content than those treated with LAB. Although the use of additives does not directly improve the ratio of NDF during the ensiling process, it can increase the ratios of hemicellulose and reduce the ratios of cellulose. Hemicellulose and cellulose could be highly correlated with the fermentation and degradability of forages in the rumen.

REFERENCES

- AOAC. (2005). Official Methods of Analysis of AOAC International (OMA). AOAC International, Gaithersburg, MD.
- ANKOM Technol. (2016). Method 6, 20/01/16: Neutral Detergent Fiber in Feeds Filter Bag Technique (For A200, A200I). Ankom Technology, Macedon, N NY. Disponible en: [https://www.ankom.com/sites/default/files/document-files/Method 13 NDF Method A2000 RevE 4 10 15.pdf](https://www.ankom.com/sites/default/files/document-files/Method%2013%20NDF%20Method%20A2000%20RevE%204%2010%2015.pdf).
- Ajila, C. M., Sarma, S. J., Brar, S. K., Godbout, S., Cote, M., Guay, F., Verma, M., Valéro, J. R. (2015). Fermented apple pomace as a feed additive to enhance growth performance of growing pigs and its effects on emissions. *Agriculture* 5:313-329.
- Arriola, K.G., Oliveira, A.S., Ma, Z.X., Jean, I.J., Giurcanu, M.C., Adesogan A.T. (2017). A meta-analysis on the effect of dietary application of exogenous fibrolytic enzymes on the performance of dairy cows. *Journal of Dairy Science* 100(6):4513-4527.
- Beauchemin, K.A., Colombatto, D., Morgavi, D.P., Yang, W.Z. (2003). Use of the exogenous fibrolytic enzyme to improve feed utilization by ruminants. *Journal of Animal Science* 81: E37-E47.
- Carrillo-Díaz, M.I., Miranda-Romero, L.A., Chávez-Aguilar, G., Zepeda-Batista, J.L., González-Reyes, M., García-Casillas, A.C., Tirado-González, D.N., Tirado-Estrada, G. (2022). Improvement of ruminal

- neutral detergent fiber degradability by obtaining and using exogenous fibrolytic enzymes from White-Rot Fungi. *Animals* 12: 843. <https://doi.org/10.3390/ani12070843>
- Chen, L., Yuan, X., Li, J., Wang, S., Dong Z., Shao T. (2016). Effect of lactic acid bacteria and propionic acid on conservation characteristics, aerobic stability and *in vitro* gas production kinetics and digestibility of whole-crop corn based total mixed ration silage. *Journal of Integrative Agriculture* 15(7):1592-1600.
- Cobos, M.A., Yokoyama, M.T. (1995). Clostridium paraputrificum var ruminantum: colonization and degradation of shrimp carapaces in vitro observed by scanning electron microscopy. In: Wallas, R.J., Lahlou-Kassi, A. (Eds.), Rumen Ecology Research Planning, Proceedings of a Workshop. International Livestock Research Institute, Addis Abeba, Ethiopia, pp. 151-161.
- Del Valle, T.A., Antonio, G., Zenatti, T.F., Campana, M., Zilio, E.M., Ghizzi, L.G., Gandra, J.R., Osório, J.A.C., Morais, J.P.G. (2019). Effects of xylanase on the fermentation profile and chemical composition of sugarcane silage. *The Journal of Agricultural Science* 156(9):1123-1129.
- Gado, H.M., Salem, A.Z.M., Camacho, L.M., Elghandour, M.M.Y., Salazar M.C. (2013). Influence of exogenous enzymes on *in vitro* ruminal degradation of ensiled straw with DDGS. *Animal Nutrition and Feed Technology* 13:569-574.
- Guo, X.S., Undersander, D.J., Combs, D.K. (2013). Effect of Lactobacillus inoculants and forage dry matter on the fermentation and aerobic stability of ensiled mixed-crop tall fescue and meadow fescue. *Journal of Dairy Science* 96(3):1735-1744.
- Jung, H.G., Casler, M.D. (2006a). Maize stem tissues: cell wall concentration and composition during development. *Crop Science* 46:1793-1800.
- Jung, H.G., Casler, M.D. (2006b). Maize stem tissues: impact of development on cell wall degradability. *Crop Science* 46:1801-1809.
- Khan, N.A., Yu, P., Ali, M., Cone, J.W., Hendricks W.H. (2015). Nutritive value of maize silage in relation to dairy cow performance and milk quality. *Journal of the Science of Food and Agriculture* 95(2):238-252.
- Knapp, J.R., Laur, G.L., Vadas, P.A., Weiss, W.P., Tricarico J.M. (2014). Invited review: enteric methane in dairy cattle production: quantifying the opportunities and impact of reducing emissions. *Journal of Dairy Science* 97:3231-3261.
- Kholif, A.E., Elghandour, M.M.Y., Rodríguez, G.B., Olafadehan, O.A., Salem, A.Z.M. (2017). Anaerobic ensiling of raw agricultural waste with a fibrolytic enzyme cocktail as a cleaner and sustainable biological product. *Journal of Cleaner Production* 142:2649-2655.
- Li, J., Yuan, X., Dong, Z., Mugabe, W., Shao, T. (2018). The effects of fibrolytic enzymes, cellulolytic fungi and bacteria on the fermentation characteristics, structural carbohydrates degradation, and enzymatic conversion yields of *Pennisetum sinense* silage. *Bioresource Technology* 264:123-130.
- Lynch, J.P., Kiely, P.O., Waters, S.M., Doyle, E.M. (2012). Conservation characteristics of corn ears and stover ensiled with addition of *Lactobacillus plantarum* MTD-1, *Lactobacillus plantarum* 30114, or *Lactobacillus buchneri* 11A44. *Journal of Dairy Science* 95:2070-2080.
- McGinn, S.M., Beauchemin, K.A., Coates, T., Colombatto D. (2004). Methane emissions from beef cattle: effects of monensin, sunflower oil, enzymes, yeast, and fumaric acid. *Journal of Animal Science* 82:3346-3356.
- Menke, K.E., Steingass H. (1988). Estimation of the energetic feed value obtained from chemical analysis and *in vitro* gas production using rumen fluid. *Animal Research Development* 27:7-55.
- Mora de Alba, M.E., Tirado-González, D.N., Quezada-Tristán, T., Guevara-Lara, F., Jáuregui-Rincón, J., Larios-González, R., Tirado-Estrada, G. (2018). Calidad nutricional del bagazo de manzana ensilado con fuentes nitrogenadas orgánicas e inorgánicas. *Revista Mexicana de Ciencias Agrícolas* 9(1):229-235.
- Oba, M., Allen, M. (1999). Evaluation of the importance of the digestibility of NDF from forage: effects on dry matter intake and milk yield of dairy cows. *Journal of Dairy Science* 82:589-596.
- Oba, M., Allen, M. (2000a). Effects of brown midrib 3 mutation in corn silage on productivity of dairy cows fed two concentrations of dietary neutral detergent fiber: 1. Feeding behavior and nutrient utilization. *Journal of Dairy Science* 83:1333-1341.
- Oba, M., Allen, M. (2000b). Effects of brown midrib 3 mutation in corn silage on productivity of dairy cows fed two concentrations of dietary neutral detergent fiber: 2. Digestibility and microbial efficiency. *Journal of Dairy Science* 83:1350-1358.
- Phakachod, N., Suksombat, W., Colombatto, D., Beauchemin, K.A. (2013). Use of fibrolytic enzymes additives to enhance *in vitro* ruminal fermentation of corn silage. *Livestock Science* 157: 100-112.
- Petri, R.M., Schwaniger, T., Penner, G.B., Beauchemin, K.A., Forster, R.J., McKinnon, J.J., McAllister, T.A. (2013). Characterization of the core rumen microbiome in cattle during transition from forage to concentrate as well during and after an acidotic challenge. *PlosOne* 8(12):e83424.

- Salem, A.Z.M., G. Buendía-Rodríguez, G., Elghandour, M.M.M., Berasain, M.A.M., Jiménez, F.J.P., Pliego, A.B., Chagoyan, J.C.V., Cerrillo, M.A., Rodríguez, M.A. (2015). Effects of cellulase and xylanase enzymes mixed with increasing doses of *Salix babylonica* extract on *in vitro* rumen gas production kinetics of a mixture of corn silage with concentrate. *Journal of Integrative Agriculture* 14(1):131-139.
- Saleem, F., Ametaj, B.N., Bouatra, S., Mandal R., Zebeli, Q., Dunn, S.M., Wishart, D.S. (2012). A metabolomics approach to uncover the effects of grain diets on rumen health in dairy cows. *Journal of Dairy Science* 95: 6606-6623.
- Schofield, P., Pitt, R.E., Pell, A.N. (1994). Kinetics of fiber digestion from *in vitro* gas production. *Journal of Animal Science* 72:2980-2991.
- Shroeder, J.W. (2013). Silage fermentation and preservation. NDSU Extension Service: AS1254.
- Skládanka, J., Mikyska, F., Dolezal, P., Seda, J., Havlíček, Z., Mikel, O., Hosková, S. (2012). Effect of the technology of the additional sowing of drought-resistant clover-grass mixture and silage additives on fermentation process quality and nutritive value of baled grass silages. *African Journal of Agriculture Research* 7(2): 325-333.
- Tabacco, E., Piano, S., Revello-Chion, A., and Borreani, G. (2011). Effect of *Lactobacillus buchneri* LN4627 and *Lactobacillus buchneri* LN40177 on the aerobic stability, fermentation products, and microbial populations of corn silage under farm conditions. *Journal of Dairy Science* 94: 5589-5598.
- Tawari, S. P., Narag, M. P., Dubey, M. (2008). Effect of feeding apple pomace on milk yield and milk composition in crossbred (Red Sindhi × Jersey) cow. *Livestock Research for Rural Development* 4(29): 293-297.
- Tirado-Estrada, G., Mejía-Haro, I., Cruz-Vázquez, C.R., Mendoza-Martínez, G.D., Tirado-González, D.N. (2015). Degradación in situ y patrones de fermentación del rastrojo de maíz (*Zea mays* L.) tratado con enzimas exógenas en vacas Holstein. *Interciencia* 40(10): 716-724.
- Tirado-Estrada, G., Tirado-González, D.N., Medina-Cuéllar, S.E., Miranda-Romero, L.A., González-Reyes, M., Sánchez-Olmos, L.A., Castillo-Zúñiga, I. (2020). Global effects of maximizing the forage in production and quality of bovine milk and meat. A meta-analysis. *Interciencia* 45(10):461-468.
- Tirado-González, D.N., Tirado-Estrada, G., Miranda-Romero, L.A. (2015). Sobre el efecto de enzimas fúngicas en la alimentación de rumiantes. *Interciencia* 40(11):758-766.
- Tirado-González, D.N., Jauregui-Rincón J., Tirado-Estrada G., Martínez-Hernández, P.A., Guevara-Lara, F., Miranda-Romero L.A. (2016). Production of cellulases and xylanases by white-rot fungi cultured in corn stover media for ruminant feed applications. *Animal Feed Science and Technology* 221:147-156.
- Tirado-González, D.N., Miranda-Romero, L.A., Ruiz-Flores, A., Medina-Cuéllar, S.E., Ramírez-Valverde, R., Tirado-Estrada, G. (2018). Meta-analysis: effects of exogenous fibrolytic enzymes in ruminants. *Journal of Applied Animal Research* 46(1):771-783.
- Tirado-González, D.N., Tirado-Estrada, G., Miranda-Romero, L.A., Ramírez-Valverde, R., Medina-Cuéllar, S.E., Salem, A.Z.M. (2021). Effects of addition of exogenous fibrolytic enzymes on digestibility and milk and meat production- A Systematic Review. *Ann. Anim. Sci.* 21(4): 1159-1192.
- Van Soest, P.J., Robertson, J.B., Lewis, B.A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74: 3583-3592.