



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.



A Cross Sectional Study into Drug Abuse among Youth and Its Association with Socio-demographic and Risk Factors in District Kangra of Himachal Pradesh, India.

Madhur Katoch ^{a++}, Nitima ^{a#} and Raj Pathania ^{a##}

^a Department of Human Development and Family Studies, CSKHPKV, Palampur, Himachal Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAEES/2023/v41i41889

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97459>

Original Research Article

Received: 21/01/2023

Accepted: 23/03/2023

Published: 01/04/2023

ABSTRACT

Aim: Substance abuse has a detrimental impact on an individual, families and communities posing major public health challenges. Therefore, the present study has been designed to study the level of prevalence of drug abuse among youth and its association with socio demographic and various risk factors.

Methodology: The present research was conducted among 500 students in the age range of 12-25 years. The study was conducted in district Kangra of Himachal Pradesh, India. Out of total 15

⁺⁺ Assistant Professor;

[#] Research Assistant;

^{##} Professor & Head;

*Corresponding author: E-mail: madhur.katoch@gmail.com;

blocks from Kangra, five blocks were randomly selected. From these selected blocks five schools and five colleges (one school and one college from each block) were selected for data collection. "The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST)" and self-structured questionnaire were administered to the sample to collect data.

Results: It was revealed that tobacco was the most abused substance (29.2%) followed by alcohol (28.2%) and inhalants (11.8%). Less than 10 per cent of sample used other substances at least once in their lifetime. Moderate level of drug prevalence of drugs was found among study subjects. A highly significant association was observed between factors like age, type of educational institutes, mother's occupation, bad company, peer pressure, curiosity, and use of drugs by parents with prevalent drug abuse among youth.

Conclusion: It was concluded from the results that use of varied drugs like tobacco, alcohol, inhalants was found to be prevalent among youth. Moderate level of prevalence was reported for drugs like tobacco, cannabis, sedatives, opioids, inhalants and amphetamines. It is a matter of concern for them as they are at risk of health and other problems from their current pattern of substance use. A brief intervention has to be provided for them in terms of awareness and treatment from health professionals or by a specialist in drug and alcohol treatment service.

Keywords: Curiosity; peer pressure; risk factors; substance use/abuse; tobacco.

1. INTRODUCTION

Drug abuse is one of the most serious social problems we are facing these days. The World Health Organization [1] defines substance abuse as persistent, sporadic or hazardous use of psychoactive substances including alcohol and illicit drugs, whose repeated use can lead to dependence syndrome - a cluster of behavioral, cognitive, and physiological phenomena which involves a strong desire to take the drug, and difficulties in controlling its use. Drugs have the ability to change an individual's consciousness, perception, mood, thinking process, behaviour or motor function. Use of any substance once or twice may happen without having notable ramifications, but with repeated substance use the tolerance is built and it results in substance use disorder, where individual's functioning is effected at the expense of work, relationships, education, health or safety. Substance use disorders are marked as growing addiction or dependence on the substance and inability to restraint from it [2]. About 5 % of adult population globally uses drugs at least once during their life time. Whereas, 0.6 % of adult population suffers from disorders associated with drug use. According to The United Nations Office on Drugs and Crime (UNODC)'s World Drug Report [3] it was estimated that around 284 million people worldwide use drugs. The report further claims that India has one of the world's single-largest opiate markets. Youth in India are the most affected by this menace. Punjab state has become the hub of illicit drugs. About 60 % of all illicit drugs seized in India are from Punjab. Dependence of drug not only create economic

burden because of increasing costs of health care, but also social costs in the form of loss of productivity and family income, violence, security problems, traffic and workplace accidents. The regular use of intoxicating psychoactive substances leads to dependence which has not only an adverse effect on the physical and mental health of an individual but also disrupt his family life, social relationship and social development [4]. It has a detrimental impact on the society by increase in the crime rate like eve teasing, group clashes, assault, impulsive murder, stealing to pay for their drugs etc. Apart from affecting the financial stability, addiction increases conflicts and causes untold emotional pain for every member of the family. The injecting drug users are vulnerable to acquire HIV/AIDS, due to sharing of needles and risky sexual behaviour. Substance abuse is considered as a complicated and multidimensional problem. It is not entirely a problem of an individual person or any specific drug or a community, but is interplay between the triad [5]. The epidemic of substance abuse among youth has assumed alarming dimensions in India. Drug abuse in India is as old as elsewhere, if not older. Numerous references are well stocked in Ancient books regarding various intoxicants such as "soma rasa ", "dev booty", "madira" etc. Opium became popular during the Mughal period. The post-war period saw the rise of synthetic drugs-both stimulants and depressants. Hard drugs such as heroin and lysergic acid diethylamide (LSD) are in use. Recently discovered, hallucinogens such as phencyclidine hydrochloride that may be known as (Angel Dust) to certain users in metropolitan

areas. The traditional drugs like opium, charas, bhang and ganja were used by sections of the society partly as leisure time activity and partly as part of the religious ceremony. Therefore, their consumptions did not invite much negative sanction from the society. Abuse of alcoholic beverages and tobacco are endemic in many societies, whilst the abuse of other psychoactive substances present in epidemic features and drug abuse is transmitted from person to person like an infectious disease [6].

Substance use leads to definitive socio-economic burden and has become a major public health concern worldwide. In spite of realizing the adverse effects and repercussions of drug use, youth especially the adolescents have a proclivity to continue the habit [7]. Early initiation into substance/drug use is generally associated with a poor prognosis and a lifelong pattern of trickery and irresponsible behavior. There are multifarious causes of substance use disorder like genetic, poor impulse control, peer pressure, uninvolved parenting, neurological vulnerability to addiction or various mental health issues like anxiety or depression UNODC, [8]. Various other factors like breakdown of traditional values, peer pressure, curiosity, media, unemployment, industrialization and rural urban migration, availability of drugs have contributed to increase the number of drug abusers. With majority of drug users in the productive age group of 15-35 years, the loss in terms of human potential is immeasurable.

Himachal Pradesh is also becoming notorious with increasing number of drug abusers in the state. Cannabis can be growing naturally near the roadsides and even illicit cultivation of opium and cannabis in higher reaches of Kangra, Kullu, Mandi districts has become a matter of concern. Even Baddi in district Solan has emerged as pharmaceutical hub of India but few firms were caught for illegally producing synthetic drugs like adulterated heroin (called Chitta). Youth are getting hooked to 'chitta' as the peddlers are operating from small shops like ration or confectionery shops near educational institutions and chains are being formed by roping in new vulnerable targets especially young children. Illicit drug produced in the State attracts traffickers and as well as Indian and foreign tourists for consumption of narcotic drugs. 'Malana cream' produced from cannabis is grown in higher reaches of Kullu is a known brand in the international drug trafficking circles. Illicit uses of drugs in "Rave Parties or Full Moon Parties" are

also reported in some areas of Kullu and Kangra. Increase in cases of drug addiction is posing big challenge to human life, dignity, and law and order situation in the State. The present research was conducted with the objectives to study the prevalence of drug abuse among young population of Himachal Pradesh a northern state of India and to assess the association between drug abuse with various socio-demographic and risk factors.

2. METHODOLOGY

Locale of the study: The study was conducted in district Kangra of Himachal Pradesh. Out of 12 districts of Himachal Pradesh, Kangra district was purposively selected for the purpose of the study as it is the most populous district of Himachal Pradesh and it also shares its borders with Punjab state where drug abuse has acquired the proportions of a pestilence that has shaken the entire society of the state. This raging epidemic has its claws in Himachal where border areas are under the grip of this menace.

Research design: The study has adopted an as mixed method, exploratory cum descriptive research design. Under which prevalence of drug abuse among youth is explored using standardized tool and secondly the association is observed among socio demographic and risk factors with drug abuse.

Sample size: The sample for the study comprised of 500 respondents in the age range of 12-25 years.

Procedure for selection of subjects: Kangra district is further subdivided into 15 blocks. Therefore, out of 15, five blocks namely Nurpur, Shahpur, Bhawarna, Dharmashala and Dehra were selected randomly. The list of Senior Secondary schools and colleges has been procured from the portal of Education department of Himachal Pradesh. From the list 5 schools and 5 Colleges were randomly selected i.e. one school and one college each from selected blocks were identified for sample selection. From each school under the selected block 50 students in the age range of 12-18 years and similarly 50 students from each college in the age range of 19-25 years under the respective block were selected for collecting the data. Therefore, 250 students were selected from schools and 250 were selected from colleges, thus making a total of 500 sample size. The principals of these schools and colleges were

contacted through phone and prior permission was taken for data collection.

2.1 Tools Used for the Study

Background information Proforma: This is a self-structured proforma that comprises the socio demographic factors like name, age, gender, Name of school/college, Parental educational and occupational status, religion, caste, type of family, number of family members and Family income (Rs) of the respondent.

The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): was developed under the auspices of the World Health Organization (WHO)[9] by an international group of researchers and clinicians who work in the field of substance abuse and addiction. This scale was generated in lieu of the overwhelming public health concern associated with drug abuse globally. This questionnaire comprises of 8 item designed to be administered to a client using paper and pencil. This scale is culturally neutral and useable across a variety of cultures to screen for use of the following substances: tobacco products, alcohol, cannabis, cocaine, amphetamine-type stimulants (ATS), sedatives and sleeping pills (benzodiazepines), hallucinogens, inhalants and opioids.

General Information about drug awareness and abuse: This is a self-structured questionnaire where information is generated about the curiosity regarding drugs, awareness about drugs and problems associated with it, their attitude towards drug use, information about drug abuse in family or friends their association with bad company of friends etc.

Method of Data Collection: Primary data was collected through Questionnaire method in group setting. A group of 10 students each were given the questionnaire. The researchers gave brief introduction about the project. Consent is being taken and respondents were assured about the anonymity and confidentiality of their names along with data privacy. After seeking their approval in participating in the survey it was also notified that they can cancel their participation in survey if they want. After giving the introduction about the questionnaire, it was administered. It took about 20-30 minutes to complete the questionnaire. The researchers were present during the whole time the questionnaire was administered and any query sought was cleared there and then only. This survey followed the

Guidelines for Ethical Considerations in Social Research & Evaluation in India.

Data analysis: After the responses were generated, the information was tabulated and computed. The tables were quantified using frequency and percentages were calculated. Each drug use was measured by their levels in terms of their use i.e. (Low, Moderate, High). Data were further subjected to statistical analysis by using appropriate test. For measuring the association between socio demographic and risk factors with prevalence of drug abuse among youth chi square test was applied.

2.2 Math Formula

Chi- square test for independence of attributes in contingency tables: The chi-square test is a statistical procedure for determining the difference between observed and expected data. It is used to determine if there is a significant relationship between two nominal variables or not. The frequency of each category for one nominal variable is compared across the categories of the second nominal variable. The data of A and B displayed in 'x × y' contingency table, which contains 'xy' cell frequencies in 'x' rows represents a category for one variable (A) and 'y' columns represents a category for the other variable (B). e.g., if we want to examine the relationship between gender (male vs. female) and prevalence of drug abuse (low or high). The chi-square test of independence can be used to examine this relationship.

Test Statistic:

$$\chi^2 = \sum_{i=1}^x \sum_{j=1}^y \frac{(O_{ij} - E_{ij})^2}{E_{ij}} = \sum_{i=1}^x \sum_{j=1}^y \frac{O_{ij}^2}{E_{ij}} - N$$
, is distributed as χ^2 with (x-1) (y-1) is considered d.f.

Where, O_{ij} is the observed frequency in the (i, j) cell and E_{ij} be the expected frequency in the (i, j) cell.

N is the total frequency

d.f. is degree of freedom

3. RESULTS

Table 1 presents the socio demographic characteristics of the respondents. Among the respondents 37.6% are in the age group of 18-21 years, followed by 37.4% who belong to the age

range 15-17 years. About 12.6 per cent and 12.4 per cent belong to the age 12-14 years and 22-25 years respectively. Out of total sample of 500, 71.6% were males and remaining 28.4% were females. In case of type of educational institute 60 per cent of respondents and remaining 40 per cent were from government and private schools respectively. Among colleges 80 per cent of youth belong to government institutes and remaining 20 per cent were from private colleges. Regarding educational status of the fathers of respondents 24.4% had education upto graduation level followed by senior secondary i.e. 24.2 per cent. About 23.6 % were educated upto 10th standard. Only 10.4 per cent educated upto post graduate level and above. In case of father's occupational status of the respondent's majority i.e. 29.6% were working in private sector followed by 27.4% who were working in government sector and 16.4 percent had run own business as their occupation. Only 5 per cent of them were non-working and 13.6% were daily wage earner. About 8 per cent had agriculture as their occupation. Regarding mother's educational status, it was seen that majority of mother's i.e. 32.8 per cent had education upto 10th standard followed by graduation i.e. 20.6 per cent. Only 5.8 per cent were highly educated i.e. post graduate and above. Mother's occupational status reported that 74.2% were homemakers, 8.6% and 6.4% in government and private jobs respectively. Majority i.e. 97.4% was Hindus and 41.2% belongs to General category followed by OBC i.e. 28.6 per cent. About 20.2 % were belonging to Scheduled caste. Only 10 per cent were from Scheduled tribes. Majority i.e. 63.2 % was living in nuclear families and remaining 36.8 per cent belonged to joint families. In case of family income majority i.e.26.4 per cent had family income between Rs10,000 – Rs 25,000, followed by 24.6% who had income less than Rs 10,000, very few i.e. 6.8% had monthly income between Rs 75000 – 1lakh and above.

From Table 2 it can be reported that majority of students from both schools and colleges (29.2 %) used tobacco at least once in their lives. If we look further it can be seen that percentage of boys is higher as compared to girls. The use of tobacco is higher in college goers (34.4%) as compared to school goers (24.0%). In case of alcohol 28.2 per cent of students consumed alcohol at least one in their life time out of that 30.2% was boys and 23.2% were girls. About 20.4 per cent were school students out of those 21.6% were boys and 16.7% were girls. There is 6.6% intake of cannabis among total students,

where 8.4% were boys and 2.1 % were girls. The intake of cocaine in school girls (13.3%) is slightly more than boys (5.3%). Only 5.95 per cent of college boys and (6.1%) girls used cocaine once in their life. Amphetamine use in school students was reported to be 2 per cent whereas 2.8% of college students took amphetamines once in life. Regarding use of inhalants 12.8 per cent of school students and 10.8 per cent of college students use inhalants. In case of opioids use 5.2% of total students abused opioids at least once in their lifetime. Use of sedatives is little bit higher in college goers i.e. 7.2 percent as compared to school students i.e. 3.2 per cent.

Table 3 showed the levels of prevalence of drug abuse among selected sample. It can be seen from the table that majority of the school sample including both boys and girls never used any type of drugs. As for different drugs 98% of students never used opioids, amphetamines and 97.6 per cent never used cannabis, whereas 87.2 per cent never used inhalants. About 76 per cent never use tobacco followed by 79.6% who never consumed alcohol. The ones who consumed these different drugs were divided into three levels i.e. Low, Moderate and High according to their usage. As we go further into the table for respective drugs in case of school sample we observed that about 14.8% of school boys have fallen under moderate level of risk caused by tobacco. A few numbers of boys i.e. (4.4%) have low level of risk. Similarly, school girls (2.8%) were also fallen under moderate level of risk for tobacco. Whereas the number of boys (12.4%) and girls (3.6%) were in low level of risk caused by alcohol followed by moderate level. A very few number of both school boys and girls have scored higher in moderate level of risk caused by cannabis, cocaine, amphetamine, inhalants, opioids and sedatives followed by boys and girls under low level of risks.

As we delve further in category of college students it can be seen that 40.4% of boys and (25.2% & 23.6%) of girls never used tobacco and alcohol in their life while maximum number of college boys and girls did not used other drugs such as cannabis, cocaine, amphetamine, inhalants and opioids and sedatives. From the table, it is clear that the number of boys and girls for the drugs such as tobacco (19.6% & 5.2%), cocaine (2% & 1.2%), amphetamine (0.8% & 0.8%), inhalants (3.6% & 3.2%), opioids (1.6% & 0.4%) and sedatives (2.4% & 1.2%) respectively were under moderate level of risks followed by

Table 1. Frequency and percentage distribution of socio-demographic factors among respondents

Variables		Sample(N=500)	Percentage (%)
Age			
12-14		63	12.6
15-17		187	37.4
18-21		188	37.6
22-25		62	12.4
Gender			
Male		358	71.6
Female		142	28.4
Type of educational institute			
School		250	50.0
College		250	50.0
Type of institute			
School	Govt. School	150	60.0
(n=250)	Private School	100	40.0
College	Govt. College	200	80.0
(n=250)	Private College	50	20.0
Father educational status			
Illiterate		5	1.0
Primary		14	2.85
Middle		68	13.6
Matric		118	23.6
Sr. Sec		121	24.2
Graduate		122	24.4
Post Graduate & above		52	10.4
Father Occupational Status			
Govt.		137	27.4
Private		148	29.6
Business		82	16.4
Agriculture		40	8
Daily wage earner		68	13.6
Non-working		25	5
Mother educational status			
Illiterate		10	2.0
Primary		26	5.2
Middle		73	14.6
Matric		164	32.8
Sr. Sec		95	19
Graduate		103	20.6

Variables	Sample(N=500)	Percentage (%)
Post Graduate & above	29	5.8
Mother Occupational Status		
Govt.	43	8.6
Private	32	6.4
Business	15	3.0
Agriculture	15	3.0
Daily wage earner	24	4.8
Home maker	371	74.2
Religion		
Hindu	487	97.4
Sikh	10	2.0
Muslim	3	0.6
Caste Category		
General	206	41.2
SC	101	20.2
ST	50	10.0
OBC	143	28.6
Type of family		
Nuclear	316	63.2
Joint	184	36.8
Family income (annual) in rupees		
Less than 10,000	123	24.6
10,000-25000	132	26.4
25,000-50000	103	20.6
50000-75000	74	14.8
75000-1 Lakh	34	6.8
1 Lakh and above	34	6.8

Table 2. Frequency and percentage distribution of overall substance use prevalence among respondents

Drugs		School (n=250)			College (n=250)			Total (%) N=500		
		Boys(%)	Girls(%)	Total(%)	Boys(%)	Girls(%)	Total(%)	Boys(%)	Girls(%)	Total(%)
Tobacco	Users	48(25.3)	12(20.0)	60(24.0)	67(39.9)	19(23.2)	86(34.4)	115(32.2)	31(21.8)	146(29.2)
	Non Users	142(74.7)	48(80.0)	190(76.0)	101(60.1)	63(76.8)	164(65.6)	243(67.8)	111(78.2)	354(70.8)
Alcohol	Users	41(21.6)	10(16.7)	51(20.4)	67(39.9)	23(28.05)	90(36.0)	108(30.2)	33(23.2)	141(28.2)
	Non Users	149(78.4)	50(83.3)	199(79.6)	101(60.1)	59(71.95)	160(64.0)	250(69.8)	109(76.8)	359(71.8)
Cannabis	Users	6(3.2)	0.0	6(2.4)	24(14.3)	3(3.65)	27(10.8)	30(8.4)	3(2.1)	33(6.6)
	Non Users	184(96.8)	60(100.0)	244(97.6)	144(85.7)	79(96.35)	223(89.2)	328(91.6)	139(97.9)	467(93.4)
Cocaine	Users	10(5.3)	8(13.3)	18(7.2)	10(5.95)	5(6.1)	15(6.0)	20(5.6)	13(9.2)	33(6.6)
	Non Users	180(94.7)	52(86.7)	232(92.8)	158(94.05)	77(93.9)	235(94.0)	338(94.4)	129(90.8)	467(93.4)
Amphetamine	Users	1(0.5)	4(6.7)	5(2.0)	4(2.4)	3(3.6)	7(2.8)	5(1.4)	7(4.92)	12(2.4)
	Non Users	189(99.5)	56(93.3)	245(98.0)	164(97.6)	79(96.4)	243(97.2)	353(98.6)	135(95.08)	488(97.6)
Inhalants	Users	20(10.5)	12(20.0)	32(12.8)	16(9.5)	11(13.4)	27(10.8)	36(10.06)	23(16.2)	59(11.8)
	Non Users	170(89.5)	48(80.0)	218(87.2)	152(90.5)	71(86.6)	223(89.2)	322(89.94)	119(83.8)	441(88.2)
Opioids	Users	5(2.6)	0.0	5(2.0)	11(6.5)	3(3.6)	14(5.6)	16(4.5)	3(2.1)	19(3.8)
	Non Users	185(97.4)	60(100.0)	245(98.0)	157(93.5)	79(96.4)	236(94.4)	342(95.5)	139(97.9)	481(96.2)
Sedatives	Users	4(2.1)	5(8.3)	8(3.2)	14(8.3)	4(4.9)	18(7.2)	18(5.03)	9(6.3)	26(5.2)
	Non Users	186(97.9)	55(91.7)	242(96.8)	154(91.7)	78(95.1)	232(92.8)	340(94.97)	133(93.7)	474(94.8)

low level of risk except for alcohol. For alcohol both college sample (13.2% & 7.2%) were found to have low level of risk followed by moderate level. As we go further 3.6 % of tobacco users 2.4% of alcohol users, 1.6% cannabis users, 0.8 % cocaine and 0.4 % college goers were in high level of risk and need intensive intervention by health professional.

The number of total participants in all types of drugs falls under moderate level of risk and are prevalent to drug abuse except for alcohol. Only for alcohol 18.2% of respondents were at low level where only brief intervention may be required. There are a very few number of respondents who are in high risk and need intervention to cure. For tobacco (1.8%), alcohol (1.6%), followed by cannabis (1.0%).

Table 4 observed the association between socio-demographic factors and prevalence of drug abuse among study subjects. As seen from the table age of respondent's $P > 0.01$ (p value is .001). Therefore, extremely significant association at 1% level of significance was found between age and prevalence of drug abuse among the selected respondents. Further it can be corroborated from the table that the type of institutes i.e. schools and colleges also were highly significantly associated at 5 % level of significance with the prevalence of drug /substance abuse among the respondents, where $P > 0.05$ (p value is .027) If we look into the table further it was seen that highly significant association at 5 % level of significance was observed between mother's occupational status and prevalence of drug /substance abuse among their children, $P > 0.05$ (p value is .029). No significant associations were observed between variables like Gender, Type of schools and college, father's education and occupation and mother's education with prevalence of drug abuse among children.

Table 5 postulated the association between risk factors and prevalence of drug abuse among subjects studied. As per table the curiosity in participants has $P > 0.05$ (p value is .021) which means that highly significant association at 5% level of significance was found between curiosity and prevalence of drug abuse among selected sample. If we look further into the table, peer pressure and prevalence of drug abuse in respondents has $P > 0.01$ (p value is .001). Therefore, extremely significant association at 1 % level of confidence was observed between peer pressure and prevalence of drug abuse among study subjects.

From above Table 5, it can be corroborated that the parental use of substances (i.e. alcohol and tobacco) of selected sample and prevalence of drug abuse has $P > 0.05$ (p value is .024) which depicts highly significant association between parent's substance use and prevalence of drug abuse at 5% level of significance. As seen from the table that in case of children attitude towards drug use and prevalence of drug abuse in them $P > 0.01$ (p value is .002) has been extremely significantly associated with each other at 1 % level of confidence. If we look further into the table, the respondents who find themselves in bad company and prevalent to drug abuse has $P > 0.01$ p value is (.000) which means that extremely significant association at 1% level of significance was found between bad company and prevalence of drug abuse. No significant association is observed between lack of awareness and prevalence of drug abuse among students.

4. DISCUSSION

This article aims to shed light on the prevalence of drug/substance use among students 12-25 years old from the 5 blocks of district Kangra, of state Himachal Pradesh in India. These findings will help to understand the prevalent drug use among sample along with associated risk factors and also the association of various socio-demographic factors with drug abuse. The findings reported that number of male students were higher as compared to female students in the selected sample. The age group of study subjects was 12-25 years, where majority belonged to 15-21 years. The sample had more number of students who were studying in Government institutes. The education level of fathers and mothers of the respondents were graduates and up to matric level respectively. In case of occupation status majority of fathers of respondents were doing private jobs and mothers were homemakers. Majority of study subjects were Hindus in General category mostly living in nuclear families. Majority of respondent's family income was between Rs 10,000-25,000.

Regarding the overall substance/drug use and level of prevalence among study subjects, majorly used substance among youth is found to be tobacco followed by alcohol. Myers and Kelly [10], reported that cigarette and alcohol use often develop simultaneously, and smoking is especially common among youth. Various studies suggested that tobacco use is closely associated with Alcohol and other drug (AOD).

Table 3. Frequency and percentage distribution of levels of prevalence of drug abuse among respondents

Variables	Sample	Levels	Tobacco	Alcohol	Cannabis	Cocaine	Amphetamine	Inhalants	Opioids	Sedatives
School (n=250)	Boys (%)	Never	142(56.8)	149(59.6)	184(73.6)	180(72.0)	189(75.6)	170(68.0)	185(74.0)	186(74.4)
		Low	11(4.4)	31(12.4)	2(0.8)	1(0.4)	1(0.4)	3(1.2)	0.0	1(0.4)
		Moderate	37(14.8)	8(3.2)	3(1.2)	8(3.2)	0.0	16(6.4)	5(2.0)	3(1.2)
		High	0.0	2(0.8)	1(0.4)	1(0.4)	0.0	1(0.4)	0.0	0.0
	Girls (%)	Never	48(19.2)	50(20.0)	60(24.0)	52(20.8)	56(22.4)	48(19.2)	60(24.0)	56(22.4)
		Low	5(2.0)	9(3.6)	0.0	1(0.4)	0.0	2(0.8)	0.0	3(1.2)
		Moderate	7(2.8)	1(0.4)	0.0	7(2.8)	4(1.6)	10(4.0)	0.0	1(0.4)
		High	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Total (%)	Never	190(76.0)	199(79.6)	244(97.6)	232(92.8)	245(98.0)	218(87.2)	245(98.0)	242(96.8)
		Low	16(6.4)	40(16.0)	2(0.8)	2(0.8)	1(0.4)	5(2.0)	0.0	4(1.6)
		Moderate	44(17.6)	9(3.6)	3(1.2)	15(6.0)	4(1.6)	26(10.4)	5(2.0)	4(1.6)
		High	0.0	2(0.8)	1(0.4)	1(0.4)	0.0	1(0.4)	0.0	0.0
College (n=250)	Boys(%)	Never	101(40.4)	101(40.4)	144(57.65)	158(63.2)	164(65.6)	152(60.8)	157(62.8)	154(61.6)
		Low	10(4.0)	33(13.2)	6(2.4)	3(1.2)	1(0.4)	6(2.4)	6(2.4)	7(2.8)
		Moderate	49(19.6)	29(11.6)	15(6.0)	5(2.0)	2(0.8)	9(3.6)	4(1.6)	6(2.4)
		High	8(3.2)	5(2.0)	3(1.2)	2(0.8)	1(0.4)	1(0.4)	1(0.4)	1(0.4)
	Girls(%)	Never	63(25.2)	59(23.6)	79(31.6)	77(30.8)	79(31.6)	71(28.4)	79(31.6)	78(31.2)
		Low	5(2.0)	18(7.2)	2(0.8)	3(1.2)	1(0.4)	3(1.2)	1(0.4)	1(0.4)
		Moderate	13(5.2)	4(1.6)	0.0	2(0.8)	2(0.8)	8(3.2)	2(0.8)	3(1.2)
		High	1(0.4)	1(0.4)	1(0.4)	0.0	0.0	0.0	0.0	0.0
	Total(%)	Never	164(65.6)	160(64.0)	223(89.2)	235(94.0)	243(97.2)	223(89.2)	236(94.4)	232(92.8)
		Low	15(6.0)	51(20.4)	8(3.2)	6(2.4)	2(0.8)	9(3.6)	7(2.8)	8(3.2)
		Moderate	62(24.8)	33(13.2)	15(6.0)	7(2.8)	4(1.6)	17(6.8)	6(2.4)	9(3.6)
		High	9(3.6)	6(2.4)	4(1.6)	2(0.8)	1(0.4)	1(0.4)	1(0.4)	1(0.4)
Total (N=500)		Never	354(70.8)	359(71.8)	467(93.4)	467(93.4)	488(97.6)	441(88.2)	481(96.2)	474(94.8)
		Low	31(6.2)	91(18.2)	10(2.0)	8(1.6)	8(1.6)	14(2.8)	7(1.4)	12(2.4)
		Moderate	106(21.2)	42(8.4)	18(3.6)	22(4.4)	22(4.4)	43(8.6)	11(2.2)	13(2.6)
		High	9(1.8)	8(1.6)	5(1.0)	3(0.6)	3(0.6)	2(0.4)	1(0.2)	1(0.2)

Table 4. Association between socio-demographic factors and prevalence of drug abuse among sample

Factors	Components	Prevalence					Value	Df	p
		N=500 (%)	Never(%)	Low(%)	Moderate(%)	High(%)			
Age	12-14 years	63 (12.6)	29(5.8)	13(2.6)	19(3.8)	2(0.4)	27.680	9	.001***
	15-17 years	187(37.4)	115(23.0)	21(4.2)	50(10.0)	1(0.2)			
	18-21 years	188(37.6)	91(18.2)	26(5.2)	65(13.0)	6(1.2)			
	22-25 years	62(12.4)	21(4.2)	11(2.2)	24(4.8)	6(1.2)			
Gender	Male	358(71.6)	186(37.2)	49(9.8)	109(21.8)	14(2.8)	4.389	3	.222
	Female	142(8.4)	70(14.0)	22(4.4)	49(9.8)	1(0.2)			
Type of educational institute	School	250(50.0)	143(28.6)	33(6.6)	70(14.0)	4(0.8)	9.185	3	.027**
	College	250(50.0)	113(22.6)	38(7.6)	88(17.6)	11(2.2)			
Type of school (n=250)	Government	150(60.0)	84(33.6)	23(9.2)	40(16.0)	3(1.2)	1.484	3	.686
	Private	100(40.0)	58(23.2)	11(4.4)	30(12.0)	1(0.4)			
Type of college (n=250)	Government	200(80.0)	88(35.2)	32(12.8)	69(27.6)	11(4.4)	3.228	3	.358
	Private	50(20.0)	23(9.2)	7(2.8)	20(8.0)	0.0			
Father education	Illiterate	5(1.0)	3(0.6)	0.0	2(0.4)	0.0	25.696	18	.107
	Primary	14(2.8)	1(0.2)	6(1.2)	7(1.4)	0.0			
	Middle	68(13.6)	37(7.0)	5(1.0)	23(4.6)	3(0.6)			
	Matric	118(23.6)	62(12.4)	13(2.6)	38(7.6)	5(1.0)			
	Sr Secondary	121(24.2)	63(12.6)	16(3.2)	38(7.6)	4(0.8)			
	Graduate	122(24.4)	67(13.4)	21(4.2)	32(6.4)	2(0.4)			
	Post Graduate & Above	52(10.4)	23(4.6)	10(2.0)	18(3.6)	1(0.2)			
Father occupation	Government	137(27.4)	74(14.8)	21(4.2)	38(7.6)	4(0.8)	14.448	15	.492
	Private	148(29.6)	70(14.0)	27(5.4)	47(9.4)	4(0.8)			
	Business	82(16.4)	47(9.4)	6(1.2)	28(5.6)	1(0.2)			
	Agriculture	40(8.0)	19(3.8)	4(0.8)	14(2.8)	3(0.6)			
	Daily wage earner	68(13.6)	34(6.8)	11(2.2)	20(4.0)	3(0.6)			
	Non-working	25(5.0)	12(2.4)	2(0.4)	11(2.2)	0.0			
Mother education	Illiterate	10(2.0)	4(0.8)	1(0.2)	5(1.0)	0.0	11.737	18	.861
	Primary	26(5.2)	13(2.6)	5(1.0)	8(1.6)	0.0			
	Middle	73(14.6)	40(8.0)	8(1.6)	24(4.8)	1(0.2)			
	Matric	164(32.8)	87(17.4)	20(4.0)	49(9.8)	8(1.6)			
	Sr Secondary	95(19.0)	45(9.0)	14(2.8)	34(6.8)	2(0.4)			
	Graduate	103(20.6)	53(10.6)	19(3.8)	29(5.8)	2(0.4)			
	Post Graduate & Above	29(5.8)	14(2.8)	4(0.8)	9(1.8)	2(0.4)			
Mother occupation	Government	43(8.6)	22(4.4)	7(1.4)	12(2.4)	2(0.4)	26.949	15	.029**
	Private	32(6.4)	12(2.4)	7(1.4)	12(2.4)	1(0.2)			

Factors	Components	Prevalence					Value	Df	p
		N=500 (%)	Never(%)	Low(%)	Moderate(%)	High(%)			
	Business	15(3.0)	7(1.4)	3(0.6)	4(0.8)	1(0.2)			
	Agriculture	15(3.0)	6(1.2)	3(0.6)	3(0.6)	3(0.6)			
	Daily wage earner	24(4.8)	9(1.8)	6(1.2)	9(1.8)	0.0			
	Home-Maker	371(74.2)	200(40.0)	45(9.0)	118(23.6)	8(0.4)			
Religion	Hindu	487(97.4)	249(49.8)	70(14.0)	153(30.6)	15(3.0)	4.772	6	.573
	Sikh	10(2.0)	5(1.0)	0.0	5(1.0)	0.0			
	Muslim	3(0.6)	2(0.4)	1(0.2)	0.0	0.0			
Caste category	General	206(41.2)	105(21.0)	28(5.6)	67(13.4)	6(1.2)	2.569	9	.979
	Schedule Caste	101(20.2)	54(10.8)	11(2.2)	32(6.4)	4(0.8)			
	Schedule Tribe	50(10.0)	24(4.8)	9(1.8)	16(3.2)	1(0.2)			
	OBC	143(28.6)	73(14.6)	23(4.6)	43(8.6)	4(0.8)			
Family Type	Nuclear	316(63.2)	165(33.0)	47(9.4)	94(18.8)	10(2.0)	1.458	3	.692
	Joint	184(36.8)	91(18.2)	24(4.8)	64(12.8)	5(1.0)			
Family income	Less than 10,000	123(24.6)	63(12.6)	17(3.4)	41(8.2)	2(0.4)	20.981	15	.137
	10,000-25,000	132(26.)	73(14.6)	20(4.0)	33(6.6)	6(1.2)			
	25,000-50,000	103(20.6)	54(10.8)	13(2.6)	35(7.0)	1(0.2)			
	50,000-75,000	74(14.8)	37(7.4)	6(1.2)	30(6.0)	1(0.2)			
	75,000-1 Lakh	34(6.8)	15(3.0)	6(1.2)	11(2.2)	2(0.4)			
	1 Lakh & above	34(6.8)	14(2.8)	9(1.8)	8(1.6)	3(0.6)			

Significant level 99%***95%** 90%*

Table 5. Association between risk factors and prevalence of drug abuse among sample

Risk Factors (N=500)	Curiosity		Peer Pressure		Substance use by parents (Alcohol and Tobacco)		Attitude towards drug use		Lack of awareness		Bad company	
	1(%)	0(%)	1(%)	0(%)	1(%)	0(%)	1(%)	0(%)	1(%)	0(%)	1(%)	0(%)
Never	198(39.6)	58(11.6)	225(45.0)	31(6.2)	204(40.8)	52(10.4)	154(30.8)	102(20.4)	196(39.2)	60(12.0)	248(49.6)	8(1.6)
Low	47(9.4)	24(4.8)	54(10.8)	17(3.4)	48(9.6)	23(4.6)	32(6.4)	39(7.8)	56(11.2)	15(3.0)	64(12.8)	7(1.4)
Moderate	101(20.2)	57(11.4)	118(23.6)	40(8.0)	124(24.8)	34(6.8)	72(14.4)	86(17.2)	126(25.2)	32(6.4)	144(28.8)	14(2.8)
High	11(2.2)	4(0.8)	9(1.8)	6(1.2)	8(1.6)	7(1.4)	4(0.8)	11(2.2)	11(2.2)	4(0.8)	9(1.8)	6(1.2)
Total	357(71.4)	143(28.6)	406(81.2)	94(18.8)	384(76.8)	116(23.2)	262(52.4)	238(47.6%)	389(77.8)	111(22.2)	465(93.0)	35(7.0)
Chi-square	9.722		17.549		9.453		14.642		.794		32.729	
Df	3		3		3		3		3		3	
P value	.021**		.001***		.024**		.002***		.851		.000***	

Significant level 99%*** 95%**

The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): developed by World Health Organization (WHO)[9] gave level of drug abuse in terms of Low, Moderate and High. From the results it was observed that tobacco was the most consumed drug among youth and majority fall in moderate level of consumption both in case of boys and girls from schools and colleges. But in case of alcohol majority were in low level of use the rest of the users fell under moderate level of drug use. It is studied that use of psychoactive substances significantly escalates during the high school years especially 14–18 years old according to Johnston et al. [11]. Another study conducted by Sivapuram et al. [12] on prevalence of alcohol and tobacco use in India found that the prevalence of alcohol abuse (8.7%) was higher than the tobacco abuse (7.9%). When compared among the genders, both alcohol and tobacco consumption was higher among males (15.8% alcohol and 13.1% tobacco) as compared to females (3.2% tobacco and 2.4% alcohol). The prevalence of smoking among 13,329 respondents (44.9% males and 55.1% females) under cross-sectional GATS-2survey in India, aged 15 to 24 years found that overall, 11.9% of respondents were using tobacco [13]. Contrast to our study findings in one study alcohol was the most common substance used by abusers (95.4%), followed by tobacco (46.5%) but the age of participants in this study were 30-50 years. It is further discussed that tobacco acts as gateway drug for initiation for the use and abuse of other substances, because those who smoke are 3 times more likely than non-smokers to use alcohol, marijuana and cocaine[14]. Every year, approximately 55,000 children start using tobacco generally hailing from low socio-economic status. This behavior is often initiated during among young population as 70% of adult smokers reported that they started smoking daily prior to age of 18 years [15]. The third most abused drug comes under inhalants (11.8%) of students were found using inhalants. It is also clear from the study that inhalant abuse is not uncommon among youth. Inhalants like glue, turpentine thinners, paints are commonly used by adolescents and youth because of their easy availability; Sadock and Sadock [16]; Dhawan and Pattanayak [17]; Narayanaswamy et al. [18]. Inhalants cause a momentary sense of well-being that reinforces repetition among users. Their effects generally last only a few minutes, where user experience exhilaration and a fleeting phase of tranquility. According to the National

Drug Dependence Treatment Centre, AIIMS, New Delhi's 2019 conducted National Survey on Extent and Pattern of Substance Use in India, the prevalence of inhalant misuse is 0.7% nationwide [19]. The percentage of cannabis, cocaine, opioids and amphetamine user was less than 10 per cent but the level of prevalence for these drugs was found at moderate level, which is quite alarming.

A significant association was found between socio-demographic variable like age, type of educational institute such as school and college and mother's occupation with prevalence of drug abuse have significant association. In a similar study conducted by Gordon et al. [20] results showed that age of the respondents was significantly associated with substance use, where youth being more likely to engage in substance use ($P < 0.001$); having a parent/guardian employed were negatively associated with substance use ($P = 0.021$). The results also aligned with study of Mahmood et al. [21] who found that age group of 17-19 years of male adolescents were found to be significantly associated with tobacco intake or cigarette smoking. In another study of Mohammadpoorasl et al. [22], increasing age of the students was significantly associated with substance use. The reason behind this could be the easy accessibility of substances with increasing age. In contradictory to our findings age was not statistically significant on drug abuse among the students indicating that abuse of drugs takes place across any age. The result showed that there was no significant influence of age on drug abuse among undergraduate students of Benue State University, Makurdi. This is an indication that students at whatever age exhibit the same level of drug abuse among undergraduate students in Benue State University, Makurdi [23]. According to Gudaji et al. [24] study, found that being of younger age, being single, and having a father who smoked were significantly associated with psychoactive substance use. Participants who were single were more likely to use inhalants, opiates, cannabis and stimulants. No relationship was observed between the age of the respondents and specific substance use. In another study prevalence of use of substance was found to be 1.3 times more among boys from the government schools in comparison to private school ($P < 0.05$), whereas alcohol use was 1.5 times more among girls from government schools in comparison to girls from private schools ($P < 0.05$); Narain et al. [25].

Association of prevalent substance use was found to be significantly associated with curiosity to use certain drugs, peer pressure, parental use of substances like alcohol and tobacco, attitude towards drugs and bad company. Similar results are found in the study of Mahmood et al. [21] who revealed significant association between student's cigarette and alcohol consumption with parental use of substance like tobacco and alcohol. The study of Rukundo et al.[26] confirmed that peer influence is the strongest risk factor associated with substance use in school students. Gudaji et al. [24] in their study showed that the fathers who smoked were significantly associated with psychoactive substance use. Webetu et al. [27] reported that children who had family members or peers exposed to substance use were more likely to use substance/drugs as compared to those whose family members are non-users. The odds of experiencing lifetime legal substance use were 2.5 times higher among students, who had a substance user family than those who don't have substance user in the family. Whitesell et al. [28] reported that having a father, mother or siblings who are substance users increased the odds of cigarette smoking by 1.5 times, 5 times for alcohol consumption, compared with those who did not have a substance user in the family. Family members, especially parents act as a role model for children and this creates indirectly the impression that substance use is acceptable behaviour among adults.

5. CONCLUSION

The study concluded that tobacco was the main drug used by the youth followed by alcohol and inhalants. Moderate level of drug use was found among majority of study subjects. Socio demographic factors like age, type of educational institute like schools and colleges and mother's occupation was found to be associated with prevalent drug use. Various risk factors like peer pressure, curiosity and parental use of drugs were found associated with use of substances among them. Strict measures should be taken by the police regarding prohibition of these substances nearby to school and college premises. Educational awareness pertaining to the harmful effects of substance use should be addressed in schools and colleges along with mandatory units regarding drug abuse in school curriculums starting from 6th standard. Awareness programs need to be conducted at regular intervals at community level to address the issues of drug abuse.

CONSENT

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

This work is supported by the Indian Council of Social Science Research (ICSSR), New Delhi for the year 2022-24.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization-Regional Office for South-East Asia. Current information on use and harm from alcohol in the South East Asian Region. Alcohol control series 6. New Delhi: WHO-SEARO; 2007. ISBN 978-92-9022-246-0
Available:<https://apps.who.int/iris/handle/10665/204906>
2. American Psychiatric Association (APA). Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Arlington, VA: American Psychiatric Publishing; 2013.
Available:<https://10.1176/appi.books.9780890425596>
3. United Nations Office on Drugs and Crime (UNODC)- World Drug Report. Vienna: United Nations Publications; 2022.
Available:<https://www.unodc.org/unodc/frontpage/2022/June/unodc-world-drug-report-2022>
4. Kumar N, Kanchan T, Unnikrishnan B, Thapar R, Mithra P. Profile of substance use among patients attending de-addiction centers in a coastal city of Southern India. Plos One. 2013;8(2):1-4.
5. Ortiz A. Development of a system for registry of information on drug use in Mexico. Bulletin of the Pan American Health Organization. 1990;24(1):46-52.
6. Medina-Mora ME, Tapia CR, Rascón ML, Solache G, Otero BR, Lazcano F, Mariño

- MC. Epidemiologic status of drug abuse in Mexico. Bulletin of the Pan American Health Organization. 1990;24(1):1-11.
7. Tsering D, Pal R, Dasgupta A. Substance use among adolescent high school students in India: A survey of knowledge, attitude, and opinion. Journal of Pharmacy and Bio Allied Sciences. 2020;2(2):137-40.
Available: <https://doi.org/10.4103/0975-7406.67005>.
8. United Nations Office on Drugs and Crime (UNODC)- World Drug Report. Vienna: United Nations Publications; 2018. ISBN: 978-92-1-148304-8; eISBN: 978-92-1-045058-4
9. WHO ASSIST Working Group. The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): Development, Reliability and Feasibility. Addiction. 2002; 97:1183-94.
10. Myers MG, Kelly JF. Cigarette smoking among adolescents with alcohol and other drug use problems. Alcohol Research & Health: The Journal of the National Institute on Alcohol Abuse and Alcoholism. 2006;29(3):221-7.
11. Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. Monitoring the future national results on adolescent drug use: Overview of key findings. National Institute on Drug Abuse, U. S. Dept. of Health and Human Services. (Bethesda) Maryland: National Institutes of Health; 2006. NIH publication no. 06-5882
12. Sivapuram MS, Nagarathna R, Anand A, Patil S, Singh A, Nagendra HR. Prevalence of alcohol and tobacco use in India and implications for COVID-19-Niyantrita Madhumeha Bharata Study Projections. Journal of Medicine and Life. 2020;13(4):499-09.
Available: <https://doi.org/10.25122/jml-2020-0079>.
13. Grover S, Anand T, Kishore J, Tripathy JP, Sinha DN. Tobacco use among the youth in India: Evidence from global adult tobacco survey-2 (2016-2017). Tobacco Use Insights. 2020;13:1-7.
Available:<http://dx.doi.org/10.1177/1179173X20927397>
14. Sims TH, Committee on Substance Abuse. From the American Academy of Pediatrics: Technical report- Tobacco as a substance of abuse. Pediatrics. 2009;124(5):1045-53.
Available:<https://doi.org/10.1542/peds.2009-2121>
15. Ghosh G. Substance abuse among young people in India - Approaches at curbing the Menace; 2013. Retrieved from Available:<http://dx.doi.org/10.2139/ssrn.2235028> or Available:<https://ssrn.com/abstract=2235028>
16. Sadock BJ, Sadock VA. Kaplan and Sadock's Synopsis of Psychiatry-Behavioral sciences/clinical psychiatry, 10th ed. Indian Journal of Psychiatry. 2009;51(4):331.
17. Dhawan A, Pattanayak RD. Synopsis of the clinical practice guidelines on management of inhalant use disorders. In: P. K. Dalal & D. Basu, editors. Clinical Practice Guidelines for Assessment and Management of Substance Use Disorders. New Delhi: Indian Psychiatric Society 2015:91-105.
18. Narayanaswamy J, Viswanath B, Ravi M, Muralidharan K. Inhalant dependence-Data from a tertiary care center in South India. Indian Journal of Psychological Medicine. 2012;34(3):232-6.
Available:<http://dx.doi.org/10.4103/0253-7176.106017>
19. Patil RC, Tavaragi MS, Sushma C. Inhalant abuse in adolescents in North Karnataka: A case series. Journal of Psychiatry Spectrum. 2022;1(2): 133-5.
Available:http://dx.doi.org/10.4103/jopsys.jopsys_1_22
20. Gordon K, Kutywayo A, Frade S, Naidoo N, Mullick, S. Socio-demographic and social support factors related to substance use in South African in-school adolescents- Insights from the girls achieve power (GAP Year) trial in three peri-urban settings. Gates Open Research. 2021;5(154).
Available:<https://doi.org/10.12688/gatesopenres.13422.1>
21. Mahmood N, Othman S, Al-Tawil N, Al-Hadithi T. Substance use among high school students in Erbil City, Iraq-Prevalence and potential contributing factors. Eastern Mediterranean Health Journal. 2019; 25(11):806-12.
Available:<https://doi.org/10.26719/emhj.19.022>
22. Mohammadpoorasl A, Nedjat S, Fakhari A, Yazdani K, Foroushani AR, Fotouhi A. Substance abuse in high school students in association with socio-demographic

- variables in North-West of Iran. Iranian Journal of Public Health. 2012;41(12):40–6.
23. Ajonye AA, Idoko E. Socio-demographic factors and depression on drug abuse amongst undergraduate students of Benue State University, Makurdi. International Journal of Social Sciences and Humanities Review. 2022;12(1):513–21.
 24. Gudaji MI, Habib ZG. Socio-demographic factors associated with psychoactive substance use among commercial motorcycle operators in Kano, Nigeria. Open Journal of Psychiatry. 2016;6:76-5.
Available:<http://dx.doi.org/10.4236/ojpsych.2016.61009>
 25. Narain R, Sardana S, Gupta S. Prevalence and risk factors associated with substance use in children- A questionnaire-based survey in two cities of Uttar Pradesh, India. Indian Journal of Psychiatry 2020;62(5):517–23.
Available:https://doi.org/10.4103/psychiatry.IndianJPsychiatry_595_19
 26. Rukundo A, Kibanja G, Steffens K. Factors influencing psychoactive substance use among adolescents in public secondary schools in Uganda. The International Journal of Alcohol and Drug Research. 2017;6(1):69-76.
Available:<http://dx.doi.org/10.7895/ijadr.v6i1.237>
 27. Wubetu AD, Getachew S, Negash W. Substances use and its association with socio-demographic, family, and environment-related factors among technical and vocational education and training college students in Ataye, Ethiopia- An institution-based cross-sectional study. BMC Public Health. 2020; 20(1):1691.
Available: <https://doi.org/10.1186/s12889-020-09797-w>
 28. Whitesell M, Bachand A, Peel J, Brown M. Familial, social, and individual factors contributing to risk for adolescent substance use. Journal of Addiction. 2013;2013:19
Article ID-579310.
Available:<http://dx.doi.org/10.1155/2013/579310>

© 2023 Katoch et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97459>