



***The World's Largest Open Access Agricultural & Applied Economics Digital Library***

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search  
<http://ageconsearch.umn.edu>  
[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*



# **Knowledge of the Respondents towards Paddy Production Technology in Mahabubnagar District of Telangana, India**

**Bheempogu Monica <sup>a++\*</sup> and Syed H. Mazhar <sup>a#</sup>**

<sup>a</sup> Department of Agricultural Extension and Communication, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, 211007, U.P., India.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/AJAEES/2023/v41i92017

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101862>

**Received: 17/04/2023**

**Accepted: 19/06/2023**

**Published: 20/06/2023**

**Original Research Article**

## **ABSTRACT**

The present study aims to highlights the Knowledge of respondents towards paddy production technology in Mahabubnagar district of Telangana, India. A structured interview was conducted among that 120 farmers were selected from 20 villages, with six respondents from each village. Data were collected by using the pre-tested schedule and analysed using appropriate statistical tools. It was found that most respondents were having a medium level of socio-economic profile. On analysing the knowledge level, the education level, the highest number of respondents i.e., 53.33 percent were having a medium level of knowledge followed by percent had high and 21.67 percent had low levels. On analysing the education level, the highest number of respondents i.e.,

<sup>++</sup> PG Final Year Student;

<sup>#</sup> Associate Professor;

\*Corresponding author: E-mail: monicabheempogu98@gmail.com;

53.33 percent were having a medium level of education followed by 33.33 percent having high and 12.5 percent having low levels. Independent variables namely age education annual income land holding, mass media exposure extension contacts, and economic motivation had positive and significant relationships with adoption respectively. The major constraints faced were delay in doing due to monsoon prolong, complexity, lack of skill and knowledge, unavailability of laborers, and high cost of input and storage facilities respectively.

**Keywords:** Knowledge; paddy growers; production; technology.

## 1. INTRODUCTION

"Paddy, also called rice paddy, small, level, flooded field used to cultivate rice in southern and eastern Asia. Wet-rice cultivation is the most prevalent method of farming in the far east, where it utilizes a small fraction of the total land yet feeds most of the rural population. Rice was domesticated as early as 3500 BC, and by about 2,000 years ago it was grown in almost all the present-day cultivation areas, predominantly deltas, floodplains, and coastal plains, and some terraced valley slopes.

Rice, (*Oryza sativa*), edible starchy cereal grain, and the grass plant (family Poaceae) by which it is produced. Roughly one-half of the world's population, including virtually all East and Southeast Asia, is wholly dependent upon rice as a staple food: 95 percent of the world's rice crop is eaten by humans. Rice is cooked by boiling or it can be ground into flour. It is eaten alone and in a great variety of soups side dishes, and main dishes in Asian, middle eastern, and many other cuisines. Other products in which rice is used are breakfast cereals, noodles, and such alcoholic beverages as Japanese sake" [1,2].

"The cultivated rice plant is an annual grass and grows to about 1.2 meters (4 feet) in height. The leaves are long and flattened and are borne on hollow stems. The fibrous root system is often broad and spreading. The panicle, or inflorescence (flower cluster), is made up of spikelets bearing flowers that produce the fruit, or grain. Varieties differ greatly in the length, shape, and weight of the panicles and the overall productivity of a given plant" Bassim [3], Dipak and Das [4].

"In the 1960s the so-called Green Revolution, an international scientific effort to diminish the threat of world hunger, produced improved strains of numerous food crops, including that known as miracle rice. Breed for disease resistance and increased productivity, this variety is characterized by a short sturdy stalk that

minimizes loss from drooping. Poor soil conditions and other factors, however, inhibited its anticipated widespread success" Kirar and Mehta [5]; Arvind and Pyasi [6].

Many cultures have evidence of early rice cultivation, including China, India, and the civilizations of Southeast Asia. However, the earliest archaeological evidence comes from central and eastern China and dates to 7000-5000 BCE. More than 90 percent of the world's rice is grown in Asia, principally in China, India, Indonesia, and Bangladesh, with smaller amounts grown in Japan, Pakistan, and various Southeast Asian nations. Rice is also cultivated in parts of Europe, North and South America, and in Australia.

## 2. METHODOLOGY

The study was conducted in the Mahabubnagar district of Telangana state, India. The study attempts to describe and analyse the impact of technology utilization behaviour of paddy growers.

There are 15 Blocks in Mahabubnagar district, 30 villages, out of which Jadcherla were selected purposely for study. In that 6 villages were selected purposely i.e (Kaverampet, Gangapur, Ranipet, Lingampet, Midgil, Jagboinpalli), Thus in all 120 paddy growers constituted sample for the investigation. Based on objectives of study an interview schedule was prepared. The information was elucidated from respondents with the help of pre structured schedule through descriptive research design. The information was collected by personally interviewing respondents using pre structured interview schedule.

## 3. RESULTS AND DISCUSSION

The study was undertaken to assess the knowledge of the respondents towards paddy production technology.

**Table 1. Knowledge of the respondent towards improved paddy production practices**

Sl. No.	Statements	Evaluation		
		Fully Correct (%)	Partially Correct (%)	Not Correct (%)
1.	Field Preparation: i. Traditional method-2-3 times plow ii. Use of zero tillage machine iii. Surface seeding method iv. Puddling v. Transplantation	58 (48.33%)	44 (36.67%)	18 (15.00%)
2.	Improved variety: i. Telangana Sona ii. Kunaram - 1638 iii. Kunaram – 118 iv. Jagital – 94423 v. Tella Hamsa	42 (35.00%)	44 (36.67%)	34 (28.33%)
3.	Seed and its treatment: i. Vitavax ii. Carbendazim iii. Pseudomonas fluorescens	26 (21.67%)	64 (54.16%)	30 (25.00%)
4.	Sowing time: i. July to October ii. Feb to May	40 (33.33%)	48 (40.00%)	32 (26.67%)
5.	Spacing: i. 22.5 to 23cm ii. 15 to 18cm iii. 20-22.5cm	40 (33.33%)	54 (45.00%)	26 (21.67%)
6.	Fertilizers: i. 120:60:40KgNPK/ha ii. 90:60:40KgNPK/ha	28 (23.33%)	70 (58.33%)	22 (18.33%)
7.	Irrigation: i. 2 times ii. 3 times iii. 6 times	36 (30.00%)	60 (50.00%)	24 (20.00%)
8.	Weeding and hoeing operations: i. 2 times ii. 3 times iii. 4 times	30 (25.00%)	43 (35.83%)	47 (39.16%)
9.	Weed control: i. 2,4-D ethyl ester ii. Butachlor iii. Benthiocarp iv. Anilophos	28 (23.33%)	62 (51.67%)	30 (25.00%)
10.	Diseases: i. Brown spot ii. Sheath blight iii. Bacterial blight iv. Blast of rice v. Tundu vi. Bacterial streak	32 (26.67%)	48 (40.00%)	40 (33.33%)
11.	Harvesting i. 80-90 days ii. 90-100 days iii. 100-110 days iv. 110-120 days	30 (25.00%)	38 (31.67%)	52 (43.33%)
12.	Yield: i. 20-25 quintal/ha ii. 25-30 quintal/ha iii. 30-35 quintal/ha	26 (21.67%)	70 (58.33%)	24 (24.00%)

It can be reported that regarding field preparation 48.33 per cent, 36.67 per cent and 15.00 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding Improved variety 35.00 per cent, 36.67 per cent and 28.33 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding Seed treatment 21.67 per cent, 54.16 per cent and 25.00 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding Sowing time 33.33 per cent, 40.00 per cent and 26.67 per cent of respondents reportedly fully correct, partially correct and not correct response respectively.

Meanwhile regarding spacing 33.33 per cent, 45.00 per cent and 21.67 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding fertilizers 23.33 per cent, 58.33 per cent and 18.33 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding irrigation 30.00 per cent, 50.00 per cent and 20.00 per cent of respondents reportedly fully correct, partially correct and not correct response


respectively. Regarding weeding and hoeing 25.00 per cent, 35.83 per cent and 39.16 per cent of respondents reportedly fully correct, partially correct and not correct response respectively [7-11].

Similarly regarding weed control 23.33 per cent, 51.67 per cent and 25.00 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding diseases 26.67 per cent, 40.00 per cent and 33.33 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding harvesting 25.00 per cent, 31.67 per cent and 43.33 per cent of respondents reportedly fully correct, partially correct and not correct response respectively. Regarding yield 21.67 per cent, 58.33 per cent and 24.00 per cent of respondents reportedly fully correct, partially correct and not correct response respectively.

It was clearly visible that majority 53.33 per cent of paddy growers had medium level of knowledge on paddy production technology, 21.67 per cent and 25.00 per cent of the paddy growers had low and high level of knowledge on paddy production technology respectively.

**Table 2. Distribution of respondents according to their overall knowledge level**

Knowledge level	Frequency	Percentage
Low (13-20)	26	21.67
Medium (21-27)	64	53.33
High (28-34)	30	25.00
<b>Total</b>	<b>120</b>	<b>100.00</b>



**Fig. 1. Distribution of respondents based on their overall knowledge level**

#### 4. CONCLUSION

It was concluded that majority 53.33 per cent of the paddy growers had medium level of knowledge on paddy production technology, 21.67 per cent and 25.00 per cent of the paddy growers had low and high level of knowledge on paddy production technology respectively. Knowledge level of majority of respondents was medium. Minimum support price of paddy should be declared well in advance by Government to increase the knowledge of growers towards paddy production technology.

#### CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Dhenge, Mankar. Knowledge level of farmers about integrated pest management practices of paddy. The Ecoscan. An International Quarterly Journal of Environmental Sciences. 2014;6(5):27-33.
2. Ajay Kumar and Sandeep. Knowledge level of farmers regarding system of rice intensification (SRI) method in Puri district of Odhisa. Indian Journal of Pure Applied and Biosciences. 2020;8(3):270-277.
3. Bassim. Knowledge level of rice farmers regarding cultivation practices in Mahanawiyah district, AL-Qadisiya Province, Iraq. International Journal of Agricultural Science, Research and Technology in Extension and Education Systems (IJASRT in EES). 2018;8(2): 71-78.
4. Dipak, Das. Knowledge on SRI (System of Rice Intensification) of Farmers in Tripura, India. International Journal Curriculum Microbiological Applied Sciences. 2018; 7(3):3586-3592.
5. Kirar, Mehta. Extent of knowledge of tribal farmers about rice production technology. Indian Research Journal of Extension Education. 2016;9(1):32-35.
6. Arvind and Pyasi. A study on knowledge about organic farming practices possessed by the farmers and their adoption in selected blocks of Balaghat district (MP). Plant Archieves. 2020;20(2):4621-4626.
7. Iqbal, Amardev. A study on the Knowledge level and constraints faced by the paddy growers of Jammu District of J&K State. Agro Economist. 2016;3(1):31-35.
8. Ram and Upadhyay. Knowledge of improved rice production technology by the farmers in Dungarpur District of Rajasthan. Rajasthan Journal Extension Education. 2012;20:97-101.
9. Randir, Ramesh. Knowledge and adoption level of farmers of Haryana about scientific rice cultivation practices. Journal of Community Mobilization and Sustainable Development. 2014;8(1):24-28.
10. Sharma, Rampal. Farmers knowledge towards aerobic rice cultivation in Muktsar district of Punjab. Indian Research Journal of Extension Education. 2017;17(4):83-86.
11. Singh, Yadav. Knowledge and adoption gap of tribal farmers of Bastar towards rice production technology. American International Journal of Research in Humanities, Arts and Social Sciences. 2014;5(1):54-56.

© 2023 Monica and Mazhar; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sciarticle5.com/review-history/101862>