



AgEcon SEARCH

RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

VOL. 6

MAY, 1931

NO. 1

HILGARDIA

A Journal of Agricultural Science

PUBLISHED BY THE

California Agricultural Experiment Station

CONTENTS

The Digestibility of Bur Clover as Affected by
Exposure to Sunlight and Rain

H. R. GUILBERT and S. W. MEAD

The Effect of Leaching on the Nutritive
Value of Forage Plants

H. R. GUILBERT, S. W. MEAD, and H. C. JACKSON

UNIVERSITY OF CALIFORNIA PRINTING OFFICE
BERKELEY, CALIFORNIA

EDITORIAL BOARD

C. B. HUTCHISON, M.S.

F. T. Bioletti, M.S.
Viticulture

R. E. Clausen, Ph.D.
Genetics

H. E. Erdman, Ph.D.
Agricultural Economics

H. M. Evans, M.D.
Nutrition

H. S. Fawcett, Ph.D.
Plant Pathology

G. H. Hart, M.D., D.V.M.
Animal Husbandry

D. R. Hoagland, M.S.
Plant Nutrition

W. L. Howard, Ph.D.
Pomology

W. P. Kelley, Ph.D.
Chemistry

H. A. Jones, Ph.D.
Truck Crops

C. S. Mudge, Ph.D.
Bacteriology

H. J. Quayle, M.S.
Entomology

H. S. Reed, Ph.D.
Plant Physiology

W. W. Robbins, Ph.D.
Botany

H. R. Tolley, B.S.
Agricultural Economics

J. Traum, D.V.M.
Veterinary Science

F. J. Veihmeyer, Ph.D.
Irrigation

H. B. Walker, C.E.
Agricultural Engineering

THE EFFECT OF LEACHING ON THE NUTRITIVE VALUE OF FORAGE PLANTS¹

H. R. GUILBERT² S. W. MEAD,³ AND H. C. JACKSON⁴

INTRODUCTION

A significant decrease in digestibility of bur clover after exposure to sunlight and rain has been reported in the first paper and evidence was presented which indicated that the greater part of the decrease could be accounted for by the loss of soluble constituents through the action of rain. Field observations support these findings.

After late rains on cured range feed, cattle have been observed to cease gaining and to require supplemental feeding in order to fatten sufficiently to be marketable. Under such conditions it is difficult, in many cases, even to maintain breeding stock. This situation prevailed over a large area of California in 1929. Extensive supplemental feeding was required in many areas to fatten the cattle for beef, and stock cattle, generally, suffered from the poor feed. Among the abnormal conditions reported in cattle from some areas were pica, particularly bone craving, deformed calves, difficult parturition, and retained placenta. Many ewes which apparently were unable to lactate abandoned their lambs. The indications are that these troubles were directly associated with the poor quality of the feed.

Rain followed by warm, humid weather is favorable to the development of molds. Frequently, however, the feed dries quickly with little or no molding; and yet deterioration has occurred, as evidenced by the condition of livestock. Field observations therefore indicate that the leaching effect of rain may be the most important factor in the loss of nutritive value.

¹ This work became cooperative with the Bureau of Animal Industry, United States Department of Agriculture, July 1, 1929.

² Assistant Animal Husbandman in the Experiment Station.

³ Associate Animal Husbandman in the Experiment Station.

⁴ Cooperative Agent, Bureau of Animal Industry, United States Department of Agriculture.

REVIEW OF LITERATURE

Wolff⁽¹⁾ in Germany reported that 20 per cent of the dry substance of hay may be lost by simply soaking in cold water, and that clover hay suffers from rain more than meadow hay because from 25 to 40 per cent of its dry substance may be dissolved. He reported analyses by Ritthausen at Möckern on two samples of clover hay which were cut at the beginning of the flowering period from the same field at the same time. One was quickly dried and the other left to lie for two weeks exposed to intermittent rains. The percentage composition was reported as follows:

	Water	Crude protein	Nitrogen-free extract and fat	Crude fiber	Ash
Not rained upon.....	16.0	14.6	36.1	25.3	8.0
Rained upon	16.0	15.8	23.4	37.4	7.5

The principal change was a decrease in the most soluble carbohydrates and ash and a resultant increase in crude fiber.

Headden⁽²⁾ of the Colorado Experiment Station reports the percentage composition of alfalfa hay before and after being exposed to rain as follows:

	Protein	Nitrogen-free extract	Crude fat	Crude fiber	Ash
Not rained upon.....	18.71	38.71	3.94	26.46	12.18
Rained upon	11.01	33.64	3.81	38.83	12.71

The latter hay was damaged by three rains at intervals of two days or more. The author states: "The mechanical loss of leaves and stems would tend to change the composition of the hay in the direction indicated by the analyses, but for good reasons we do not consider this to enter largely into this particular case, but attribute the changes in composition to the action of heat and moisture."

Henry and Morrison⁽³⁾ state: "Exposure to the sun reduces the palatability by bleaching and causes a loss of aromatic compounds, dew works injury and rain carries away the more soluble portions. The actual damage from rain is even greater [than analysis shows], for the nutrients lost were those most soluble and hence most easily digested." According to Piper⁽⁴⁾ the destruction of the green chlorophyll by sunlight is increased by the action of dew. He also makes the following statement:

“Westgate sprinkled perfectly cured crimson clover hay with water to imitate rain for one hour on each of three successive days. On analysis it was found in comparison with a sample unsprinkled to have lost about three-fourths of its sugar, one-ninth of its protein and three-fourths of its ash constituents.”

Le Clerc and Breazeale⁽⁵⁾ harvested a sample of greenhouse barley at the heading period and subjected the whole plant to leaching. They state: “The plant was placed in a large evaporating dish and soaked with water for several minutes. After drying, this operation was again repeated. The plant was then dried and analyzed. The washings were also analyzed, the results showing that 1.6 per cent of the whole nitrogen of the plant was lost on washing or soaking, 36 per cent of the phosphoric acid, 65 per cent of the potash, 52 per cent of the soda, 45 per cent of the magnesia, and 75 per cent of the chlorin.”

Various other plants, among them rice, wheat, apple twigs, oats, and potatoes, were experimented upon in a number of ways and at different stages of growth. The largest loss of nutrients by leaching was found to occur when the plant was at maturity. In the growing state, however, some losses occur. When wheat plants were in bloom, the amounts washed out of the straw and leaves were as follows: “Nitrogen, 1.4 per cent; phosphoric acid, nothing; potash, 4.4 per cent; soda, 12.7 per cent; lime, nothing; magnesia, 10.3 per cent; chlorin, 7.6 per cent.” From this Le Clerc and Breazeale state: “It is not contended that the green plants give off very much of their plant food by such treatment, for it is very probable that most of the ash ingredients removed by washing are those which were in the dead or wilted tissue, as it is well known that when plants dry or wilt, the inorganic constituents exude to the surface, where they may be easily washed off if subjected to the action of rain, dew, etc. As illustrative of this, an experiment made with freshly cut grass showed that when the grass was dried previous to treatment with water a much larger amount of ash materials was washed out. This explains why it is that when freshly cut hay has been rained upon it is only slightly injured, whereas if the rain comes after the hay has been dried the loss is considerable, sometimes as much as half of the ash ingredients being thus removed.”

Digestion experiments⁽⁶⁾ and field observations indicate that the leaching effect of rain is an important factor in loss of nutritive value. Although the effect of leaching is indicated in the literature cited, these experiments do not appear directly applicable to range conditions. Experiments were therefore undertaken with species of forage

plants common to California ranges for the purpose of showing the extent and character of the losses which could be caused by rain, and thus contribute further information on the reasons for the observed deterioration in nutritive value.

LEACHING EXPERIMENTS WITH BUR CLOVER

A quantity of bur clover was cut at an advanced stage of maturity but was still green in color. It was dried in the sun for one day in a thin layer and then cured for 6 days in cocks. This was the lot designated as No. 1, in the digestion experiments reported in the first paper.

For the leaching experiment a sample weighing 172 grams was placed in a soil percolator for one hour with two liters of distilled water; it was then washed twice with one-liter portions of water, and the final volume of the extract made up to four liters. The extract was first filtered with suction through linen, and the portion used for analysis was filtered through filter paper to remove any solids in suspension. The results are summarized in table 1.

TABLE 1

PERCENTAGE OF CHEMICAL CONSTITUENTS EXTRACTED FROM BUR CLOVER No. 1; DRY BASIS	
Crude protein	16.20
Nitrogen-free extract	28.32
Calcium	30.43
Phosphorus	45.71
Chlorine	86.02
Total silica-free ash	59.11
Total solids	19.94

As shown in table 1, nearly 20 per cent of the dry matter of the forage was extracted by water. The silica-free ash, the various ingredients of the ash, and the nitrogen-free extract were most susceptible to leaching. Although the percentage loss of protein was least, the amount extracted appears significant particularly since the digestion experiments show a decreased availability of the protein after exposure to weathering and to rain.

Although the nutritive value of bur clover is markedly affected by rain, it is still much higher than that of the dried grasses and other forage under similar conditions; and stock can be maintained fairly

well so long as the clover burs are obtainable in adequate quantities. The relative effect of leaching on the burs and the enclosed seeds as compared with the remainder of the plant is, therefore, of interest, and another experiment was carried out to determine this.

A sample from another lot of bur clover was used in this experiment. The burs were separated from the stems and leaves, and samples of each were thoroughly mixed and quartered, opposite quarters being taken for leaching and analysis. The samples for leaching were weighed and placed in soil percolators, which were then filled with distilled water and allowed to stand for approximately one hour. The extract was then drawn off, and the residue washed with distilled water until it came off practically colorless. The extract was then made up to the nearest convenient volume. A portion of each of the original samples and of the extracts were analyzed. The results are shown in table 2.

TABLE 2

PERCENTAGE OF CHEMICAL CONSTITUENTS EXTRACTED FROM CLOVER BURS AS COMPARED TO STEMS AND LEAVES; DRY BASIS

	Clover burs	Leaves and stems
Crude protein	9.3	11.2
Nitrogen-free extract	15.3	35.3
Calcium	9.5	19.6
Phosphorus	16.3	58.7
Total silica-free ash.....	26.5	34.4
Total solids	10.8	21.7

The clover burs were much more resistant to leaching than the stems and leaves, the loss of total solids being only half as great. The difference in the percentage of calcium, phosphorus, and carbohydrate extracted is especially large.

LEACHING EXPERIMENT WITH OAT HAY

A sample of good quality red-oat hay was cut into 2 to 3-inch lengths, and a weighed quantity placed in a soil percolator with distilled water and allowed to stand for approximately one hour. The extract was then drawn off and the residue washed with distilled water until it came off practically colorless. The extract was filtered and made up to a convenient volume. Samples of the original, of the residue after leaching, and of the extract were analyzed.

The percentage of chemical constituents extracted is shown in table 3.

TABLE 3
PERCENTAGE OF CHEMICAL CONSTITUENTS EXTRACTED
FROM RED-OAT HAY; DRY BASIS

Crude protein	1.1
Nitrogen-free extract	14.2
Calcium	31.3
Phosphorus	21.4
Chlorine	67.2
Total silica-free ash.....	59.2
Total solids	10.4

The loss of total solids was less than from bur clover under similar conditions of extraction. The oat hay was lower in protein, and a small amount was removed by leaching. The percentage of carbohydrate soluble in water was also less than in the clover. The loss of ash, however, was nearly 60 per cent of the total.

LEACHING EXPERIMENTS WITH NATURALLY-CURED RANGE FORAGE

All of the previous experiments were conducted on samples cut in the green stage and dried. Because changes in composition occur during the latter stages of maturity; it appeared desirable to experiment with samples of naturally-cured range feed. The following is a description of the samples used:

Sample No. 200 was soft chess (*Bromus hordeaceus*). It was dry and bleached, and the seeds were mostly shattered. The sample was collected on June 3, 1930.

Sample No. 212 was a composite in which stork's-bill alfilaria (*Erodium botrys*) predominated. It was collected on June 5, 1930; it was dry and bleached, and the seeds were completely shattered.

Sample No. 215 was dry, bleached bur clover (*Medicago hispida*). Many of the burs had fallen to the ground, but a large percentage of these were included in the sample in order to have it as nearly representative of the material grazed as possible. It was collected on June 9, 1930.

All the samples were ground to pass through a 40-mesh screen. Fifty-gram samples of each were placed in flasks with 500 cc. of distilled water and allowed to stand for approximately one hour. The

extracts were filtered with suction and the residues washed with 150 to 200 cc. of distilled water. The final volume of extract varied from 500 to 590 cc. The purpose of the experiment was to ascertain whether or not the relative loss of the various nutrients would be similar to that found in previous experiments.

The percentage of each nutrient extracted is shown in table 4.

TABLE 4
PERCENTAGE OF CHEMICAL CONSTITUENTS EXTRACTED FROM NATURALLY-CURED FORAGE; DRY BASIS

	No. 200, soft chess	No. 212, alfilaria	No. 215, bur clover
Crude protein	18.2	12.0	12.9
Nitrogen-free extract	12.1	12.6	15.0
Calcium	30.5	11.8	9.5
Phosphorus	37.0	45.4	31.9
Chlorine	72.6	63.0	76.9
Total silica-free ash	62.7	28.3	32.8
Total solids	10.7	8.0	10.6

The greatest percentage loss was in the ingredients of the silica-free ash, a fact which is in agreement with the other experiments. The percentage loss of ash from soft chess was approximately double that from alfilaria and bur clover. Compared with all the previous experiments, a higher percentage of protein relative to other ingredients was extracted.

Another lot of four naturally-cured samples, each weighing 25 grams, was prepared in the same way as the previous samples and extracted with distilled water for 5 hours in a Soxhlet apparatus. At the end of this time the water which came over was colorless. The treatment of these samples probably approached complete extraction. A brief description of the samples follows:

- No. 221. Soft chess, dry, bleached, and the seeds partly shattered.
- No. 222. Wild oats, dry, bleached, and the seeds mostly shattered.
- No. 226. Bur clover, dry, bleached, and consisting largely of burs and some stems. Most of the leaf material had been shattered and lost.
- No. 228. Stork's bill alfilaria, dry, bleached, and the seeds shattered.

All of the samples were collected in the same locality on June 10, 1930. They were taken from a different area than the samples reported in table 4, and had been dry somewhat longer. The results of the Soxhlet extraction are shown in table 5.

TABLE 5

PERCENTAGE OF CHEMICAL CONSTITUENTS EXTRACTED FROM NATURALLY-CURED FORAGE WITH SOXHLET APPARATUS; DRY BASIS

	No. 221, soft chess	No. 222, wild oats	No. 226, bur clover	No. 228, alfilaria
Crude protein	6.8	11.8	14.3	12.2
Nitrogen-free extract	6.1	8.9	15.6	17.1
Calcium	26.3	47.2	10.2	11.7
Phosphorus	18.8	24.6	27.7	42.5
Total silica-free ash.....	30.0	66.9	34.8	24.6
Total solids	5.0	8.3	11.9	12.1

The loss of total solids varied from 5 to 12 per cent, and the greatest percentage loss was in the silica-free ash. There was a relatively greater loss of calcium than of phosphorus in wild oats and soft chess, whereas the reverse was found in bur clover and alfilaria. The variation in chemical composition between this lot of samples and the first lot of naturally cured forage, may account for some of the variations in leaching. The soft chess sample, No. 221, for example, was significantly lower in protein and nitrogen-free extract and higher in fiber than sample No. 200. The general trend of the results is the same as in previous experiments.

The results of this experiment indicate that the lower amount of total solids extracted in the range samples as compared to bur clover No. 1 results not so much from the method of leaching as from a lower content of soluble material. The loss of total solids in the wild oats and in soft chess, sample No. 200, was not far from that found in red-oat hay, table 3. The loss from bur clover sample No. 226 was similar to that found for a pure sample of burs, table 2. Because of the shattering of leaves under field conditions, the bur clover and alfilaria samples used in these experiments are not representative of the entire plant. It is not possible, therefore, from the data available, to compare directly the solubility of nutrients in forage cut in advanced stages of maturity but still green, with the fully matured and naturally dried forage.

THE COMPOSITION OF RESIDUES AFTER LEACHING

The foregoing data have shown the percentage loss of the various nutrients through leaching and hence indicate the possible loss in tonnage of cured feed. The utilization of the material which is left, however, is a most important consideration. The composition of the residues are shown in tables 6, 7, and 8.

TABLE 6

COMPARISON OF THE PERCENTAGE COMPOSITION OF UNLEACHED PORTIONS OF BUR CLOVER AND RED-OAT HAY WITH THE RESIDUES AFTER LEACHING; DRY BASIS

	Bur clover No. 1*		Bur clover burs		Bur clover stems and leaves		Red-oat hay	
	Unleached sample	Leached residue	Unleached sample	Leached residue†	Unleached sample	Leached residue†	Unleached sample	Leached residue
Crude protein.....	16.86	17.77	15.12	15.38	16.33	18.51	5.15	5.62
Nitrogen-free extract	45.54	41.67	52.21	49.73	46.93	38.76	60.35	57.68
Ether extract	3.31	3.94	2.54	2.85	2.54	3.24	3.43	3.83
Crude fiber	25.31	31.25	23.78	26.66	23.81	30.41	25.00	28.22
Silica-free ash....	7.56	3.89	5.84	4.81	9.53	7.98	3.07	1.08
Acid-insoluble ash	1.42	1.48	0.51	0.57	0.86	1.10	3.00	3.57
Calcium	0.95	0.94	0.83	0.84	1.20	1.23	0.23	0.18
Phosphorus	0.30	0.22	0.40	0.38	0.20	0.10	0.18	0.16
Chlorine	0.60	0.20	0.76	0.29

* Results in the first two columns were obtained from a different sample from that reported in table 1.

† Computed from the original and the extract.

Nitrogen-free extract represents a large percentage of the total dry matter in the plant. In the leaching experiments shown in table 6 there was a considerable loss of this constituent, which has the effect of increasing the percentage of other ingredients of the residue in which the percentage loss is less.

The protein in the residue after leaching was in every case higher than in the original material. The nitrogen-free extract was lower in the residue in every case, while the ether extract and crude fiber were higher than in the unleached sample. The silica-free ash was reduced in every instance, and in the oat hay it was reduced to a little more than 1 per cent, which is extremely low.

Of the ingredients of the ash, calcium was least affected by leaching except in red-oat hay, where the calcium loss was greater than that of phosphorus. The phosphorus of bur clover No. 1 was reduced from 0.30 per cent to 0.22 per cent. In the case of the bur clover leaves and stems, the amount was reduced to 0.10 per cent, which is definitely low. Animals grazing on leached bur clover in which the per cent of burs eaten is less than in these samples may be ingesting less than optimum amounts of phosphorus.

The ratio of calcium to phosphorus in bur clover No. 1 was changed from approximately 3:1 to 4.5:1, while in the stems and leaves it was changed from 6:1 to 12:1. The clover burs contained twice as much phosphorus as the stems and leaves, and the ratio of calcium to phosphorus was not affected by leaching. The quantity of burs available may thus have a distinct bearing on the nutrition of animals grazing on dried range feed which has been subjected to rain.

The composition of the unleached material and of the residue after leaching of the samples of naturally-dried range feed is given in tables 7 and 8. The residues were not analyzed in this case but have been computed from the analyses of the original and of the extract.

TABLE 7

COMPARISON OF THE PERCENTAGE COMPOSITION OF THE UNLEACHED PORTIONS WITH LEACHED RESIDUES OF NATURALLY-CURED FORAGE; SILICA AND MOISTURE-FREE BASIS

	No. 200, soft chess		No. 212, alfilaria		No. 215, bur clover	
	Unleached sample	Leached residue	Unleached sample	Leached residue	Unleached sample	Leached residue
Crude protein	9.43	8.64	5.74	5.49	15.02	14.62
Nitrogen-free extract	60.44	59.55	53.33	51.41	43.19	41.04
Ether extract	1.69	1.89	2.65	2.88	1.91	2.14
Crude fiber	25.70	28.78	34.68	37.70	33.37	37.31
Silica-free ash	2.74	1.14	3.60	2.52	6.51	4.89
Calcium	0.36	0.28	1.27	1.23	1.26	1.27
Phosphorus	0.27	0.19	0.11	0.07	0.25	0.19
Chlorine	0.22	0.07	0.27	0.11	0.39	0.10

Table 7 shows that the protein and nitrogen-free extract of the residue is lower than that of the original samples, and the ether extract and crude fiber is greater. The most significant change was in the amount of silica-free ash. The calcium content of soft chess was reduced by leaching, but there was no appreciable change in the calcium content of alfilaria and bur clover. In all samples the phosphorus was lowered and the chlorine greatly reduced.

TABLE 8
COMPARISON OF THE PERCENTAGE COMPOSITION OF THE UNLEACHED PORTIONS WITH
LEACHED RESIDUES OF NATURALLY-CURED FORAGE (SOXHLET
EXTRACTION); SILICA AND MOISTURE-FREE BASIS

	No. 221, soft chess		No. 222, wild oats		No. 226, bur clover		No. 228, alfilaria	
	Unleached sample	Leached residue	Unleached sample	Leached residue	Unleached sample	Leached residue	Unleached sample	Leached residue
Crude protein	6.78	6.65	4.56	4.39	18.23	17.74	5.24	5.23
Nitrogen-free extract.....	55.29	54.67	59.92	59.50	46.80	44.79	54.91	51.78
Ether extract	2.39	2.52	1.85	2.02	3.63	4.12	2.82	3.21
Crude fiber	31.54	33.21	30.05	32.78	25.71	29.18	28.72	32.65
Silica-free ash....	4.00	2.95	3.62	1.31	5.63	4.17	8.31	7.13
Calcium	0.24	0.19	0.29	0.17	1.20	1.23	1.81	1.82
Phosphorus	0.25	0.21	0.13	0.11	0.30	0.25	0.08	0.05

In agreement with the previous experiments, table 8 shows that the greatest change was in the amount of silica-free ash. The phosphorus was lowered in the residues of all the samples; the calcium was reduced in wild oats and soft chess; but the percentage remained practically unchanged in the alfilaria and bur clover. The protein and nitrogen-free extract of the residues were slightly lower than that of the unleached samples, and the ether extract and crude fiber were greater, as was found with the other samples of naturally-cured forage.

DISCUSSION

It was recognized from the beginning that it would be impossible to conduct these experiments so that the leaching would be comparable to that resulting from a given amount of rainfall on the range. The methods which have been used are therefore purely arbitrary and intended only to show the relative losses of the various constituents in order that the reasons for the observed decrease in nutritive value might be more clearly understood.

The results of some of the experiments probably represent nearly complete extractions. Probably the amount extracted was not in excess of that sometimes occurring on the range, when the feed remains saturated for one or two days and is leached by intermittent showers totaling one to three inches or more of rainfall.

The loss of a high percentage of the silica-free ash is significant. Elliot, Orr, and Wood⁽⁷⁾ in part II of their investigation on the mineral content of pasture grass in the British Isles concluded that there was no striking difference in the total energy value of good and poor pastures, but that wide differences existed in the proportion of the mineral constituents and that high mineral content was associated with high nutritive value.

Aside from the specific functions of inorganic elements in metabolism, the concentration of mineral salts in the intestinal tract appears to have important functions in the processes of digestion. In regard to this, Orr⁽⁸⁾ states: "The ebb and flow of fluid between the lumen of the gut and the blood stream is controlled by the concentration of mineral salts in the intestinal contents and the membrane lining the intestines. An increased amount of mineral salts in the intestinal contents tends to cause a flow of fluid from the blood to the intestines which in extreme cases causes diarrhea." It is a common observation on the ranges during the dry season that the feces of cattle become dry and comparatively hard. There is evidence that the mineral content of the feed may be responsible.

The loss of chlorine together with that of sodium (Le Clere and Breazeale⁽⁵⁾) undoubtedly accounts for the increase in salt consumption observed in cattle grazing upon forage which had been damaged by rain.

The removal of the most soluble carbohydrates and proteins may leave in the residue the more complex compounds and nutrients which are protected from water and enzyme action by cellulose walls, thus resulting in lower digestibility. The increased fiber content may also have the effect of depressing the digestibility of the other constituents,^(9, 10) in addition to being itself more difficult of digestion than other forms of carbohydrate. The reduction of soluble salt and of the soluble carbohydrate may have a marked effect on palatability.

A comparison of the composition of the unleached samples and the residues after leaching does not indicate clearly the extent to which leaching has occurred. A slight increase in one nutrient in the residue might mean a considerable loss in certain other nutrients which would be shown only by an analysis of the extract. An increase of from 3 to 6 per cent in crude fiber for example, was coincident, in these experiments, with losses of from 10 to 20 per cent of total solids.

SUMMARY

The greatest percentage loss caused by leaching was of the constituents of the silica-free ash, which represents that portion of the mineral in the plant which is available to the animal. This loss varied from 25 to 67 per cent in the different samples.

Of the ingredients of the ash which were analyzed, chlorine was lost in greatest amount. The percentage loss amounted to 67 per cent in oat hay and as high as 86 per cent in bur clover. The experiments indicate that practically all of the chlorine may be leached out of dried pastures by excessive rainfall. This is in agreement with the observed salt requirements of stock after feed has been damaged by rain.

In the bur clover and alfalfa samples the percentage of calcium in the forage after leaching was not significantly different from the unleached portion. Phosphorus was distinctly lower in the leached material, particularly in the case of the bur clover leaves and stems. The ratio of calcium to phosphorus in these species thus tends to be widened by leaching. In the grass species a larger percentage of calcium was lost, and the percentage in the residue was lower than in the unleached sample. This may be significant from the standpoint of nutritive value because these dry grasses are in general probably below optimum in calcium. Since the phosphorus is also reduced, the Ca:P ratio remained practically unchanged.

The amount of nitrogen-free extract lost by leaching varied from 6 per cent of the total in a sample of dry bleached soft chaff to 35 per cent in bur clover stems and leaves. This loss represents largely the sugars, which are easily digested and which may also influence palatability.

The amount of crude protein lost varied from 1 per cent of the total in oat hay to 16 per cent in bur clover and 18 per cent in soft chaff. The loss of protein by leaching from the samples which were cut green and dried was relatively less than that of other constituents, so that there was a higher per cent of protein in the residue than in the original sample. In the naturally-cured samples there was a reduction in the per cent of protein in the leached residue as compared with the original sample. In general the change in percentage of protein does not appear very significant, but there may be a very

significant difference in availability of the residual material as compared with that extracted.

Ether extract is influenced only slightly by leaching. The percentage in the residue is higher than in the original material.

Crude fiber remained entirely in the residue. The decrease in other nutrients caused a very significant increase in the percentage of this material in the dry matter after leaching. An increase of from 3 to 6 per cent in crude fiber was coincident with losses of from 10 to 20 per cent of total solids. The increased fiber content may have a depressing effect upon digestibility of other nutrients in addition to being, itself, difficult of digestion.

LITERATURE CITED

- ¹ WOLFF, EMIL.
1895. *Farm Foods*: 357 p. English ed. trans by H. B. Cousins. Gurney and Jackson, London.
- ² HEADDEN, WM. P.
1896. *Alfalfa*. Colorado Agr. Exp. Sta. Bul. 35:1-92; plates 17.
- ³ HENRY and MORRISON.
1923. *Feeds and feeding*. 18th ed., unabridged: 770 p. The Henry-Morrison Co., Madison, Wisconsin.
- ⁴ PIPER, C. V.
1924. *Forage plants and their culture*: 671 p. Macmillan Co., New York.
- ⁵ LE CLERC, J. A., and J. F. BREAZEALE.
1909. Plant food removed from growing plants by rain or dew. U. S. D. A. Yearbook 1908:389-402.
- ⁶ GUILBERT, H. R., and S. W. MEAD.
1931. The digestibility of bur clover as affected by exposure to sunlight and rain. *Hilgardia* 6(1):1-12.
- ⁷ ELLIOT, W. E., J. B. ORR, and T. B. WOOD.
1926. Investigation on the mineral content of pasture grass and its effect on herbivora. *Jour Agr. Sci.* 16:59-104.
- ⁸ ORR, J. B.
1925. The importance of mineral matter in nutrition. Rowett Res. Institute, collected papers 1:189-215.
- ⁹ ARMSBY, H. P.
1922. *The nutrition of farm animals*. 741 p. The Macmillan Co., New York.
- ¹⁰ MUMFORD, H. W., H. S. GRIDLEY, L. D. HALL, and A. D. EMMET.
1914. A study of the digestibility of rations for steers. Illinois Agr. Exp. Sta. Bul. 172:246-255.

The titles of the Technical Papers of the California Agricultural Experiment Station, Nos. 1 to 20, which HILGARDIA replaces, and copies of which may be had on application to the Publication Secretary, Agricultural Experiment Station, Berkeley, are as follows:

4. Effect of Sodium Chlorid and Calcium Chlorid upon the Growth and Composition of Young Orange Trees, by H. S. Reed and A. R. C. Haas. April, 1923.
5. Citrus Blast and Black Pit, by H. S. Fawcett, W. T. Horne, and A. F. Camp. May, 1923.
6. A Study of Deciduous Fruit Tree Rootstocks with Special Reference to Their Identification, by Myer J. Heppner. June, 1923.
7. A Study of the Darkening of Apple Tissue, by E. L. Overholser and W. V. Cruess. June, 1923.
8. Effect of Salts on the Intake of Inorganic Elements and on the Buffer System of the Plant, by D. E. Hoagland and J. C. Martin. July, 1923.
9. Experiments on the Reclamation of Alkali Soils by Leaching with Water and Gypsum, by P. L. Hibbard. August, 1923.
10. The Seasonal Variation of the Soil Moisture in a Walnut Grove in Relation to Hygroscopic Coefficient, by L. D. Batchelor and H. S. Reed. September, 1923.
11. Studies on the Effects of Sodium, Potassium, and Calcium on Young Orange Trees, by H. S. Reed and A. R. C. Haas. October, 1923.
12. The Effect of the Plant on the Reaction of the Culture Solution, by D. E. Hoagland. November, 1923.
14. The Respiration of Potato Tubers in Relation to the Occurrence of Black-heart, by J. P. Bennett and E. T. Bartholomew. January, 1924.
16. The Moisture Equivalent as Influenced by the Amount of Soil Used in its Determination, by F. J. Veihmeyer, O. W. Israelsen and J. P. Conrad. September, 1924.
17. Nutrient and Toxic Effects of Certain Ions on Citrus and Walnut Trees with Especial Reference to the Concentration and Ph of the Medium, by H. S. Reed and A. R. C. Haas. October, 1924.
18. Factors Influencing the Rate of Germination of Seed of *Asparagus Officinalis*, by H. A. Borthwick. March, 1925.
19. The Relation of the Subcutaneous Administration of Living Bacterium abortum to the Immunity and Carrier Problem of Bovine Infectious Abortion, by George H. Hart and Jacob Traum. April, 1925.
20. A Study of the Conductive Tissues in Shoots of the Bartlett Pear and the Relationship of Food Movement to Dominance of the Apical Buds, by Frank E. Gardner. April, 1925.