



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

*Department of Agricultural &
Resource Economics, UCB*
CUDARE Working Papers
(University of California, Berkeley)

Year 2008

Paper 1074

The Impact of Water Quality on
Southern California Beach Recreation: A
Finite Mixture Model Approach

James Hilger *

W. M. Hanemann †

*Bureau of Economics, Federal Trade Commission

†University of California, Berkeley

This paper is posted at the eScholarship Repository, University of California.

http://repositories.cdlib.org/are_uchb/1074

Copyright ©2008 by the authors.

The Impact of Water Quality on Southern California Beach Recreation: A Finite Mixture Model Approach

Abstract

This paper uses a finite mixture logit (FML) model to investigate the heterogeneity of preferences of beach users for water quality at beaches in Southern California. The results are compared with conventional approaches based conditional logit (CL) and random parameters logit (RPL). The FML approach captures variation in preferences by modeling individual recreator choices using a mixture of several distinct preference groups, where group membership is a function of individual characteristic and seasonal variables. The FML parameter estimates are used to calculate welfare measures for improvements in beach quality through a reduction of water pollution. The FML segment specific welfare measures bound the traditional CL and RPL mean welfare estimates, and have the advantage of highlighting the distribution of the population sample's preferences. Analysis of beach recreation site choice data indicates the existence of four representative preference groups within the survey respondent sample. As a result, willingness to pay measures for improvements in water quality and other beach site attribute changes can be weighted across individuals to calculate the distribution of individual welfare measures. One group of recreators is characterized as people who go to the beach and engage in water recreation with children. An interesting finding is that this group has a lower mean WTP for improving water quality than groups who go without children. This may well be an example of cognitive dissonance: parents find they go to the beach more often than others who don't have children, since that keeps the children occupied and happy, and they adapt their perception of the water quality to be consistent with their behavior.

Previous environmental and resource economic applications of the FML have been limited to applications with small choice sets (6) and group membership variables (4). This paper extends the FML model through the estimation of a

large (51) choice set with 9 membership variables. This application is the first to incorporate seasonal variables into the group membership function to capture seasonal heterogeneity.

Estimated welfare changes are calculated using the compensating variation measure for several hypothetical beach closure and water quality degradation scenarios. Estimation results indicate that the FML welfare estimates differ from those calculated using the traditional logit or RPL models. The FML model sheds light onto which subsets of beach recreators are likely to be impacted by different scenarios of resource change.

The Impact of Water Quality on Southern California Beach Recreation: A Finite Mixture Model Approach*

James Hilger and W. Michael Hanemann[†]

March, 2008

Abstract

This paper uses a finite mixture logit (FML) model to investigate the heterogeneity of preferences of beach users for water quality at beaches in Southern California. The results are compared with conventional approaches based conditional logit (CL) and random parameters logit (RPL). The FML approach captures variation in preferences by modeling individual recreator choices using a mixture of several distinct preference groups, where group membership is a function of individual characteristic and seasonal variables. The FML parameter estimates are used to calculate welfare measures for improvements in beach quality through a reduction of water pollution. The FML segment specific welfare measures bound the traditional CL and RPL mean welfare estimates, and have the advantage of highlighting the distribution of the population sample's preferences. Analysis of beach recreation site choice data indicates the existence of four representative preference groups within the survey respondent sample. As a result, willingness to pay measures for improvements in water quality and other beach site attribute changes can be weighted across individuals to calculate the distribution of individual welfare measures.

One group of recreators is characterized as people who go to the beach and engage in water recreation with children. An interesting finding is that this group has a lower mean WTP for improving water quality than groups who go without children. This may well be an example of cognitive dissonance: parents find they go to the beach more often than others who don't have children, since that keeps the children occupied and happy, and they adapt their perception of the water quality to be consistent with their behavior.

Previous environmental and resource economic applications of the FML have been limited to applications with small choice sets (6) and group membership variables (4). This paper extends the FML model through the estimation of a large (51) choice set with 9 membership variables. This application is the first to incorporate seasonal variables into the group membership function to capture seasonal heterogeneity.

Estimated welfare changes are calculated using the compensating variation measure for several hypothetical beach closure and water quality degradation scenarios. Estimation results indicate that the FML welfare estimates differ from those calculated using the traditional logit or RPL models. The FML model sheds light onto which subsets of beach recreators are likely to be impacted by different scenarios of resource change.

*I am grateful to the National Marine Fisheries Service and California Sea Grant for providing support that made this research possible. I thank Michael Hanemann, Guido Imbens, Jeffrey LaFrance, Cindy Thomson, Miguel Villas-Boas, Sofia Villas-Boas and seminar participants at the 2005 AAEA meetings for helpful comments and suggestions. I also thank Chris Busch, Koichi Kurisawa, David Layton, Craig Mohn, Linwood Pendleton, and Felipe Vasquez for their involvement with the Southern California Beach Valuation Project, which this project has greatly benefited from. All remaining errors are mine.

[†]Hilger is an economist at the Federal Trade Commission. Hanemann is a Chancellor's Professor at the Department of Agricultural & Resource Economics, University of California at Berkeley. This article was written independent of the author's duties at the FTC and thus expresses solely the views of the author and not those of any Bureau, any individual commissioner or the Commission. Federal Trade Commission, Bureau of Economics, 600 Pennsylvania Ave., NW, Washington DC, 20580; jhilger@ftc.gov

1 Introduction

Environmental resource protection and management requires the ability to assign values to non-market goods. While the literature has generally focused on the *average* valuation and preferences for these goods, the importance of the *distribution* of preferences for environmental amenities for populations with diverse preferences has often been neglected.

Not only is there a loss of welfare due to impaired use and enjoyment of the public beaches, but there can also be health impacts from swimming in polluted ocean waters, including upper respiratory infection and other illnesses.¹ This paper investigates the willingness to pay for a reduction in beach water pollution and illustrates how these values vary by recreator characteristics and season.

Varying preferences of recreational users and the multiple use nature of beach sites complicate the estimation of willingness to pay measures for improvements in water quality and other beach attributes. Systematic preference heterogeneity can lead to bias in parameter estimates if left unaccounted for. This paper addresses systematic preference heterogeneity by utilizing a finite mixture logit (FML) random utility model. A panel trip diary data set documenting 4,462 Southern California winter and summer beach trips for 595 recreators from December 2000 to November 2001 is analyzed using the FML approach. Estimation results indicate that beach recreators can be characterized by one of four distinct representative groups by beach recreational decisions, and individual and seasonal attributes. This information is then used in calculating welfare estimates for each individual in the sample and the weighted average measure for the population. I find that the welfare estimates associated with an environmental improvement vary significantly both within the population and across seasons.

1.1 Contributions of this Research

While FML models have been estimated previously in the environmental and resource economics literature this research makes three main contributions. First, this work is the first FML application focused on modeling the welfare and behavioral impacts of an environmental good associated with health outcomes. Recreational swimming is the second most popular recreation activity in the United States with over 90 million participants, and these recreators often swim in coastal and fresh water that does not meet the EPA health standards (NRDC, 2004). This drives an increasing interest in determining what draws recreators to specific beaches (Hanemann et. al., 2004, and Lew and Larson, 2005) and what influences where they choose to recreate once at a particular beach (Pendleton, 2001). This research furthers the understanding of the impact that water pollution has on beach recreation through the estimation of a recreation choice model for a diverse group of beach recreators. This model can be used to increase understanding of recreational beach choice and the composition of behavioral groups and as a forecasting tool in the resource and public health arenas.

Second, this application contributes to the modeling of heterogeneity with the FML through the incorporation of a seasonal variable in the beach choice occasion preference membership function. This enables the analyst to capture seasonal variation in preferences for beach attributes. Other studies have not utilized the model to account for seasonal changes in site attribute preferences.

¹A large epidemiological study, The Santa Monica Bay Restoration Project study, found an increase in the risk of contracting an illness when swimming near storm drains. Recreators that swam near storm drains were 57% more likely to suffer symptoms of a fever than other swimmers (Haile et. al., 1996). For a recent review of health risks associated with beach water pollution see NRDC (2005).

Lastly, this research represents a substantial step forward in the technique’s empirical application. Previous applications have been applied to fairly restrictive choice sets, primarily modeling binary participation choice or multinomial choice for up to 6 options (Boxall and Adamowicz, 2002). In contrast, this application models recreator decisions among a choice set of 51 beaches using a revealed choice data set. This model specification utilizes 9 individual trip membership function variables consisting of seasonal, activity participation, and demographic variables. This marks a substantial increase in the number of parameters estimated relative to other applications in the literature (Provencher et. al. use 3 (2002), Boxall and Adamowicz use 6 (2002), and Shonkwiler and Shaw use 3 (2003)).

Comparison of the welfare estimation results from the competing models indicate that the FML model provides an important insight into the heterogeneity of individual’s willingness to pay (mWTP) for improvements in water quality. The conditional logit and RPL mean mWTP estimates are bounded by the individual FML mWTP estimates for individual recreator trips, which are a function of recreator and trip characteristics.

The remainder of this paper is organized as follows. The remainder of this section further motivates the application and the model, respectively. Section 2 starts with an overview of discrete choice models, the challenges of modeling consumer heterogeneity within this framework, and several techniques that have been developed as solutions. This section concludes with a discussion of the finite mixture logit model. Section 3 describes the trip, beach site and recreator data and is followed by the presentation of the estimation results for the competing models and the corresponding welfare estimates for several beach closure and water quality degradation scenarios. Section 4 concludes.

1.2 Application Background

Coastal and marine health play an important role not only in the prosperity of the fisheries industry but also in the welfare of the communities which border the California coast and rely on the coastal environment for recreation and tourism.² Beach trips serve as a primary recreational activity for some and as a source of income for others.

However, coastal environmental resources are increasingly strained and affected by pollution and overuse. In California there were 6,568 beach closures and advisories in 2001. This was a 14% statewide increase from 2000 and marked the fifth consecutive year that beach closures and advisories have increased (NRDC, 2002). The public awareness of poor water quality is so widespread in the Los Angeles area that, in a focus group a few years ago, eight out of ten participants said that they do not go into the water when they go to local beaches (Hanemann, 2005).

The risk of becoming ill while swimming at the beach reduces the welfare of those who venture in the water and contract an illness,³ and diminishes the welfare for those who forgo swimming because of the risk.⁴ Public concern regarding this coastal degradation has prompted several studies focusing on the adverse health effects of coastal pollution and has generated the approval of several Legislative and Assembly bills (NRDC, 2002).

The differences in the values placed on beach recreation by different user groups can have important practical implications for beach management. For example, sunbathers may place importance on different

²Lew and Larson estimate the mean value of a recreational beach day to be \$28.28 (2004).

³Rabinovici et. al. (2004) review the valuation of health status literature and report that Mauskopf and French (1991) estimate the WTP for government programs to aid in the avoidance of gastrointestinal symptoms at \$280 for a 2-4 day case and \$1,125 for a 5-7 day case.

⁴Walsh et. al. (1992) report a mean value per visitor day of recreational swimming at \$35.60 (\$2001)

characteristics of the beach than surfers or mothers with young children; what is considered an amenity to one may be unimportant or possibly an unwanted nuisance to others. Moreover, the resource manager may be forced to make trade-offs in meeting the needs of different groups.

Coastal resource managers should find it useful to have welfare measures of the values associated with the alternative uses of harbors, piers, and docks since beach recreation values may swing the direction of the overall coastal management plan. The model developed through this research will facilitate the implementation of balanced and equitable resource management through the increased understanding of taste differences across users.

Given that different user groups value different characteristics of recreation sites and demand different services from them it is useful to be able to account for the variation in preferences among different user groups. Accurate welfare and usage estimates can serve as a useful tool to resource managers concerned with understanding the equity implications of specific policies. Furthermore, robust welfare and usage estimates are increasingly called upon in a litigation setting for the enumeration of damages caused by resource degradation. The development and implementation of methods and techniques used to capture and control for heterogeneity is the key objective of this paper.

1.3 Modeling Background

A rich diversity of preferences among decision makers challenges the accurate modeling recreational site choice and estimation of the economic value associated with a change in resource attributes. Diverse user groups often value different attributes of recreation sites and demand different services from them. If preference heterogeneity can be easily controlled by segmenting the sample population by a variable known to the analyst, a standard logit random utility model (RUM) can be used to estimate coefficients and welfare measures for each group separately. For example, beach recreators who swim in the ocean are likely to have different preferences for water quality and other beach attributes than those lying on the sand. However it is often unclear where to draw the line in defining sub-samples of the population.⁵ This may lead to bias in welfare measures for changes in site attributes and hinder proper aggregation of welfare measures across individuals or time periods and adversely affect policy and management decisions.

The logit model handles variation in preferences by averaging over the individuals. In cases where the population is fairly homogenous in their preferences this may not cause a major problem; however, if the population is characterized by considerable systematic preference heterogeneity, the model's results may be misleading due to an averaging out effect over preferences from distinct groups. Additionally, the distribution of preferences over individuals or time is commonly lost due to the restrictive single point or modal distribution which the model imposes on the data. The preservation of the preference distribution may aid analysts in the estimation of the welfare impact to individual users and those between user groups which arise from changes in choice set.

Suppose there are two different groups of beach users: sunbathers and surfers, who prefer calm water and large waves respectively. Membership in either group is unobservable to the analyst, but may be statically correlated with observable demographic and seasonal data. Imagine further that there are several means of undertaking a coastal project which can have the secondary effect of impacting wave size. Estimation with

⁵If the analyst differentiated between beach and water users there would still be heterogeneity within users. For example, among water users, surfers may care about different aspects of the beach recreation experience than mothers taking young children to the beach to swim. Although both of these groups likely view clean water as desirable, they may differ in the level of importance they place on water quality. Additionally, users within primary groups may often have preferences for a subset of attributes that closely resemble those of another group.

the standard logit model causes the opposing preferences for waves of the two user groups to be averaged out, resulting in model estimates that call for the medium sized waves, which are not preferred by either group. In contrast, a model that could account for the preference heterogeneity between the two user groups through the estimation of separate preferences could lead to a policy of a variety of waves being maintained at specific beaches, resulting in a welfare improvements for both groups.

The finite mixture logit (FML) model implemented in this paper accounts for systematic heterogeneity in recreator preferences by estimating the probability of latent segment behavioural group membership for individual recreators.⁶ Where a latent segment type is a construct of a preference traits and is termed "latent" since individual membership in a particular segment, as well as the segments themselves, are not observable. Estimation of the FML model simultaneously generates the probability of membership to each latent segment and the choice probabilities for each agent's choice occasion. This approach captures the variation in preferences across the population through modeling them as an unrestricted discrete distribution. The model is distinctive in that it not only accounts for heterogeneity, but it provides insight into the composition of the primary behavioural groups.

2 Random Utility Models

2.1 Basic Framework

Random utility models have a long history as a powerful tool for resource managers. The random utility model is the standard statistical framework used to estimate the value of the change in consumer welfare due to an incremental change in the level of resource attributes in a setting characterized by consumer choice between several alternative sites with varying attributes.⁷

Consider the utility maximization problem that an individual solves in relation to a choice occasion between a set of J alternatives ($j = 1, \dots, J$):

$$Max_j : u_i = v_i(M_i - C_{ij}, Q_j, Z_i) + \epsilon_{ij}. \quad (1)$$

Where $u(.)$ is a function of individual income, M_i , the cost of individual i visiting site j , C_{ij} , the quality and attribute mix of the chosen site, j , in the choice set, Q_j , and individual socioeconomic characteristics, Z_i . The unobservable portion of utility is denoted by ϵ_{ij} and is assumed to be a random variable. The decision to visit a particular area is viewed as the decision to consume, or incorporate into one's utility function, the specific attributes that uniquely identify the chosen site from others in the choice set. When individual i chooses to consume bundle j out of her choice set J , $j \in J$, it is assumed that u_{ij} is the maximum of the J possible utilities in the choice set. The conditional probabilities of individual i choosing choice j can be derived as

$$Pr_{ij} = \Pr(u_{ij} > u_{ik}) \forall k \neq j. \quad (2)$$

The outcome of an individual choice occasion, designated by Y_i , is a random variable. If and only if the disturbances associated with j , $\forall j \in J$, are independently and identically distributed with the generalized

⁶This model was first proposed by McFadden (1986) and implemented by Swait (1994).

⁷The Conditional Logit Random Utility Model (CL RUM) is a widely used research tool. An early application of this model to recreational choice application is Hanemann (1978). For technical discussions refer to Greene (2000) and Wooldridge (2002). For a discussion of the application of RUMs to environmental economics refer to Haab and McConnell (2003).

extreme value distribution, $F(\epsilon_{ij}) = \exp(-e^{-\epsilon_{ij}})$. The choice probabilities are

$$\Pr_{ij} = \Pr(Y_i = j) = \frac{e^{\beta' \Gamma_{ij}}}{\sum_{j=1}^J e^{\beta' \Gamma_{ij}}} \quad (3)$$

where Γ_{ij} is a vector of individual and alternative specific variables (McFadden, 1973). This model is known as the conditional logit model (CL). This formulation of the CL model causes individual variables, Z_i , that do not vary over the choice set to drop out of the choice probability. The choice probability is then determined by choice specific variables.⁸

The parameters of the indirect utility function, $v_i(\cdot)$, can be estimated using maximum likelihood techniques.

$$\Pr_{ij} = \frac{e^{v_i(C^j, Q_j)}}{\sum_{j=1}^J e^{v_i(C^j, Q_j)}} \quad (4)$$

Several econometric and modeling issues commonly arise with the Random Utility Model. Econometric consideration should be given to the independence of irrelevant alternatives property⁹ and to identification issues surrounding the scaling parameter. In terms of modeling, the construction of the travel cost variable¹⁰ and the formation of the choice set are major issues that have been the focus of considerable research.¹¹

2.2 Econometric accounting of Heterogeneity

The behavioral response and economic value associated with a change in resource characteristics can vary over individuals due to the rich diversity, or heterogeneity, among individual decision makers. Heterogeneous preferences are difficult to account for in behavioral choice models due to the formulation of the conditional logit (CL) model which has historically been the base tool for random utility models. Within demand system models, the analyst can directly incorporate demographic, temporal, or other individual characteristic data directly into the individual's utility function to address preference heterogeneity. However under the specification of the CL, these characteristics drop out of the probability of an individual selecting a specific choice, thus preventing the direct identification of these characteristics in the model.

If heterogeneity is not accounted for, RUM estimates are characterized by bias and lead to inaccurate forecasts pertaining to changes in resource attribute levels and management policies (Chamberlain, 1980; and

⁸However, through construction of interaction variables that vary over both individuals and choice attributes, individual specific information can be retained as an argument in the choice probabilities.

⁹The analyst must take note that in the standard multinomial or conditional logit models the odds ratios for a specific pair of choices, \Pr_j / \Pr_k , is independent of the remaining alternatives. This property is known as independence of irrelevant alternatives (IIA). This property is fairly restrictive because it implies that the relative probability of choosing between alternatives remains constant after the introduction or removal of a perfect substitute of one of the alternatives to the choice set. Several models such as the nested logit and random parameters logit models have been developed, in part, as a solution to IIA (Haab and McConnell (2003)).

¹⁰The assumption that travel cost prices are exogenously determined deserves comment, as the endogeneity in prices assumption is one of the primary issues critiqued in the discrete choice literature (Berry, Levinsohn, and Pakes, 1995; and Nevo, 2000). However as discussed in Train (2003), this issue is not of great importance outside of market-level demand models. Within customer-level demand models it is assumed that individual demand does not affect price. Moreover, within the recreational demand literature the price associated with choosing a specific good is determined by the cost of travel to that location. One alternative is that the consumption of the good is of large enough proportion in the individual's utility function that the individual incorporates the location of the recreational site as an important argument in the residential location decision making process. Secondly, site characteristics to some degree all relate to visitation. For most site attributes individual trips do not affect the attribute level. However, some attributes, such as solitude, offered by the site are highly sensitive to small changes in the number of trips taken to the site. Assuming that individual residence location and travel infrastructure is determined exogenously, the travel cost price is exogenous. See Parsons (1991) for a discussion on housing location.

¹¹For a thorough review on the optimal size of the choice set see Kurisawa (2003).

Jones and Landwehr, 1988). This bias adversely affects welfare estimates for simulated changes in resource attributes and/or management decisions.

Early work addressing heterogeneity focused on structural approaches requiring the *a priori* selection of typically demographic or choice variables. In "cluster models" individuals are segmented into demographically homogenous/similar groups. An alternative method incorporates into the indirect utility function and interaction variable composed of individual demographic variables, such as income and various choice attributes (Adamowicz et al., 1997). These methods are limited by the assumption that preference groups can be accurately determined *a priori* by demographic variables, and theoretical issues pertaining to the choice of an interaction variable (Boxall and Adamowicz, 2002). Other related solutions to this problem include the fixed effects and random effects specification of the conditional logit model (McFadden, 1986). However, these methods are difficult to employ when the heterogeneity structure is complex and the sample consists of a large number of decision makers.

An additional structural method, the Generalize Extreme Value (GEV) Logit (or nested logit) disaggregates the decision between alternatives into subsets of similar alternatives, relaxing the IIA restriction (McFadden, 1978). In the context of beach recreation, the GEV framework has been used to model recreational beach choice conditional on the type of activity engaged in during the beach visit (Hanemann et. al., 2004). The primary benefits of this approach are that the model may be useful in highlighting the differences in choice behavior and welfare estimates for different user groups and the relaxation of the IIA property. However, the approach requires that the "nesting" rules are defined *a priori* by the analyst.

Another approach, the random parameter logit (RPL), controls for heterogeneity across preferences by allowing estimated coefficients to randomly vary across individuals according to a continuous probability distribution, typically the normal or log-normal. By allowing for variation in coefficients over people, the unobserved portion of the respondent's utility is correlated over sites and time (Train, 1997). Additionally, the RPL model is not restricted by the IIA property due to interactions within the choice probabilities of the attributes of all elements in the choice set (Train, 2003). The RPL approach has two weaknesses. First, it assumes that preferences vary continuously across economic agents. Second, it does not offer a behavioral explanation for the source of the heterogeneity across people. Although the continuous distribution assumption is likely to be valid in many applications, there are situations where actual preferences may be more accurately estimated by a less restrictive distribution with multiple probability masses. When preferences differ sharply between user groups, for instance the importance of water quality for swimmers versus cyclists on a beachside bikepath or the presence of motorized watercraft for users and non-users, a model which provides coarse grouping of preferences may provide more accurate behavioural forecasting and welfare estimates.

2.3 Finite Mixture Logit Approach

An alternative solution is the finite mixture logit (FML), or latent segmentation approach which simultaneously accounts for heterogeneity, models preference composition and site choice. This approach was suggested in a RUM setting by McFadden (1986), and was implemented by Swait (1994). There has been a recent increase in the application of this approach, including several recreational choice models applications (Provencher et. al (2002), Boxall and Adamowicz (2002), and Shonkwiler and Shaw (2003). Use of the FML approach is motivated by two primary assumptions. First, individual preferences are neither homogeneous nor continuously distributed, but can vary between population segments which can be represented by discretely distributed multiple probability mass points. Second, variation in preferences between user groups

is not purely a function of demographic variables, but of perceptions, attitudes, behavior, past experiences, and unobserved variables. Utilization of the FML approach allows the exploration of preference variation across individuals conditional on the probability of membership to a latent preference group. The gained explanatory power can be exploited by managers in terms gains in use forecasting and welfare analysis.

Each "latent segment" represents like-minded individuals with homogeneous preferences. The segments are termed latent because individual membership to a particular segment is not observable, nor are the segments themselves. The FML model simultaneously assigns the economic agent the probability of membership to each latent segment and estimates the discrete choice probability for the random utility model. This approach captures the variation in preferences across the population through a discrete distribution with multiple probability masses. The model is unique in that it not only accounts for heterogeneity, but is able to explain the sources of that heterogeneity. This is of particular importance in regards to management decisions where user groups may either be demographically homogenous or where there is little correlation between user group preferences and the standard demographic variables. The FML model additionally estimates the composition of the latent segments and can be used to help researchers and managers understand the processes involved in the formation of behavioral groups.

The FML RUM is an extension of the CL model, and follows the assumption that individual i 's indirect utility is maximized on a choice occasion by selecting alternative $j \in J$. The probability that alternative j is chosen is the probability that the utility gained from choice j is greater than or equal to the utility forgone by not picking one of the other alternatives in the choice set, J .

Under the assumption that there exists some degree of heterogeneity in preferences across the sample, let S be the number of segments that the population is to be grouped into.¹² Individuals are assumed to belong to a segment $s(s = 1, \dots, S)$ within the sample population. Individuals within a segment are assumed to be characterized by homogeneous preferences. Additionally, in all but the trivial case, $S = 1$, the probability ratio between any two alternatives includes arguments from all other alternatives in the complete choice set, J . It has been shown that in these cases the FML model is not constrained by the IIA property.(Shonkwiler and Shaw, 2003).

2.3.1 Single Choice Occasion

In a cross sectional data setting, the optimal solution to the choice decision for individuals represented by a given segment s , is to maximize

$$u_{i|s} = v(\beta_s X_{ij}), \quad (5)$$

where the β_s vector is the coefficients representing individual preferences conditional on individual i 's membership in segment s .

The parameter coefficients for a specific segment of the population are estimated using the following probabilities.

$$\Pr_{ij|s} = \frac{e^{v_{i|s}(-Cji, Q_j)}}{\sum_{j=1}^J e^{v_{i|s}(-Cji, Q_j)}}. \quad (6)$$

Consider a latent membership likelihood function M^* that assigns individuals to segment $s \in S$ (Swait,

¹²The optimal choice of S is discussed below.

1994). Arguments to M^* can include variables associated with the unobservable tastes, attitudes, and preferences of the members of the group, socioeconomic variables, and characteristics of the choice occasion represented by the vector Z_i . Segments can be identified using standard demographic variables, behavioral and preference data, and choice occasion specific or temporal data. Assume the following equation:

$$M_{is}^* = \gamma_s' Z_i + \zeta_{is}, \quad s = 1, \dots, S, \quad (7)$$

where γ_s is a vector of segment specific parameters and ζ_{is} represents the error terms.

The membership likelihood function, M^* , is a random variable. To use the function in an econometric model requires assumptions about the distribution of its error terms. Following Kamakura and Russell (1989), Swait (1994) and Boxall and Adamowicz (2002) the error terms are assumed to be independently distributed across individuals with Type I extreme value distribution. The probability of individual i belonging to segment s can then be estimated utilizing a multinomial logit framework where the independent variables in this function vary over individuals, unlike the conditional logit where the variation is in the choice specific variables. Addressing an indeterminacy in the model caused by the lack of normalization the following restriction must be imposed:

$$\pi_{is} = \frac{e^{\gamma_s' Z_i}}{1 + \sum_{s=2}^S e^{\gamma_s' Z_i}} \quad \text{for } s = 2, \dots, S, \quad (8)$$

$$\pi_{i1} = \frac{1}{1 + \sum_{s=2}^S e^{\gamma_s' Z_i}} \quad \text{for } s = 1, \text{ and} \quad (9)$$

$$0 \leq \pi_{is} \leq 1, \text{ such that } \sum_{s=1}^S \pi_{is} = 1.$$

To model choice behavior under the assumption that the sample population can be represented as a weighted average of a finite number of representative segments, the researcher estimates individual i 's utility maximizing choice between J alternatives conditional on membership to a specific segment, s . The joint probability \Pr_{ins} that an individual i is a member of segment s , and chooses alternative j for all $s \in S$ and $j \in J$ is defined as

$$\Pr_{ijs} = \pi_{is} \Pr_{ij|s}. \quad (10)$$

It follows that for a single choice occasion the probability of individual i choosing alternative j unconditional on segment membership can be written as

$$\Pr_{ij} = \sum_{s=1}^S \pi_{is} \Pr_{ij|s}. \quad (11)$$

Defining d_{ij} as an indicator variable that takes the value of 1 if an individual i chooses site j and 0 if not, allows the writing of the individual likelihood function as

$$L = \sum_{s=1}^S \left[\pi_{is} \left(\prod_{j=1}^J \Pr_{ij|s}^{d_{ij}} \right) \right]. \quad (12)$$

The individual likelihood function can be rewritten as

$$L = \prod_{j=1}^J \left[\sum_{s=1}^S \left(\pi_{is} \Pr_{ij|s} \right) \right]^{d_{ij}}, \quad (13)$$

which yields the log likelihood function for cross sectional data

$$\ln L = \sum_{i=1}^I \sum_{j=1}^J d_{ij} \ln \left[\sum_{s=1}^S \pi_{is} \Pr_{ij|s} \right]. \quad (14)$$

2.3.2 Segment Membership Time Consistency

The extension of the single choice occasion likelihood function to incorporate a time dimension utilizing panel data introduces a few complications in terms of the assumptions of segment membership independence across choice occasions. One assumption is that preferences are constant over time, although there is preference heterogeneity across individuals. A second modeling assumption is that preferences can be allowed to vary both over individuals and time.

Constant over Time FM Membership In the constant over time framework individuals agents are modeled to be characterized by the same preference segment for all choice occasions. The constant over time assumption is most appropriate when the set of choice occasions are temporally close (such as multiple decision choice occasions), or when preferences and choice attributes are stable over time. This specification has been applied in both the marketing and transportation literature (Ramaswamy et al., 1999, Greene and Hensher, 2003). Following this assumption the probability of individual i choosing the set of alternatives j at each time t over the set T choice occasions is

$$\Pr_{ijt} = \sum_{s=1}^S \pi_{is} \left(\prod_{t=1}^T \Pr_{ijt|s} \right).^{13} \quad (15)$$

This assumption appears plausible in cases where the set of choice occasions are short in time duration, where the population segment's characteristics are constant over time, when the arguments of the segment membership function do not vary over time, and when there may be one choice occasion that is made up of several individual decisions.

Variation over Time FM Membership An alternative modeling specification, variation over time, can be useful as preferences often tend to vary with seasonal tastes as the underlying choice decision changes. This assumption is implemented in this paper and assumes that preferences can be allowed to vary both over individuals and time. Allowing for variation over time in preference membership relaxes the correlation between individual segment membership.

Seasonal variation in unobserved or unmeasured attributes necessitates the need to allow for seasonal variation in the segment membership function, allowing individual segment membership to change over time. For example, the surf is generally better in the winter and the weather is warmer in the summer. This may result in a winter preference set that gives high weight to water quality and surf variables, and a summer preference set that base choice on attributes that are important to sun bathers.

¹³Note this can also be written as $\Pr_{i\varphi}$, where φ is a vector of length that represents the sequence of site choices over time T .

On a shorter time scale, allowing for variation between individual segment membership helps the model capture the correlation between segment membership between time periods. If a respondent was in a very active preference group during one period (swim), and they go to the beach the next day they are more likely in the second period to be in a more sedate group (lie on sand). In this case, segment membership is a function of both the previous segment classification and the time elapsed since the last choice occasion. Serial correlation of this type has been investigated in the marketing (Haaijer and Wedel, 2000) and recreational fishing literature (Provencher et al, 2002).

Write the probability of individual i choosing alternative j at time t as

$$\Pr_{ijt} = \sum_{s=1}^S \pi_{is} \Pr_{ijt|s} . \quad (16)$$

This leads to the likelihood function

$$L = \prod_{i=1}^I \prod_{t=1}^T \left[\sum_{s=1}^S \pi_{ist} \left(\prod_{j=1}^J \left(\Pr_{ijt|s} \right)^{d_{ijt}} \right) \right] \quad (17)$$

which simplifies as

$$L = \prod_{i=1}^I \prod_{t=1}^T \prod_{j=1}^J \left[\sum_{s=1}^S \pi_{ist} \Pr_{ijt|s} \right]^{d_{ijt}} , \quad (18)$$

and leads to the log likelihood function¹⁴

$$\ln L = \sum_{i=1}^I \sum_{t=1}^T \sum_{j=1}^J d_{ijt} \ln \left[\sum_{s=1}^S \left(\frac{e^{\alpha \gamma'_s \mathbf{Z}_{it}}}{\sum_{s=1}^S e^{\alpha \gamma'_s \mathbf{Z}_{it}}} \right) \left(\frac{e^{\beta'_s \mathbf{X}_{ijt}}}{\sum_{j=1}^J e^{\beta'_s \mathbf{X}_{ijt}}} \right) \right] . \quad (19)$$

The above likelihood function has been utilized in both the marketing (Swait, 1994) and recreation (Boxall and Adamowicz, 2002) literature. Both applications utilized stated preference data where each respondent made a series of sequential choices from a structured choice experiment where all choice decisions are made at the same time, weakening the basis for the preference variation over time assumption. The basis of the FML is that decisions made by different members of the same preference segment will be more correlated than decisions made by members of different segments.¹⁵ This holds true unless there is a mechanism for an individual's segment membership to change between choice decisions (Morey, 2003).

The choice of time specification is dependent on the goals of the analysis and what data is used. As a general rule, the constant over time specification is appropriate for models over short time durations which do not utilize membership covariates that vary over time and where preferences are assumed to be constant. The varying over time specification better suits applications that seek to model FM membership as a function of seasonality, the effect of previous choices, or individual characteristics that vary over time (the decision to get into the water on a specific beach trip). This paper utilizes the varying over time specification, as individual preferences are expected to vary over time due to both seasonal effects and variety seeking throughout the survey year. It is noted that the constant over time model specification can be implemented by restricting the time varying individual characteristic variable parameters to zero. The consistency over time of estimated

¹⁴Note the individual demographic variables, \mathbf{Z}_{it} , have a time index.

¹⁵This assumes that the information set and individual characteristics are constant across choice decisions.

individual recreator segment membership is discussed with the application estimation results.

2.3.3 Additional Econometric Issues

Scale Parameter In addition to attribute preference parameters, the variance of the disturbance terms may also differ across segments of the population. In the standard CL framework the analyst assumes that the unobserved factors have constant variance, hence utility is of the same scale across respondents. However, this restriction is not implicitly held in the FML specification. Therefore FML model parameter estimates cannot be compared across segments directly. Researchers that do not take the differences in scaling parameters into account may incorrectly infer that the members of the segment with a larger coefficient estimate care about the attribute more than those individuals in the other segment. To properly interpret parameter results across segments analysts can compare the signs or ratios of parameter point estimates.¹⁶

Determining the Number of Segments The appropriate number of segments is not identifiable in the FM class of models and is treated as exogenous. However, one can statistically test for modeling improvements conditional on the number of segments by estimating a series of models which iteratively increase in S . Modeling improvements due to changes in the number of latent segments defined in the model can be tested for through the use of McFadden's ρ^2 , Bayesian Information Criterion, and Akaike Information Criterion test statistics. The use of traditional Likelihood Ratio tests in determining the number of segments should be used with caution as the regularity conditions are violated (Ben-Akiva and Swait, 1986, Jedidi, 1997, and Boxall and Adamowicz, 2002). In addition to the statistical tests, the analyst's judgment in regards to which model specification in terms of the number segments best describes the respondent population and addresses the relative policy questions should be applied.

Upon inspection of the FML model it is clear that through the selection of the appropriate number of segments the above model can mimic both the traditional CL and the RP models.¹⁷ For instance, when $\gamma_s = 0, \beta_s = \beta, u_s = u, \forall s$, the FML reduces to the CL.

3 Welfare Estimation

The generation of welfare measures associated with a change in the attributes of the choice alternatives is a primary use of the RUM. The economic marginal value of site attributes and the compensating variation measure of consumer surplus associated with changes in site choice characteristics, such as water quality grades, can be calculated for each segment membership group using model parameter estimates. The marginal value measure offers a readily assessable rule thumb welfare measure for changes in quality attributes. Whereas the compensating variation measure of consumer surplus takes into account the substitution patterns associated with a change in the choice set.

The FML model provides a framework for the calculation of willingness to pay measures associated with changes in the choice set attributes using parameter estimates for each membership segment. The resulting willingness to pay calculations provide a detailed estimate of the willingness to pay distribution of individual welfare measures calculated through a weighted average of their component latent representative consumers.

¹⁶Alternatively, the scaling parameter can be normalized for one segment so that the variance of the disturbance term is the same across both segments. This leads to the identification of the scaling parameter (Train, 2003).

¹⁷In the present form FML is theoretically similar to the RPL where each respondent undertakes one choice occasion.

Marginal Value Measure Changes in welfare due to a marginal change in a given attribute can be calculated using the marginal mean willingness to pay measure (mWTP). This measure is defined as the maximum amount of income a person will pay in exchange for an improvement in the level of a given attribute provided and can be calculated as:

$$mWTP_i^* = \frac{\beta}{\gamma} \quad (20)$$

where β is the parameter on the attribute of interest and γ is the travel cost parameter. Both parameters measure the marginal utility of the object in question. This result can easily be applied using FML parameter estimates:

$$mWTP_{i|s}^* = \frac{\beta_s}{\gamma_s} \quad (21)$$

Because the degree of heterogeneity in preferences is assumed to be considerable in many recreational choice optimization problems, the ability to segment the changes in welfare over latent user types is important. However if the resource managers are interested in aggregate welfare measures over the sample, these can be calculated by adding up the welfare measures weighted by the latent segment probability (Boxall and Adamowicz, 2002).

$$mWTP_{ji}^* = \sum_{s=1}^S \pi_s \left[\frac{\beta_s}{\gamma_s} \right] \quad (22)$$

Compensating Variation Measure of Consumer Surplus Changes in welfare due to the attribute/quality mix of the chosen bundles on one choice occasion can be calculated using the compensating variation measure and the estimated parameters of the indirect utility function (Small and Rosen, 1981; Hanemann 1982).

$$v_i(M_i - Cji - CV_{ji}^*, Q_j^1, Z_i) + \epsilon_{ji} = v_i(M_i - Cji, Q_j^0, Z_i) + \epsilon_{ji} \quad (23)$$

This results in the per trip marginal change in welfare due to a decrease in some site attributes.

$$CV_{ji}^* = \frac{\ln \left[\sum_{j=1}^J e^{v(\beta Q_j^1)} \right] - \ln \left[\sum_{j=1}^J e^{v(\beta Q_j^2)} \right]}{\gamma} \quad (24)$$

This result is readily extended for use with FML parameter estimates. Analysts interested in the welfare effect to specific groups can generate welfare measurements for an arbitrary change in choice set attributes for each latent segment though the use of parameter estimates for the segment of interest.

Because the degree of heterogeneity in preferences is assumed to be considerable in many recreational choice optimization problems, the ability of segmenting the changes in welfare over latent user types is important. However if the resource managers are interested in aggregate welfare measures over the sample, these can be calculated by adding up the welfare measures weighted by the latent segment probability (Boxall and Adamowicz, 2002).

$$CV_{ji}^* = \sum_{s=1}^S \pi_s \left[\frac{\ln \left[\sum_{j=1}^J e^{v(\beta_s Q_j^1)} \right] - \ln \left[\sum_{j=1}^J e^{v(\beta_s Q_j^2)} \right]}{\gamma_s} \right] \quad (25)$$

Utilization of the FML for welfare analysis provides an improvement over the traditional welfare calculation using the logit and RPL models. Choice attribute and membership variable coefficients can be used to estimate the appropriate mWTP and CV welfare measures for each choice occasion and the distribution of these measures as a function of individual and trip characteristics.

4 Data

The empirical choice model application utilizes an extensive recreational panel data set for recreational beach trips to 51 Southern California beaches (Table 1). The data come from a survey of households in Southern California. Respondents were asked to keep a diary of all their trips to beaches in Southern California from December 2000 through November 2001. The data consists of observation over a 12 month period for 4,642 beach recreation choice occasions of 595 beach recreators living in Southern California (Figure ??). Recreators include fishers, boaters, divers, surfers, sunbathers, runners, cyclists and other beach users. Beach recreator data contains demographic and behavioral data. An attribute data set contains individual beach attributes including water quality data and the travel times and distances between each beach and respondent residence.¹⁸ The CL and RPL models are estimated using the same choice probability specification as the FML model.

Modeling individual site choices for beach recreation requires explanatory variables in terms of how the beaches in the choice set differ from one another. Beach attributes incorporated into the model specifications include beach location, water quality, presence of children's playgrounds, restaurants, tide pools, rest rooms, and foot or bike paths. The choice set for the complete panel consisted of 304 beaches, which were then aggregated into a set of 53 beaches. Properly defining the choice set is of great importance in model estimation. This is increasingly important when dealing with large choice sets. To help address these issues, respondents were asked questions to determine their familiarity and subjective quality opinions of the beaches included in the complete choice set. Summary statistics are included in Table 2.

Beach sites that had zero trips, and low name recognition are kept in the choice set. This decision is made based on the observation that these beaches are within close proximity to other beaches in the choice set that were visited. Actual beach choice often includes a degree of search; one may know the general area that they wish to visit, but their final choice is not made until a degree of "window shopping" is undertaken.

Each respondent is assigned a unique numeric identifier in order to link survey responses from all segments of the project and thus create a large panel data set. The screener and recruitment surveys collect standard socioeconomic household data, as well as beach and non-beach recreation data. Respondents were asked to

¹⁸The complete data set consists of a screener and recruitment survey, 6 bi-monthly diary surveys, and 7 supplementary modules that focus on a variety of topics. The original data set comes from a random telephone sample of 1,848 respondents. Of these, 824 respondents were classified as non-beach users and 202 declined to take part in the survey. The remaining 822 respondents agreed to be included in a large panel data set. Analysis shows that the demographics of the final sample is similar to those who declined to participate and therefore it is assumed that there is not a substantial amount of systematic self-selection bias. For a thorough discussion of the data see Hanemann et. al. (2003).

Table 1: Beach Sites in Study

1	San Onofre South	18	Bolsa Chica	35	Mother's
2	San Onofre North	19	Sunset	36	Venice
3	San Clemente State	20	Surfside	37	Santa Monica
4	San Clemente City	21	Seal	38	Will Rogers
5	Poche	22	Alamitos Bay	39	Topanga
6	Capistrano	23	Belmont Shores	40	Las Tunas
7	Doheny	24	Long Beach	41	Malibu (Surfrider)
8	Salt Creek	25	Cabrillo	42	Dan Blocker (Corral)
9	Aliso Creek	26	Point Fermin	43	Point Dume
10	Laguna	27	Royal Palms	44	Free Zuma
11	Crystal Cove	28	Abalone Cove	45	Zuma
12	Corona Del Mar	29	Torrance	46	El Matador
13	Balboa	30	Redondo	47	La Piedra
14	Newport	31	Hermosa	48	El Pescador
15	Santa Ana River	32	Manhattan	49	Nicholas Canyon
16	Huntington State	33	El Segundo	50	Leo Carrillo
17	Huntington City	34	Dockweiler	51	County Line

Figure 1:

Table 2: Respondent Beach Site Familiarity

	Beach	Obs.	Familiar? Number saying yes	Familiar? Percentage saying yes
1	San Onofre South	473	73	15%
2	San Onofre North	482	128	27%
3	San Clemente State	481	174	36%
4	San Clemente City	475	99	21%
6	Capistrano	484	185	38%
7	Doheny	483	162	34%
8	Salt Creek	484	76	16%
9	Aliso Creek	483	91	19%
10	Laguna	484	371	77%
11	Crystal Cove	483	133	28%
12	Corona Del Mar	484	235	49%
14	Newport	484	399	82%
16	Huntington State	484	339	70%
17	Huntington City	469	280	60%
18	Bolsa Chica	482	194	40%
19	Sunset	482	175	36%
21	Seal Beach	484	294	61%
22	Alamitos Bay	484	54	11%
23	Belmont Shores	484	191	39%
24	Long Beach	483	308	64%
25	Cabrillo	480	192	40%
27	Royal Palms	473	49	10%
28	Abalone Cove	473	69	15%
29	Torrance	482	117	24%
30	Redondo	484	350	72%
31	Hermosa	484	281	58%
32	Manhattan	484	262	54%
33	El Segundo	483	128	27%
34	Dockweiler	484	77	16%
35	Mother's	484	265	55%
36	Venice	484	352	73%
37	Santa Monica	482	382	79%
38	Will Rogers	482	147	30%
39	Topanga	482	92	19%
40	Las Tunas	479	51	11%
41	Malibu (Surfrider)	479	207	43%
42	Dan Blocker (Corral)	479	30	6%
43	Point Dume	479	74	15%
45	Zuma	479	130	27%
49	Nicholas Canyon	479	22	5%
50	Leo Carrillo	482	143	30%
51	County Line	482	85	18%

Table 3: Probability of Water Recreation by Season

	Trips	Trip %	Recreators	Recreator %	Avg Individual Seasonal %
Total	4,642	27%	595	23%	22%
Winter	987	14%	222	6%	9%
Summer	1,749	38%	378	62%	30%
Shoulder Season	1,906	23%	377	58%	22%

Table 4: Seasonal and Water Recreation Beach Trip Counts

Dry Trips by Season				
	Total	Winter	Shoulder	Summer
Min	0	0	0	0
Avg	67	17	29	21
Max	492	214	154	124
Std Dev	267	119	82	66
Total	3,409	850	1,474	1,085

Wet Trips by Season				
	Total	Winter	Shoulder	Summer
Min	0	0	0	0
Avg	24	3	8	13
Max	208	32	66	110
Std Dev	114	18	36	60
Total	1,233	137	432	664

keep a record of every Southern California beach trip in a bi-monthly diary throughout the survey period.¹⁹ For each trip, respondents were asked a series of trip details including the date of the trip, the specific beach they went to, the number of minors in their group, and information about up to four beach activities. Beach recreational activities are expected to be affected by seasonal variables. To control for this effect the data set is split into three time periods: winter (December and January), summer (June through September) and the remaining shoulder season months. Summary statistics on the seasonal distribution of trips, the probability of the average beach recreator's immersion rate, the percentage of trips that involves water immersion, and the proportion of recreators that enter the water are listed in Table 3.²⁰ Summary statistics on beach site trip counts are displayed in Table 4.

The implicit price of visiting each beach used in modeling is the travel cost construct. This construct

¹⁹Individual recreators frequented several beaches. 73% of all beach trips were to the recreator's most frequently visited beach.

²⁰Due to multiple site trips or inconsistencies among the screener, recruitment, and diary surveys 14.2% of the trip observations have been dropped from the dataset. Multiple site trips make up 3.9% of the dataset and have been excluded from this analysis due to complications in capturing the percentage cost of travel from one beach to another for the price matrix, and the proper weighting of beach attributes. Multiple site trips are commonly handled in the literature by assuming that they are independent trips.

is a function of the respondents reported income, and the estimated vehicle operational cost (\$0.145/mile), travel time and the distance between the respondent's residence and each beach in the choice set.²¹ One way travel distance and travel time between a respondent's address and the beach address are calculated using the computer program PC-Miler. The time and distance data is transformed into the round trip travel cost of each trip, and is one of the model's primary explanatory variables.²² See Table 5 and for average round trip costs to each beach recreation site.

Beach water pollution data is obtained from Heal the Bay, a Southern California non-profit group. This data contains weekly ratings on a scale of A+ to F for beach water quality for dry days at many monitoring stations throughout Southern California between June 1998 and April 2001. The A+ to F ratings are based on three biological pollutants measures: total coliform, fecal coliform, and enterococcus. The presence of these pollutant are indicators of several illnesses such as stomach flue, ear infection, upper respiratory infection, and skin rashes. The calibration of the A+ to F scores are set at levels where a D score caused by a high fecal coliform ratio is associated with a 1 in 85 chance of becoming ill; and D water caused by enterococcus is associated with a 1 in 77 chance of becoming ill. Beaches that are rated as "Failing" with an F score caused by a high fecal coliform ratio are associated with a 1 in 20 chance of becoming ill (Heal the Bay, 2005).

Table 5: Beach Site Details: Cost, Water Quality, Trips

	Beach	Avg. Cost	Avg. Water Grade	Observed Trips
1	San Onofre South	\$6.91	4.0	34
2	San Onofre North	\$8.83	3.8	40
3	San Clemente State	\$6.30	4.2	33
4	San Clemente City	\$6.09	3.0	36
5	Poche	\$5.69	2.0	1
6	Capistrano	\$5.49	1.4	17
7	Doheny	\$5.46	1.5	38
8	Salt Creek	\$5.43	4.1	70
9	Aliso Creek	\$4.98	3.8	17
10	Laguna	\$4.71	3.9	268
11	Crystal Cove	\$4.21	4.2	57
12	Corona Del Mar	\$4.07	4.0	116
13	Balboa	\$3.57	4.3	49
14	Newport	\$3.67	4.1	659
15	Santa Ana River	\$3.50	3.5	1
16	Huntington State	\$3.47	2.5	213
17	Huntington City	\$3.38	3.9	301
18	Bolsa Chica	\$3.26	4.0	206
19	Sunset	\$3.21	4.3	33
20	Surfside	\$3.21	4.2	2
21	Seal	\$3.18	3.3	240
22	Alamitos Bay	\$3.39	4.0	45
23	Belmont Shores	\$3.27	3.6	31
24	Long Beach	\$3.54	2.9	310
25	Cabrillo	\$4.04	3.0	52
26	Point Fermin	\$4.01	4.2	7
27	Royal Palms	\$4.04	4.1	13
28	Abalone Cove	\$4.35	4.2	3
29	Torrance	\$3.91	4.2	65
30	Redondo	\$3.84	3.6	191
31	Hermosa	\$3.74	4.1	249

²¹This cost is calculated as

$$Cost_{ij} = 2 * [one\ way\ travel\ dist * 0.145 + (one\ way\ travel\ time) * (0.5 * hourly\ wage)] \quad (26)$$

²²For a discussion on the percentage choice of wage rate in a travel cost model in a beach recreation application see Lew and Larson (2004).

Table 6: Water Quality Grade and Variance

	Grade				
	F	D	C	B	A
Occurrence	7	8	22	62	207
% Occurrence	3%	3%	7%	17%	58%
Trips	53	329	483	1,542	2,232
% Trips	1%	7%	10%	33%	48%

	Variance						
	0	0.25	0.5	0.75	1	1.25	1.5
Occurrence	34	6	2	0	2	0	4
% Occurrence	71%	13%	4%	0%	4%	0%	8%
Trips	3,883	369	5	0	289	0	96
% Trips	84%	8%	0%	0%	6%	0%	2%

Table 5: Beach Site Details (continued)

	Beach	Avg. Cost	Avg. Water Grade	Observed Trips
32	Manhattan	\$3.71	4.2	302
33	El Segundo	\$3.85	3.8	6
34	Dockweiler	\$3.78	3.7	16
35	Mother's	\$4.11	2.5	76
36	Venice	\$4.13	3.9	199
37	Santa Monica	\$5.06	3.3	400
38	Will Rogers	\$5.08	3.1	39
39	Topanga	\$5.10	3.0	4
40	Las Tunas	\$6.78	2.1	0
41	Malibu (Surfrider)	\$8.43	2.1	58
42	Dan Blocker (Corral)	\$9.81	4.0	10
43	Point Dume	\$9.35	3.2	22
44	Free Zuma	\$9.97	4.1	0
45	Zuma	\$10.01	4.2	79
46	El Matador	\$9.22	4.1	4
47	La Piedra	\$9.22	4.1	0
48	El Pescador	\$9.22	4.1	2
49	Nicholas Canyon	\$9.65	4.1	2
50	Leo Carrillo	\$9.62	4.1	20
51	County Line	\$10.03	4.0	6

Three water quality variables are constructed utilizing this data: yearly average grade, bimonthly average for all years, and the bimonthly worst grade reported during the survey year (Mohn et. al., 2003). See Table 5 for average yearly water quality grades. In addition to these three measures a set of discrete water quality variables, indicating an F or D grade, were constructed. Table 6 reports summary statistics on the bi-monthly occurrence of water quality grade ratings, the bimonthly within beach variance for water grades, and the number of trips taken by water quality grade and variance category.

To be included in the final data set a trip requires a valid destination and the respondent who took the trip must have supplied all of the demographic variables included in the model. This data source not only contains the necessary variables to implement the standard models, but also is rich enough in preference, choice set awareness, and past activity data to be able to implement the latent segmentation assignment of individuals.

Table 7: Composite Beach Variables and Their Components

Composite Variables		Component Variables
Developed Beach	3 or more	Street Access
Very Developed	8 or more	Public Transit
		Restaurants
		Stores
		Concessions
		Rentals
		Beach Clubs
		Houses
		Condos/Hotels
		Pier
		Concerts
		Volley Ball Tournaments
Wild Beach	1 or more	Pedestrian Access Only
		Rocky
		Tide pools
		Dogs Allowed
Ugly Beach	1 or more	Oilpumps
		Oilrigs
		PowerSewer
		Stormdrains

5 Application

5.1 Recreational Beach Choice Model

Following the literature, recreational site choice decision occasions are modeled using the discrete choice RUM as a function of site attributes, individual characteristics, and seasonal data holding the number of trips taken as exogenously determined. The CL, RPL, and FML variants of the RUM are estimated using an identical specification for the site choice probability. The FML model uses additional variables as arguments to the group membership function.

To capture the seasonal variation in preferences, a seasonal dummy is included into the segment membership function. Previous recreational modeling studies which have focused on trip temporal characteristics, such as season or part of the week, have operationalized the temporal data as an interaction variable or used it to segment the data set *a priori*. The use of the time variable in the FML enables the analyst to capture the probabilistic nature of seasonal influences on beach recreation in Southern California where there are often unseasonably warm and cold days during the winter and summer respectively.

Explanatory variables used in the RUM specifications can be categorized into beach choice and group membership variables. Modeling individual site choices for beach recreation requires explanatory variables in terms of how the beaches in the choice set differ from one another. Binary composite variables for development, very developed, wild, and ugly beaches serve to collapse twenty component attributes into four composite indicator variables (Table 7).²³

Beach choice variables incorporated into the CL, RPL, and FML model specifications include beach

²³The data set includes a large number of beach attribute variables (42) relative to the number of beaches in the choice set (51). Therefore, a composite choice variable strategy for the appropriate right hand side variables was developed in part to handle correlation within the beach attribute data set (Mohn et. al., 2003). The variables that are used to construct the composites are 0/1 indicator variables for the absence/presence of the relevant attributes. For a detailed discussion on the formation of the composite choice set, see Hanemann (2004).

Table 8: Choice Variable Summary Statistics

Choice Variables	Min	Mean	Max	Std Dev
Cost	3.183	5.546	10.027	2.375
Water Quality	1.373	3.602	4.333	0.757
Beach Length (ln)	-2.207	0.352	2.088	0.940
Developed	0.000	0.549	1.000	0.503
Very Developed	0.000	0.196	1.000	0.401
Wild	0.000	0.314	1.000	0.469
Ugly	0.000	0.275	1.000	0.451

Table 9: Correlation of Choice Variables

	Cost	Water Quality	Beach Length	Developed	Very Developed	Wild	Ugly
Cost	1	0.034	-0.347	-0.168	-0.215	-0.013	-0.329
Water	0.034	1	-0.099	-0.293	-0.009	0.094	0.038
Length	-0.347	-0.099	1	0.302	0.385	-0.137	0.108
Developed	-0.168	-0.293	0.302	1	0.448	-0.236	0.116
Very Dev	-0.215	-0.009	0.385	0.448	1	-0.227	-0.082
Wild	-0.013	0.094	-0.137	-0.236	-0.227	1	-0.132
Ugly	-0.329	0.038	0.108	0.116	-0.082	-0.132	1

travel cost, water quality, the length of the beach, and a set of binary composite variables for capturing the developed, very developed, wild, or ugly nature of the beaches. Beach attribute summary statistics and correlation matrices are displayed in Table 8 and Table 9 respectively.

Group membership dummy variables used in the FML specifications indicate whether the trip occurred during winter, the recreator got in the water, the recreator is male, kids are present on the trip, the recreator is a student, the recreator works full time, and the recreator is a college graduate (Table 10 and Table 11).

The model specification reported upon in this paper is a preliminary specification designed to illustrate the level of heterogeneity which characterizes preferences for attributes that describe beach recreation site choices. The objective of this paper is to illustrate the importance of handling systematic preference heterogeneity in a discrete choice setting characterized by diverse user groups. Estimation results indicate that the FML model is a useful tool in analyzing Southern California beach choice recreational decisions. The choice model specification reported in this paper focuses on broad composite beach attribute variables and excludes several activity specific variables. Inclusion of these omitted variables is expected to impact the parameter

Table 10: Membership Variable Summary Statistics

	Min	Mean	Max	Std Dev
Constant	0.0	1.000	1.0	0.000
Winter	0.0	0.213	1.0	0.409
Summer	0.0	0.377	1.0	0.485
In Water	0.0	0.266	1.0	0.442
Male	0.0	0.561	1.0	0.496
Kids	0.0	0.266	1.0	0.442
Student	0.0	0.175	1.0	0.380
Work Fulltime	0.0	0.649	1.0	0.477
College Grad	0.0	0.534	1.0	0.499

Table 11: Correlation of Membership Variables

	Winter	Summer	In Water	Male	Kids	Student	Work Fulltime	College Grad
Winter	1	-0.404	-0.149	0.061	-0.079	-0.032	0.068	0.089
Summer	-0.404	1	0.201	-0.056	0.098	-0.017	-0.064	-0.011
Water	-0.149	0.201	1	0.082	0.064	0.010	0.003	-0.010
Male	0.061	-0.056	0.082	1	-0.210	-0.075	0.238	0.037
Kids	-0.079	0.098	0.064	-0.210	1	-0.014	-0.058	-0.125
Student	-0.032	-0.017	0.010	-0.075	-0.014	1	-0.148	-0.116
Fulltime	0.068	-0.064	0.003	0.238	-0.058	-0.148	1	0.134
College	0.089	-0.011	-0.010	0.037	-0.125	-0.116	0.134	1

and welfare estimates reported in this paper. Additionally, inclusion is expected to strengthen the preference group separation of the FML model due to an increase in the dimensionality of preference space.²⁴

5.2 Estimation

The log likelihood functions for the three FML model specifications discussed above each have two major components: the segment membership probability, π_{is} , which is specified as a multinomial logit with individual attributes, Z_{it} , arguments; and the site choice probability, $\Pr_{ijt|s}$, which is specified as a conditional logit with site attribute, X_{ijt} , arguments.

Estimation of the preceding log Likelihood function using traditional derivative based maximum likelihood search algorithms can be troublesome. The non-linear nature of the likelihood function, and the exogenously determined number of segments, S , cause instability because the likelihood function is maximized on a ridge in parameter space if S is misspecified (Wedel, 1993). This is a common issue in the finite mixture model literature and a common solution is to implement the Expectation Maximization (EM) algorithm (see Ruud, 1991 for a thorough discussion of the algorithm and Arcidicon and Jones (2003) for a recent application to finite mixture models).

5.2.1 Observation Weighting

Due to the unbalanced panel nature of the data, observation weighting can affect the estimation results. A common approach in the literature is to weight each observation equally, however problems can arise due to the overweighting of the segments of the respondent population which have the most observations. An alternative approach would be to weight the observation of each individual by the inverse of the number of observations for that individual. Both of these approaches can be estimated and the results tested for robustness. Alternative weighting strategies can be researched in the choice avidity literature. This research will use the standard equal weighting approach.

5.3 Choice Model Estimation Results

Estimation of the CL, RPL, and FML models is implemented using numerical solutions with the GAUSS programming language and the Maxlik maximum likelihood software.²⁵ The CL and RPL model estimation

²⁴Whereas use of composite categorical data variables as a data reduction tool leads to a loss of information in the pattern of data over the attributes and respondents; as it is the pattern of data which allows the identification of latent segments (Ramaswamy, 1999).

²⁵Gauss code for the RPL is available on-line from Kenneth Train (2001).

is performed using the Newton-Raphson (NR) search algorithm and the FML is estimated using the Broyden-Fletcher-Goldfarb-Shanno method (BFGS) followed by the NR method. The model specification for beach choice variables is the base model specification from the preliminary report by the Southern California Beach Valuation Project (Hanemann et. al., 2004).²⁶ White's standard errors are calculated for all regressions to correct for violations of independence between observations from a respondent.

The CL model parameter estimates are of the expected and plausible sign, except for the 'ugly beach' dummy parameter estimate. Parameter estimates for travel cost, and very developed are negative. Parameter estimates for water quality rating, beach length, and developed beach dummy variables are positive. Counter intuitively the ugly beach dummy variable coefficient is positive.²⁷ The wild beach dummy coefficient is negative and not statistically different than zero. CL model parameter estimates are presented in Table 12.

The RPL model parameter estimates are of the same sign as those of the CL model. This result is expected. However the coefficient estimate for water quality is negative and not statistically significant, and the wild beach dummy's coefficient estimate is negative and statistically significant. RPL model parameter estimates are presented in Table 12. As expected, the RPL has greater explanatory power than the CL model indicated by high pseudo R^2 and other test statistics (Table 14).

5.3.1 Finite Mixture Logit Segment Testing and Results

Model estimation using the FML specification allows for an increased focus regarding the heterogeneous nature of the sample population's preferences. The FML is estimated iteratively with an increasing number of preference segment groups per specification. For specification of the FML model, a complete set of beach attribute coefficients is estimated for each latent segment. Additionally, a set of probabilities for each segment is estimated assigning segment membership as a function of the individual characteristics incorporated into the model.²⁸ The FML model is estimated for specifications with 2, 3, and 4 segments. Following the statistical segment testing methodology from the literature, the 4 segment model is chosen as having the greatest explanatory power. The 4 segment model (FML-4) has the highest R^2 compared to the CL, RPL, and 2 and 3 segment FML specifications. The 4 segment model also shows statistical significant improvements over the 3 segment model for several other test statistics: AIC, AIC-3, and BIC (Table 14). A 5 segment model is programmed in Gauss, but did not converge despite using a variety of parameter starting values and search techniques.²⁹ The lack of convergence with the 5 segment model signals that 5 segments is too many, as parameter estimates are known to tend towards negative and positive infinity when an $N + 1$ segment FM model is implemented on data which actually has N preference segments (Beard et. al., 1991).

The literature cautions against absolute reliance on statistical tests to determine the number of segments in a finite mixture and suggests the use of common sense (Beard et. al., 1991, and Boxall and Adamowicz, 2002). It is suggested that in most cases no more than 5 segments are needed in the FM framework (Heckman and Singer, 1984). The maximum number of feasible segments for a 7 dimensional preference space is 8 segments.

In terms of within sample forecast accuracy the 4 segment model outperforms the CL, and 2 and 3 segment models. Table 15 displays the percent of correct beach recreation site choice predictions for weighted segment membership and maximum probability single segment membership. In both cases the 4 segment

²⁶ Additional model specifications for the standard logit and nested logit are analyzed in the Southern California Beach Project reports.

²⁷ This is likely due to an omitted variable.

²⁸ The number of segments minus one set(s) of segment membership function coefficients are estimated in order to account for the indeterminacy in the model.

²⁹ FML model for 1 to 6 segments are programmed and estimated with simulated data consisting of 1 to 6 preference segments.

Table 12: Parameters on Choice and Membership Variables

Choice Variables	Logit	RPL		FML-4			
		Mean	SD	Seg 1	Seg 2	Seg 3	Seg 4
Cost	-0.085 (-50.887)	-0.182 (-34.016)	0.109 (23.921)	-0.653 (-6.074)	-0.021 (-11.919)	-0.408 (-15.980)	-0.366 (-10.415)
Water Quality	0.105 (4.316)	0.028 ^a (1.008)	-0.007 ^a (-0.055)	-7.950 (-4.395)	0.047 ^a (0.852)	10.382 (10.673)	-0.637 (-5.606)
Beach Length	0.470 (18.627)	0.567 (19.184)	-0.006 ^a (-0.114)	-0.871 (-2.320)	0.259 (5.166)	2.160 (9.73)	0.814 (7.508)
Developed	0.789 (17.456)	1.192 (5.770)	-1.885 (-4.317)	1.422 (3.541)	0.527 (5.200)	1.998 (11.693)	-0.448 (-2.226)
Very Devlp	-0.097 (-2.458)	-2.271 (-2.728)	9.546 (3.252)	8.857 (4.482)	0.637 (5.746)	-6.347 (-14.289)	1.836 (8.261)
Wild	-0.008 ^a (-0.192)	-0.662 (-4.040)	2.200 (7.537)	-2.291 (-3.995)	0.206 (2.271)	-6.253 (-7.288)	1.706 (8.754)
Ugly	0.073 (2.122)	0.100 (2.186)	-0.364 ^a (-0.889)	10.343 (4.156)	0.537 (6.220)	-8.461 (-13.663)	0.748 (6.369)
Segment Variables				Seg 1	Seg 2	Seg 3	Seg 4
Constant				-2.674 (-8.781)	-0.788 (-4.943)	-1.322 (-8.871)	0
Winter				0.299 ^a (1.392)	-0.204 ^a (-1.295)	0.704 (5.162)	0
Summer				1.195 (6.006)	0.282 (2.058)	0.516 (3.994)	0
In Water				-6.814 (-2.119)	0.294 (1.962)	0.028 ^a (0.205)	0
Male				1.907 (7.321)	-0.129 ^a (-0.929)	0.855 (6.334)	0
Kids				0.204 ^a (0.977)	0.446 (3.219)	0.120 ^a (0.888)	0
Student				0.083 ^a (0.287)	0.313 ^a (1.775)	-0.786 (-3.823)	0
Work Fulltime				-0.980 (-4.972)	0.480 (3.757)	-0.309 (-2.597)	0
College Grad				1.235 (5.764)	0.209 ^a (1.616)	1.006 (8.206)	0

^a Indicates that the parameter is not significantly different than 0 at the 5% level.
(T-statistics) are calculated using White's standard errors.

Table 13: FML-4 Membership Probabilities

Individual Seg Probabilities	Seg 1	Seg 2	Seg 3	Seg 4
Min	0.0%	5.7%	3.2%	10.2%
Mean	10.6%	29.8%	27.2%	32.4%
Max	64.2%	70.4%	69.8%	55.9%
Seg. Membership By Max Probability	6.4%	25.1%	33.8%	34.9%
Water Quality mWTP	-\$12.18	\$2.19	\$25.46	-\$1.74

Table 14: Model Selection Statistics

Estimation Results Conditional Logit, Random Parameters Logit, and Finite Mixture Models ^a					
Model	Logit	RPLb	FML ^c		
Segments	1		2	3	4
LL at Convergence	-14014.08	-13380.74	-12863.50	-12317.03	-12066.10
Convergence					
LL at 0	-18251.55	-18251.55	-18251.55	-18251.55	-18251.55
Parameters	7	14	23	39	45
AIC ^d	28042.16	26789.48	25773.01	24712.06	24222.20
AIC-3 ^e	42063.25	40184.23	38659.51	37068.08	36333.31
BIC ^f	14043.63	13439.84	12960.60	12481.66	12256.07
ρ^{2g}	0.232	0.267	0.295	0.325	0.339
mWTP			-\$6.87	-\$7.66	-\$12.18
			\$18.40	\$7.37	\$2.19
				\$21.03	\$25.46
					-\$1.74
Avg mWTP ^h	\$1.23	\$0.16	\$5.64	\$5.89	\$5.71

^aSample size is 4642 choices from 595 individuals (N).

^bRPL represents the random parameters logit model.

^cFML represents the finite mixed logit model.

^dAIC (Akaike Information Criterion) is calculated using $-2(LL-P)$.

^eAIC-3 (Akaike Information Criterion-3) is calculated using $-3(LL-P)$.

^fBIC (Bayesian Information Criterion) is calculated using $-LL+[(P/2)*\ln(N)]$.

^g ρ^2 is calculated as $1-(LL)/LL(0)$.

^hAverage Willingness to Pay is a weighted average of the willingness to pay by segment, using estimates of segment membership. Weighted WTP ranges from -\$4.06 to \$17.66.

Table 15: Within Sample Forecast Accuracy

Segments	Weighted Model % Correct	Point Estimate % Correct
1	15.2%	15.2%
2	20.6%	18.9%
3	21.7%	20.2%
4	23.6%	24.3%

model predicts a larger number of trips correctly. Taking all of the above factors into account, I conclude that a 4 segment FML model is the best model.

The ability to construct the distribution of welfare estimates for the sample population is one of the primary benefits of the FML model. In the beach choice application each trip occasion is characterized by a constant and 8 individual and trip specific binary variables. This simple characterization of each trip by agent and seasonal characteristics results in 256 different probability assignments which are used to assign beach choice preference group membership to each choice occasion triple.

The 4 segment FML model estimates the probability that an individual is a member of each preference group conditional on the season of trip and individual recreator characteristics. Each individual choice occasion in the sample thus has a probability of being in each segment.³⁰ For some choice occasions the probability is high (up to 70%), while for others it approaches zero (Table 12). Segments 4, 3, and 2 are the most likely preference groups to characterize the largest number of beach choice occasions at 34.8%, 33.8%,

³⁰Choice occasions are the individual recreator, water use, season triples that characterize each trip.

Table 16: Segment Membership Composition

Estimated Segment Composition by Membership Variable						
		Segment				Total
		1	2	3	4	
Winter Trip	0 1	521 3	1217 38	1055 578	862 368	3,655 987
Summer Trip	0 1	123 401	662 593	1109 524	999 231	2,893 1749
In Water	0 1	524 0	696 559	1067 566	1,122 108	3,409 1233
Male	0 1	287 237	1004 251	331 1,302	415 815	2,037 2605
Kids	0 1	437 87	799 456	1253 380	919 311	3,408 1234
Student	0 1	363 161	980 275	1629 4	857 373	3,829 813
Work Fulltime	0 1	485 39	117 1138	421 1,212	606 624	1,629 3013
College Graduate	0 1	232 292	781 474	33 1,600	1,118 112	2,164 2478
Total		524	1,255	1,633	1,230	4,642

and 25.1% of the total number of trip. Segment 1 is least likely preference group to characterize a choice occasion (6.4%) with the lowest mean percentage of group membership, 10.6%. However it has a 64.2% probability of characterizing some choice occasions.

Summary statistics for the composition of the estimated segment membership in terms of trip and individual characteristics are displayed in Table 16. Beach trips that are estimated to be characterized by segment 1 preferences are 77% likely to occur during the summer and 45% likely to be taken by male beach recreators. However membership in segment 1 is the lowest out of all groups. Estimated segment 2 preference type trips are 91% taken by beach recreators that are employed full time. Just under half of these trips are taken during the summer months and include water recreation. Segment 2 is characterized by summer trips, water use, kids on trips, female recreators, and full time employment. Segment 3 trips are characterized by male beach recreators that work full time. Winter trips, trips taken by male recreators, and those involving water recreation are most likely characterized by segment 3. Segment 4 trips are likely to occur during the shoulder season and have 66% male beach recreators. Trips taken by student recreators are likely to be characterized by segment 4 preferences.

Water Quality and Membership Consistency Three specifications of the water quality variable are investigated: average yearly grade, monthly grade, and a dirty water dummy variable. The results of all estimated models are qualitatively robust, however the continuous yearly grade water quality variable specification is chosen over the competing specifications based on improved measure of fit, improved coefficient robust standard errors, and ease of convergence. Table 17 reports the log likelihood score at convergence of

Table 17: LL scores for Yearly and Month Water Quality

	Logit	FML-2	FML-3	FML-4	RPL
Yearly	-14014	-12864	-12317	-12066	-13381
Monthly	-14017	-13209	-12915	-12356	-13392

Table 18: Segment Membership Time Consistency

	Number of Different Segments Per Individual				
	1	2	3	4	Total
Individuals	427	140	28	0	595
Trips Taken	2650	1394	598	0	4642

the CL, RPL, and FML models for the yearly and monthly grade specification. While it is noted that the competing specifications are not nested, the log likelihood scores indicate that the yearly water quality grade variable provides an improved fit. This result indicates that beach recreators may base their recreational decisions based on impressions about water quality that are formed over many years as opposed to current information. Hanemann et. al. report a similar finding regarding GEV beach choice model estimation (2004).

Estimation results indicate that segment membership consistency is characterized by variation over time preferences. 28% of individuals accounting for 43% of the trips took trips that are characterized by more than one preference segment (Table 18). Individuals that are characterized by one segment type take an average of 6.2 trips, those that are characterized by two or three segment types take an average of 10 or 21.4 trips respectively. No beach recreator in the sample took trips characterized by all four segment types. Additionally, statistically significant parameter estimates on time varying attributes in the segment membership function indicate membership variation over time.

5.4 Welfare Estimates for Water Quality Changes

5.4.1 Marginal Value of Estimates

The average beach recreator in the sample has an estimated marginal value or willingness to pay (mWTP) of \$5.71 for a water pollution rating increase of one letter grade when estimated using the 4 segment FML specification. This FML estimate is 4.64 times greater relative to the CL specification estimate of \$1.23 (the mWTP measure for the RPL is \$0.16³¹). However, this valuation estimate ranges from negative to \$17.66 for individual beach recreators (roughly 14.35 times the CL mWTP measure).³² See Table 12.

Latent groups 3 and 1, respectively, have the highest and lowest mean mWTP estimates for a one letter grade increase in water quality. With a mean mWTP point estimate of roughly 20 times the CL mWTP estimate, Group 3 membership is particularly likely for winter trips taken by male college graduates that work

³¹Note the parameter estimate on water quality is not statistically different than zero for the RPL model.

³²Theoretically I expect that WTP is greater or equal to 0. However, a non-negativity constraint is not imposed during the process of estimation. In the case of RPL, although the RPL mWTP is positive, a portion of the distribution of the mWTP takes on negative values. In the case of FML, I believe the negative estimates of mWTP for Group 1 and 4 are likely due to an omitted variables bias, because the model fitted here does not include certain activity-specific beach characteristic variables that are expected to impact the parameter estimates.

Figure 2: Marginal Value of Water Quality

full time and do not have children accompanying them to the beach. Individuals with Group 3 preferences are likely to choose beaches that have long beach length, development, but are not very developed, wild, or ugly (Table 12 and Table 14).

On the bottom half of the mWTP distribution, Group 1 has a mean mWTP point estimate of roughly negative 10 times the CL mWTP point estimate. Trips that occur during the winter, where the respondent went into the water by recreators that work full time are less likely to be characterized by Group 1 preferences. Additionally, recreators that are male, students, do not work full time, and are not college graduates are more likely to be characterized by Group 1 than Group 3. Those with Group 1 preferences are likely to choose beaches that are very developed, ugly, and have poor water quality.

The existence of multiple preference groups allows the construction of a multi-modal welfare distribution. A major strength of the FML approach is that the location within the distribution of specific welfare measures is recoverable conditional on individual and trip specific characteristics. The mWTP distribution for an improvement in water quality of one letter grade illustrates the heterogeneity in preferences for coastal water quality (Figure 2). Trips that occur during the winter, involve getting in the water, and are taken by male college graduates are associated with the representative groups that have a high valuation for an improvement in water quality. Conversely, trips taken during by students are strongly associated with representative groups with low mWTP for water quality.

Table 19: Estimated mWTP Regression

Regressors from Group Membership Function						
	OLS		GLS		GLS panel	
Intercept	2.451 (0.058)	2.366 (0.058)	2.680 (0.041)	2.691 (0.039)	2.294 (0.149)	2.234 (0.148)
Winter	3.803 (0.058)	3.780 (0.057)	3.640 (0.400)	3.746 (0.035)	4.120 (0.213)	4.043 (0.211)
Trip					0.247	0.228
Summer	0.138 (0.049)	0.129 (0.048)	0.464 (0.040)	0.324 (0.038)		
Trip					(0.182)	(0.180)
Water	3.626 (0.050)	4.016 (0.058)	2.241 (0.049)	3.023 (0.063)	3.941 (0.141)	4.273 (0.160)
Male	1.085 (0.046)	1.057 (0.045)	1.478 (0.032)	1.365 (0.031)	1.050 (0.100)	1.028 (0.098)
Kids	-0.145 (0.050)	0.237 (0.058)	0.033 (0.038)	0.023 (0.029)	-0.165 (0.110)	0.218 (0.142)
Student	-3.656 (0.057)	-3.660 (0.056)	-3.355 (0.045)	-3.516 (0.045)	-3.654 (0.125)	-3.657 (0.123)
Work	-0.272 (0.047)	-0.260 (0.046)	-0.369 (0.039)	-0.263 (0.036)	-0.273 (0.102)	-0.261 (0.101)
Fulltime						
College	3.154 (0.044)	3.165 (0.043)	2.769 (0.033)	2.671 (0.031)	3.134 (0.096)	3.149 (0.095)
Graduate						
Kids		-1.318 (0.105)		-1.119 (0.088)		-1.312 (0.312)
Water						
Regression Statistics						
R Sqr	0.828	0.833	0.855	0.874		
R Sqr-all					0.826	0.832
Adj R Sqr	0.827	0.833	0.855	0.873		
Obs	4642	4642	4177	4264	4642	4642

All coefficient estimates are significant at the 1% level.

Standard Errors are in parenthesis.

Bold indicates significantly different from 0, at the 1% level.

5.5 Second Stage Estimated Marginal Value Regression

To analyze the relationship between the estimated mWTP for individual trip occasions and the group membership variables. The weighted estimated mWTP for each beach trip are regressed on individual and seasonal characteristics of the trip with ordinary least squares (OLS), and both cross-section and panel specifications of generalized least squares (GLS)(Table 19). Coefficient estimates for the winter trip, in the water, and college graduate variables are positive for all three estimators. The coefficient estimate for the student variable is negative for all three estimators.

The coefficient estimate on the children present on trip variable is of particular interest. The OLS coefficient estimate for this variable is negative, whereas both GLS models produce coefficient estimates that are not significantly different than zero. However, the introduction of an interaction term for trips characterized by both the presence of children and getting into water produces negative and significant coefficient estimates for all three estimators. One would expect that beach trips that are taken with children and involve water recreation would have a higher probability of occurring at beaches with higher levels of water quality and would be associated with higher mWTP for water quality. One explanation for this result may be that the polluted beaches are characterized by features that are perceived by parents to provide safer environments for their children to swim, such as a lack of surf, but at the same time perpetuate water pollution. This result may be an example of cognitive dissonance and suggests the need for further research.

As illustrated by the paradoxical above result above the ability to construct the distributions of the

relative importance which site attributes have on site choice is an important tool for resource and health officials charged with the management of resources used by diverse user groups.

5.6 Compensating Variation Simulation Estimates

The compensating variation (CV) measure can be used to estimate the welfare change or consumer surplus (CS) resulting in from a change in the composition of site quality attributes. The CV measure captures the substitution effects due to a change in the choice set; where as the marginal value measure (mWTP) illustrates welfare changes for a marginal change in one attribute holding all others constant. Consumer surplus measures are calculated for four hypothetical attribute scenarios to illustrate the difference in consumer surplus measures calculated based on logit and FML choice model estimates. The four scenarios are: A) the closure of Santa Monica and Venice beaches to all beach use; B) the closure of 13 popular beaches;³³ C) degrading the water quality at all beaches to a D score; and D) dropping the water quality at Newport, Bolsa Chica, and Manhattan beaches by one letter grade to roughly a B score.

The estimated change in consumer surplus for each of the four scenarios is displayed in Table 20 for the logit and FML models. The simulated closure of beaches result in an estimated loss in consumer surplus for all beach recreators, regardless of model choice or segment membership. As expected the simulated closure of additional beaches result in a greater welfare loss.

The change in consumer surplus for degradation in water quality is negative on average for both scenarios C and D. However CS estimates for segments 1 and 4 for the 4 segment model are positive for both scenarios. The large CS gain for segments 1 and 4 are mathematically expected in scenario C, "D" grade water quality at all beaches. The cause of the negative welfare measure for segments 1 and 4 is likely due to omitted variable bias and not consumer preferences for poor water quality. In terms of water based recreation, segment 1 and 4 account for 0% and 9% respectively of trips. It follows that beach recreators characterized by segment 1 and 4 preferences will be less adversely impacted by a degradation in water quality.

Scenario D narrowly focuses and a degradation of one water grade, from roughly A to B, for three popular swimming beaches. Preference segments that are characterized by engaging in water based recreation have proportionally greater welfare changes than the preference segments that do not engage in water based recreation.

For the two beach closure scenarios the two competing models provide CV welfare measures of -\$0.95 and -\$7.91 (CL model) versus -\$1.16 and -\$11.96 (FML-4 model) for the simulated closures of 2 and 13 popular beaches respectively. For the two water quality degradation scenarios the two competing models provide CV welfare measures of -\$3.27 and -\$0.17 (CL model) versus -\$17.41 and -\$1.31 (FML-4 model) for the degradation of water quality to a 'D' grade for all area beaches and the dropping of one water quality grade for 3 popular swimming beaches respectively.

While the magnitude of the welfare loss generally increases with the number of segment groups estimated in the model, it is noted that this does not always hold. For instance, for scenario A, the estimated welfare loss calculated with the logit model estimates is greater than that estimated using the FML-2 model, and the welfare loss for the FML-3 and FML-4 are the same. Likewise, for scenario D, the ranking of the models with the largest estimated welfare loss is {2, 3, 4, 1} segments, with the 2 segment model resulting in the largest estimated change in consumer surplus. Interestingly, the 'unordered' welfare estimates are observed in the two scenarios that model a small change in site attributes and not the two scenarios with greater

³³These are: Laguna, Corona Del Mar, Newport, Huntington State, Huntington City, Bolsa Chica, Seal, Long Beach, Redondo, Hermosa, Manhattan, Venice, and Santa Monica Beach.

Table 20: Welfare Senarios: Beach Closure and Water Degradation

A: Close Santa Monica and Venice Beaches					B: Close 13 Popular Beaches			
	Logit	FML-2	FML-3	FML-4	Logit	FML-2	FML-3	FML-4
Min	-\$9.86	-\$11.67	-\$13.60	-\$13.60	-\$18.21	-\$23.16	-\$24.10	-\$27.79
Mean	-\$0.95	-\$0.70	-\$1.16	-\$1.16	-\$7.91	-\$9.40	-\$10.58	-\$11.96
Max	\$0.00	\$0.00	-\$0.01	-\$0.01	-\$0.40	-\$0.44	-\$2.40	-\$3.03
	Segment 1 WTP					Segment 1 WTP		
Min		-\$8.08	-\$6.26	-\$1.60		-\$26.09	-\$24.69	-\$24.05
Mean		-\$0.46	-\$0.28	-\$0.03		-\$5.42	-\$4.25	-\$4.14
Max		\$0.00	\$0.00	\$0.00		\$0.00	\$0.00	\$0.00
	Segment 2 WTP					Segment 2 WTP		
Min		-\$14.02	-\$20.31	-\$20.15		-\$29.10	-\$35.38	-\$42.42
Mean		-\$0.95	-\$2.81	-\$3.54		-\$13.27	-\$21.45	-\$29.04
Max		\$0.00	-\$0.02	-\$0.08		-\$1.21	-\$10.78	-\$17.86
	Segment 3 WTP					Segment 3 WTP		
Min			-\$21.72	-\$23.11			-\$25.49	-\$27.64
Mean			-\$0.43	-\$0.39			-\$6.27	-\$6.85
Max			\$0.00	\$0.00			\$0.00	\$0.00
	Segment 4 WTP					Segment 4 WTP		
Min				-\$11.50				-\$18.38
Mean				-\$0.48				-\$3.15
Max				\$0.00				\$0.00
C: Water Quality is 'D' at All Beaches					D: 3 Swimming Beaches Fall 1 Grade			
	Logit	FML-2	FML-3	FML-4	Logit	FML-2	FML-3	FML-4
Min	-\$3.79	-\$45.85	-\$42.95	-\$51.51	-\$0.60	-\$9.15	-\$6.49	-\$6.65
Mean	-\$3.27	-\$15.84	-\$17.63	-\$17.41	-\$0.17	-\$2.15	-\$1.40	-\$1.31
Max	-\$1.57	\$7.20	\$2.62	\$8.95	\$0.00	\$0.95	\$1.63	\$5.07
	Segment 1 WTP					Segment 1 WTP		
Min		\$5.97	\$4.71	\$7.74		\$0.00	\$0.00	\$0.00
Mean		\$18.02	\$19.96	\$30.79		\$0.36	\$0.43	\$0.86
Max		\$21.41	\$24.20	\$38.83		\$5.68	\$7.31	\$12.08
	Segment 2 WTP					Segment 2 WTP		
Min		-\$56.70	-\$20.98	-\$6.08		-\$11.86	-\$1.22	-\$0.35
Mean		-\$50.20	-\$19.62	-\$5.73		-\$4.57	-\$0.55	-\$0.16
Max		-\$31.47	-\$16.46	-\$4.99		-\$0.27	-\$0.19	-\$0.07
	Segment 3 WTP					Segment 3 WTP		
Min			-\$66.76	-\$80.86			-\$12.60	-\$13.26
Mean			-\$61.73	-\$75.28			-\$4.60	-\$5.14
Max			-\$41.51	-\$50.27			\$0.00	\$0.00
	Segment 4 WTP					Segment 4 WTP		
Min				\$1.24				\$0.00
Mean				\$4.68				\$0.11
Max				\$5.57				\$1.61

attribute changes.

While the FML model has much strength, care must be taken to properly specify the utility model to avoid single preference segment estimates with omitted variable bias that can lead to biased welfare measures.

6 Conclusion

Coastal water quality impacts recreation and tourism. Southern California beach recreators cite pollution as a primary reason for abstaining from swimming, a belief supported by studies linking swimming in polluted water with illness. While there is interest in understanding the impact of water quality on beach recreation to improve resource and public health management, this task is complicated by the diversity of user preferences and the multiuse nature of the beach.

This paper implements the FML RUM to highlight the importance of capturing preference heterogeneity. Exploiting an extensive beach recreational panel data set, this paper furthers the literature by applying the FML approach to model preference heterogeneity regarding the impact of an environmental variable related to health and seasonality on recreational choice. The application also increases the number of choice alternatives and the number of variables included in the segment membership function estimated with the FML model in the literature.

Application of the FML model to the Southern California beach recreation data set recovers 4 preference groups, highlighting the variation in the importance of water pollution on beach choice for a diverse sample of beach users. For these groups, the impact that water quality has on recreational site choice, as measured by the mean mWTP, ranges from negative to \$17.66, with an average of \$5.71, for an improvement in water quality of one letter grade. The mWTP estimate calculated with the CL model is \$1.23. The RPL coefficient estimate on water quality is not significantly different from 0, and yields a mWTP estimate of \$0.16. Compensating variation measures for consumer surplus associated with changes in beach attributes tell a similar story.

The FML approach facilitates the estimation of the distribution of water quality preferences and welfare measures across a diverse user-base, and enables researchers to identify user preference groups characterized by several variables. This increases the ability of resource managers to forecast the impact that changes in site characteristics will have on the beach choice and welfare across segments of society.

Estimation results indicate that recreators who enter the water have a higher estimated mWTP for water quality. Gender, employment, education, and seasonal variables are also important in estimating one's preferences. One troubling result of the model is the finding that the presence of children on beach trips which include water activity is not associated with a higher mWTP for improvements in water quality. This result highlights the model's ability to identify groups that resource managers and public health officials may desire to concentrate their educational outreach efforts.

The FML approach is likely to become increasingly important as diversity continues to grow, and the identification of user groups by a small number of variables becomes less feasible. The application of the model to a unique beach recreation data set is of major significance from the environmental management perspective. The powerful combination of being able to specify a model which simultaneously estimates the marginal benefits associated with different attributes for different groups and assigns group membership makes FML a particularly attractive model for policy analysis.

References

- [1] Adamowicz, W.L., J. Swait, P.C. Boxall, J. Louviere, and M. Williams. 1997. Perceptions versus objective measures of environmental quality in combined revealed and stated preference models of environmental valuation. *Journal of Environmental Economics and Management* 32:65-84.
- [2] Arcidiacono, P. and J.B. Jones. 2003. Finite Mixture Distributions, Sequential Likelihood and the EM Algorithm. *Econometrica*. Vol 71, no. 3, pp 933-946.
- [3] Beard, T.R., Caudill, S.B., Gropper, D.M. 1991. Finite Mixture Estimation of Multiproduct Cost Functions. *Review of Economics and Statistics*, November, v. 73, iss. 4, pp. 654-64.
- [4] Ben-Akiva, M. and J. Swait. 1986. The Akaike Likelihood Ratio Index. *Transportation Science*. 20: 133-136.
- [5] Berry, S., Levinsohn, J. and Pakes A. 1995. Automobile Prices in Market Equilibrium. *Econometrica*, Vol. 63, No. 4, pp. 841-890
- [6] Bockstael, N.E., W.M. Hanemann, and I.E. Strand, Jr. 1986. Measuring the benefits of water quality improvements in a recreational demand framework. *Water Resources Research*. Vol. 23, no. 5, pp. 951-960
- [7] Boxall, P.C. and W.L. Adamowicz. 2002. Understanding heterogeneous preferences in random utility models: A latent class approach. *Environmental and Resource Economics*. 23: 421-446.
- [8] Chamberlain, G. 1980. Analysis of Covariance with Qualitative Data. Chamberlain, Gary; *Review of Economic Studies*, January, v. 47, iss. 1, pp. 225-38
- [9] Chapman, D.J. and W. Michael Hanemann. "Environmental Damages In Court: The American Trader Case," In Anthony Heyes' (ed) *The Law and Economics of the Environment*, pp. 319-367, 2001.
- [10] Gordon, Peter, John Kuprenas, Jiin-Jen Lee, James E. Moore, Harry W. Richardson, Christopher Williamson. An Economic Impact Evaluation of Proposed Storm Water Treatment for Los Angeles County. Los Angeles: School of Engineering and School of Policy, Planning, and Development, University of Southern California. November 2002.
- [11] Greene, W, and D. Hensher. 2003. A Latent Class Model for Discrete Choice Analysis: Contrasts with Mixed Logit. *Transportation Research Part B*. Vol 37. pp. 681-698.
- [12] Haab, T,C,, and K.E. McConnell. 2003. *Valuing Environmental and Natural Resources: The Econometrics of Nonmarket Valuation*. Edward Elgar Publishing, Massachusetts.
- [13] Haaijer, R., and M. Wedel. 2000. Habit Persistence in Time Series Models of Discrete Choice. *Marketing Letters*. Vol. 12, pp 25-35.
- [14] Haile RW, Alamillo J, Barrett K, et al. An Epidemiological Study of Possible Adverse Health Effects of Swimming in Santa Monica Bay: Final Report. Santa Monica, Calif: Santa Monica Bay Restoration Project; 1996.
- [15] Hanemann, W.M. 1978. A Theoretical and Empirical Study of the Recreation Benefits from Improving Water Quality in the Boston Area. Dissertation, Harvard University, 1978, 532p.

- [16] Hanemann, W.M. 1982. Applied welfare analysis with qualitative response models, Working Paper. No. 241. University of California, Berkeley, 26 pp.
- [17] Hanemann, W.M., J.R. Hilger, C. Mohn, and L. Pendleton. November, 2003. Data Reduction for Los Angeles Area Beach Site Attributes. Southern California Beach Project.
- [18] Hanemann, W.M., L. Pendleton, C. Mohn, J. Hilger, K. Kurisawa, D. Layton, C. Busch, and F. Vasquez. August, 2004. Using revealed preference models to estimate the effect of coastal water quality on beach choice in Southern California. Southern California Beach Project.
- [19] Hanemann, W.M. Personal communication. November, 2005. Berkeley, California.
- [20] Hausman J., D. McFadden. 1984. Specification Tests for the Multinomial Logit Model. *Econometrica*. Vol. 52. pp. 1219-1240.
- [21] Heal the Bay. 2005. Beach Report Card: Grading System. www.healththebay.org/brc/gradingsystem.asp
- [22] Heckman, J., and B. Singer. 1984. A Method for Minimizing the Impact of Distributional Assumptions in Econometric Models for Duration Data. *Econometrica*, Vol. 52, No. 2., pp 271-320.
- [23] Jedidi, K., H.S. Jagpal, and W.S. DeSarbo. 1997. Finite_mixture Structural Equation Models for Response-Based Segmentation and Unobserved Heterogeneity. *Marketing Science*. Vol. 16, pp 39-59.
- [24] Jones, J.M., and Landwehr, J. T. 1988. Removing Heterogeneity Bias from Logit Model Estimation. *Marketing Science*, Winter, v. 7, iss. 1, pp. 41-59.
- [25] Kamakura, W., and G. Russell. 1989. A probabilistic Choice Model for Market Segmentation and Elasticity Structure. *Journal of Marketing Research*. Vol. 26, pp 379-390.
- [26] Lew, D.K., Larson, D.M. 2005. Valuing Recreation and Amenities at San Diego County Beaches. *Coastal Management*, 33:71-86.
- [27] McFadden, D. 1973. Conditional Logit Analysis of Qualitative Choice Behavior. In P. Zarembka, ed., *Frontier in Econometrics*. New York: Academic Press.
- [28] McFadden, D. 1978. Modeling the Choice of Residential Location, In *Spatial Interaction Theory and Residential Location*, A. Karlquist et al., editors, Amsterdam:North Holland, 75-96.
- [29] McFadden, D. 1986. The choice theory approach to market research. *Marketing Science*. Vol. 5, pp. 275-297.
- [30] Mohn, C., M.H. Hanemann, L. Pendleton, and D. Layton. 2003. Shifting Sands: A Preliminary Exploration of Seasonal Variation in Preferences for Beaches. Working Paper.
- [31] Morey, E.R. 2003. Latent Segmentation Likelihood Functions.' Personal Communication.
- [32] Nevo, A. 2000 A Practitioner's Guide to Estimation of Random-Coefficients Logit Models of Demand. *Journal of Economics and Management Strategy*, Winter, 513-48.
- [33] Natural Resources Defense Council. 2002. Testing the waters: A guide to water quality at vacation beaches, 2002. Volume 12, pp 24-57.

- [34] Natural Resources Defense Council. 2004. Testing the Waters: A Guide to Water Quality at Vacation Beaches.
- [35] Natural Resources Defense Council. 2005. Testing the Waters: A Guide to Water Quality at Vacation Beaches.
- [36] Parsons, G.R. 1991. A Note on Choice of Residential Location in Travel Cost Demand Models. *Land Economics*, August, v. 67, iss. 3, pp. 360-64.
- [37] Pendleton, L. 2001. Managing Beach Amenities to Reduce Exposure to Coastal Hazards: Storm Water Pollution. *Coastal Management*, 29:239-252.
- [38] Provencher, B., K.A. Baerenklau, and R.C. Bishop. 2002. A Finite Mixture Logit Model of Recreational Angling with Serially Correlated Random Utility. *American Journal of Agricultural Economics*. Vol 84, pp 1066-1075.
- [39] Rabinovici, S.J.M., Bernknopf, R.L., Wein, A.M., Coursey, D.L., and Whitman, R.L. 2004. Economic and Health Risk Trade-offs of Swim Closures at a Lake Michigan Beach. *Environmental Science and Technology*. Vol. 38, No. 10, pp 2737-2745.
- [40] Ramaswamy, V., R. Chatterjee, and S. Cohen. 1999. Reply to "A Note on Ramaswamy, Chatterjee, and Cohen's Latent Joint Segmentation Models." *Journal of Marketing Research*. Vol 36, pp 115-119.
- [41] Ruud, P.A. 1991. Extension of estimation methods using the EM algorithm. *Journal of Econometrics*. Vol 49, pp 305-341.
- [42] Shonkwiler, J.S. and W.D. Shaw. 2003 "A Finite Mixture Approach to Analyzing Income Effects in Random Utility Models" in *The New Economics of Outdoor Recreation*, N. D. Hanley, ed. Edward Elgar, pp. 268-278.
- [43] Small, K.A. and H.S. Rosen. 1981. Applied Welfare Economics with Discrete Choice Models. *Econometrica*, Vol. 49, No. 1. pp. 105-130.
- [44] Sorensen, B. 1975. Energy and resources. *Science*. Vol. 189 : 255-60
- [45] Swait, J. and J. Louviere. 1993. The role of the scale parameter in the estimation and use of multinormal logit models. *Journal of Marketing Research*. Vol. 30. pp 305-314.
- [46] Swait, J.R. (1994) A structural equation model of latent segmentation and product choice for cross-sectional revealed preference choice data. *Journal of Retailing and Consumer Services* Vol. 1, no 2, pp. 77-89.
- [47] Train, K. (1997) Recreation Demand Models with Taste Differences Over People. *Land Economics*. Vol. 74, No. 2, May 1998.
- [48] Train, K. 2001. Discrete Choice with Simulation Gauss Code..[www.elsa.berkeley.edu/ Software/ abstracts/ train0401.html](http://www.elsa.berkeley.edu/Software/abstracts/train0401.html)
- [49] Train, K. 2003. *Discrete Choice Methods with Simulation*, Cambridge. Cambridge University Press.
- [50] United States Forest Service. 2000. National Survey on Recreation and the Environment: American's Participation in Outdoor Recreation. Southern Research Station.

- [51] Walsh, R. G.; Johnson, D. M.; McKean, J. R. 1992. Benefit Transfer of Outdoor Recreation Demand Studies, 1968-1988; *Water Resources Research*, March, v. 28, iss. 3, pp. 707-13.
- [52] Wedel, M., W.S. DeSarbo, J.R. Bult, and V. Ramaswamy. 1993. A Latent Class Poisson Regression Model for Heterogeneous Count Data. *Journal of Applied Econometrics*. Vol. 8, No. 4, pp 397-411.
- [53] Wooldridge, J.M. 2002. *Econometric Analysis of Cross Section and Panel Data*. MIT Press, Massachusetts.