



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.



Paper prepared for the 122nd EAAE Seminar
**"EVIDENCE-BASED AGRICULTURAL AND RURAL POLICY MAKING:
METHODOLOGICAL AND EMPIRICAL CHALLENGES OF POLICY
EVALUATION"**

Ancona, February 17-18, 2011



**Counterfactual approach for assessing agri-environmental
policy: The case of the Finnish water protection policy**

Lankoski J.¹ and Ollikainen M.¹

¹ Department of Economics and Management, University of Helsinki, Helsinki, Finland

jussi.lankoski@helsinki.fi

Counterfactual approach for assessing agri-environmental policy: The case of the Finnish water protection policy

Lankoski J. and Ollikainen M.

Abstract

This paper applies counterfactual approach to assess the impacts of agri-environmental programs. Counterfactual analysis evaluates policies answering questions: what would have happened if...? We develop a theoretical framework for counterfactual analysis based on the inter-linkages between the behaviour of agents and the response of environmental systems to the economic decisions. We apply our model to assess the performance of the Finnish Agri-Environmental Programme to reduce agricultural nutrient runoff to the Baltic Sea. Counterfactual analysis allows us to determine both the unit effectiveness of the measures included in the Programme and its preventive impact. We demonstrate that the Finnish Agri-Environmental Programme does not achieve its goals, because it fails to anticipate farmers' responses to incentives created by the Common Agricultural Policy and the Agri-Environmental Programme itself. The social cost-benefit analysis of the Program shows negative net benefits: benefits from reduced nutrient loading are much lower than support payments.

Keywords: environmental policy evaluation, counterfactual analysis, nutrient runoff, the Baltic Sea

JEL classification: Q5, H23, H43.

1. INTRODUCTION

Common agricultural policy (CAP) of the EU includes a possibility for voluntary environmental protection programs in agriculture. The programs provide an incentive payment for compensating the compliance costs and farmers' private transaction costs. In many countries these voluntary programs entail significant monetary transfers from tax payers to farmers. Thus, it is important to assess the success of these voluntary agri-environmental programs. As is well-known, it is a challenging task to assess any program, save a case where environmental effects are deeply involved. One method for assessing environmental policies is to contrast them with alternative, hypothetical policies or states of affairs and thereby reveal the impacts of policies, that is, to resort counterfactual analysis. Counterfactual analysis belongs to the basic tools for policy evaluation in economics. Counterfactual analysis answers the question: what would have happened if...? Counterfactual analysis builds on a non-observable case, called counterfactual, against which policies can be evaluated. A comparison of the counterfactual with the actual case sheds light to the critical factors explaining the impacts of policy.

In this paper we suggest a formal approach to counterfactual analysis based on behavioural assumptions, that is, the profit maximization hypothesis of farmers. The farmers maximize profits both in the presence and absence of policy. The choices over the use of inputs and land allocation between crops can then be linked to nutrient runoff and other relevant

environmental process functions to predict the environmental outcomes of different policies that can then be contrasted to the goals and measured outcomes of agri-environmental policies.

We develop a theoretical frame and derive the counterfactuals for empirical analysis from it. We apply our theoretical frame to agricultural water protection policy which aims at reducing nutrient runoff from arable lands to the Baltic Sea. In Finland, like in all Baltic Sea countries, agriculture is the main source of nitrogen and phosphorus loads (60% of the anthropogenic phosphorus and 52% of nitrogen loads) to the Sea (HELCOM 2010). Given that nutrient loads from point sources have been reduced considerably, pressure to reduce loads from agriculture is high and agri-environmental program tries to cope with this challenge. The first Agri-Environmental Programme was for 1995-1999. Although the program addressed many environmental issues (water quality, air quality, biodiversity, and landscape), improving surface water quality has been and still is the highest priority – also in the subsequent programmes (2000-2006, 2007-2013) that have only slightly fine-tuned the original programme.

Finnish authorities expected that agri-environmental program could achieve about 30 – 40% decrease of phosphorus and nitrogen runoff by 2010 (MMM 2003). Even more ambitious 50% nutrient runoff reduction targets by 2005 were set for agriculture in the Government resolution as regards surface water protection targets (YM 2000). Despite the ambitious goals and all efforts and support payments, something has gone wrong, however. No actual reductions in measured/monitored nutrient runoff can be observed and especially nitrogen runoff is reported to have increased. Clearly, a theory-guided analysis of the water protection policy in agriculture is needed to evaluate the impacts of the water protection policy and to find an explanation for the observed no-progress in actual nutrient loads. To examine the performance of agri-environmental policy, we develop theoretically and employ empirically two basic counterfactuals.

The rest of the paper is structured as follows. Section 2 provides our theoretical framework and section 3 presents an empirical framework. Results are presented in section 4 and concluding section 5 ends the paper.

2. THE FRAMEWORK: A COUNTERFACTUAL APPROACH

Consider an introduction of a new agri-environmental policy, A , starting a period $t = T$. This ends the previous policy regime, which can be no policy (free market solution), or some other policy denoted by B . Now, let x_0 , x_A and x_B denote the vector of instruments of no policy intervention and policies A and B , respectively. For no policy intervention, this instrument set is naturally equal to zero, $x_0 = \{0\}$. To keep the presentation simple, policy instruments under policy A are a fertilizer application constraint (\bar{l}^A), a buffer strip requirement (\bar{m}), the CAP area payment (a) and the environmental support payment per hectare (b), that is, $x^A = \{\bar{l}^A, \bar{m}, a, b\}$. Furthermore, the previous policy regime B is assumed to consist of direct price support (s) and a requirement for large set-aside areas (E), that is, $x^B = \{s, E\}$.

The farmers optimize their agricultural profits subject to exogenous variables and the policy instruments under each policy regime. Denote the conventional response function of crop i as $y = f_i(l_i)$ with $f'_i(l_i) > 0$ but $f''_i(l_i) < 0$. Let p_i be the price of the crop and c the price of fertilizer input, and L_i be the amount of land allocated to each crop under the three policies. Market parameters alone or together with the instruments under policy regimes A and B determine the optimal use of inputs and land allocation (including entry and exit of land in agricultural sector). We next develop the farmers choices under each policy regime.

Under no policy intervention (denoted by superscript 0), the profits of a given parcel allocated to crop i are given by $\pi_i^0 = p_i f_i(l_i) - c$, and profits from the land area allocated to crop i by $r_i = \pi_i^0 L_i^0$. The optimal solution entails $l_i^0(p_i, c)$ and $L_i^0(p_i, c, p_j)$, where p_j refers to prices of the other crops and $\sum_{i=1}^n L_i^0(p_i, p_j, c) = L^0$, where L^0 refers to overall land in cultivation in the no-policy regime.

Under the new policy regime A the per parcel profits are given by $\pi_i^A = (1 - \bar{m})[p_i f_i(l_i) - cl_i] + a + b$ subject to $l_i^A \leq \bar{l}_i$. At the optimal solution output price and fertilizer cost no longer impact fertilizer intensity, because the fertilizer application constraint is binding ($l_i^A = \bar{l}_i$); also the buffer strip is mandated $m^A = \bar{m}^A$. Using these two mandatory figures, the overall amount of fertilizer applied to each hectare is $l_i^A = (1 - \bar{m})\bar{l}_i$. Land allocation, however, continues to depend on the relative profitability of each crop, and thereby it depends on prices, so that $L_i^A(p_i, c, p_j, a, b)$, where a refers to CAP compensation payment and b refers to area-based environmental support payment. The overall profits are then given by $r_i^A = \pi_i^A L_i^A$ with $\sum_{i=1}^n L_i^A(p_i, p_j, c, a, b) = L^A$, where L^A refers to overall land in cultivation in the policy regime A.

Under the previous policy B the per parcel and total profits for land in cultivation are given by $\pi_i^B = (p_i + s)f_i(l_i) - c$ and $r_i^B = \pi_i^B L_i^B$, respectively. The optimal fertilizer intensity is a function of crop price, fertilizer price and price support, $l_i^B(p_i, s, c)$ and land allocation between crops depends on relative profitability as follows: $L_i^B = L_i^B(p_i, p_j, c, s)$. For the total amount of land in cultivation, it holds that $\sum_{i=1}^n L_i^B + E = L^B$, where L^B refers to overall amount of land in cultivation in the policy regime B (recall E is the mandatory fallow area).

The environmental quality is a function of the input use, the amount of cultivated arable land, and its allocation between the crops. Let function G represent the way the use of inputs in agriculture transform to environmental quality, nutrient runoff in our case. Then, drawing on the

above discussion environmental quality can be expressed for our three cases as a function of respective optimal choices:

$$G^0 = G^0 \left(\sum_{i=1}^n l_i^0(p_i, c) L_i^0(p_i, p_j, c) \right) \quad (1)$$

$$G^A = G^A \left(\sum_{i=1}^n l_i^A L_i^A(p_i, p_j, c, a, b) \right) \quad (2)$$

$$G^B = G^B \left(\sum_{i=1}^n l_i^B(p_i, s, c) L_i^B(p_i, p_j, c, s) + \varepsilon \bar{E} \right) \quad (3)$$

Recall, our aim is to assess the performance of the new agri-environmental policy A. Let \bar{G}^A be the announced environmental target of the new policy; in our case reduction in nutrient loads, while the observed environmental quality under this policy is G^A . Naturally, the difference between the goal and the actually measured nutrient runoff, $G^A - \bar{G}^A$, can be any sign and due to multiple reasons. The challenge of the counterfactual analysis is to explain this difference.

We can now use the above analysis to formulate our two counterfactuals, which recall, were the following. What would have happened to G^A if the land allocation between crops and green set-aside would not have changed from policy regime B? Second, what would have happened to G^A if no voluntary agri-environmental policy would have taken place when Finland joined the EU? Economic mechanisms present in (2) and (3) readily suggest how to formalize these counterfactuals (CF):

$$CF_1 = G^{CF_1} \left(\sum_{i=1}^n l_i^A L_i^B(p_i, p_j, c, s) + \varepsilon B \right) \quad (4)$$

$$CF_2 = G^{CF_2} \left(\sum_{i=1}^n l_i^0(p_i, c) L_i^A(p_i, p_j, c, a, b) \right) \quad (5)$$

Taking the difference $G^A - CF_1$ and $G^A - CF_2$ allows us evaluate the relative role of input use intensities and land use changes of the agri-environmental policy regime A. Counterfactual CF_1 allows us to define the unit effectiveness of the instruments in policy regime A and counterfactual CF_2 in turn defines the preventive impact of the policy A on nutrient loads. In the empirical part we also consider the role of some fine tunings of policy regime A on environmental impacts; they are our minor counterfactuals that are developed in a similar fashion as (4) and (5).

3. EMPIRICAL FRAMEWORK

3.1. Crop yield response to fertilizer

Per hectare crop yield is modelled as a function of nitrogen fertilization. By assumption, farmers use a compound fertilizer that contains nitrogen and phosphorus in fixed proportions and in the absence of constraints choose the application rate of fertilizer on the basis of yield response to nitrogen application. The crop yield function for spring wheat, barley, and oats is assumed to follow the Mitscherlich form,

$$y_i = \mu_i (1 - \sigma_i e^{-v_i N_i}) \quad (6)$$

where y_i is yield per hectare, N_i is nitrogen use per hectare, and μ_i , σ_i and v_i are parameters. These parameters are estimated by Bäckman *et al.* (1997) on the basis of Finnish field experiments. The yield function for rape, silage and hay is assumed to have the quadratic form

$$y_i = A_i + \chi_i N_i + \gamma_i N_i^2 \quad (7)$$

where y_i is yield per hectare, N_i is nitrogen use per hectare, and A_i , χ_i and γ_i are parameters. Parameters for rape have been estimated by Pietola *et al.* (1999) and parameters for silage and dry hay are based on Lehtonen (2001).

3.2. Optimal fertilizer use

Farmer's short-run restricted profits π^i are given by equation (8a) for spring wheat, barley, and oats and by equation (8b) for rape, silage and hay.

$$\pi^i = p_i \mu_i (1 - \sigma_i e^{-v_i N_i}) - c_i N_i \quad (8a)$$

$$\pi^i = p_i (A_i + \chi_i N_i + \gamma_i N_i^2) - c_i N_i \quad (8b)$$

Where π^i is farmers' per hectare profits, p_i is output price for a given crop (i) and c_i is nitrogen price for a given combined fertilizer (NPK). Optimal nitrogen application level can be solved by taking first-order conditions with respect to nitrogen application N and setting them to equal zero and then solving for optimal N .

3.3. Nutrient runoff

The modeling of nutrient runoff follows Lankoski *et al.* (2006) who modeled nitrogen and phosphorus runoff on the basis of Finnish data. In the case of phosphorus runoff two forms are distinguished: (i) dissolved reactive phosphorus (DRP) and (ii) particulate phosphorus (PP). As already noted, farmers use a compound fertilizer (NPK) and because these main nutrients are

in fixed proportions, nitrogen fertilizer intensity determines also the amount of phosphorus used. Part of this phosphorus is taken up by the crop, while the rest accumulates and builds up soil P. The concentration of dissolved phosphorus in surface runoff is found to depend linearly on the easily soluble soil P, and the runoff of particulate phosphorus depends on the rate of soil erosion and the P content of eroded soil material. The following nitrogen runoff function (Simmelsgaard 1991) is employed,

$$Z_N^i = \phi_i \exp(b_0 + bN_i), \quad (9)$$

where Z_N^i = nitrogen runoff at fertilizer intensity level N_i , kg/ha, ϕ_i = nitrogen runoff at average nitrogen use, $b_0 < 0$ and $b > 0$ are constants and N_i = nitrogen fertilization in relation to the normal fertilizer intensity for the crop, $0.5 \leq N \leq 1.5$. This runoff function represents nitrogen runoff generated by a nitrogen application rate of N_i per hectare and the parameter ϕ_i reflects differences in crops.

In the case of phosphorus, both dissolved and particulate runoff is modeled. Drawing on Finnish experiments (e.g. Saarela et al. 1995) it is assumed that 1 kg increase in soil phosphorus reserve increases the soil P status (i.e., ammonium acetate-extractable P) by 0.01 mg/l soil. Uusitalo and Jansson (2002) estimated the following linear equation between soil P and the concentration of dissolved phosphorus (DRP) in runoff: *water soluble P in runoff (mg/l) = 0.021*soil_P (mg/l soil) – 0.015 (mg/l)*. The surface runoff of potentially bioavailable particulate phosphorus is approximated from the rate of soil loss and the concentration of potentially bioavailable phosphorus in eroded soil material as follows: *potentially bioavailable particulate phosphorus PP (mg/kg eroded soil) = 250 * ln [soil_P (mg/l soil)]-150* (Uusitalo 2004). Thus, the parametric description of surface phosphorus runoff is given by

$$Z_{DRP}^i = \varpi_i [\psi_i (0.021(\Phi + 0.01 * P_i) - 0.015)] / 100 \quad (10a)$$

$$Z_{PP}^i = \Delta_i [\zeta_i \{250 \ln(\Phi + 0.01 * P_i) - 150\}] * 10^{-6} \quad (10b)$$

where ψ_i is runoff volume (mm), Φ is soil_P (common to all crops) and ζ_i is erosion kg/ha, and P_i is the phosphorus application rate. As in the case of nitrogen, the crop, soil textural class and field slope based differences in the runoff of dissolved and the potentially bioavailable particulate phosphorus are captured by parameters ϖ_i and Δ_i , respectively. Soil_P is fixed at 12.6 mg/l in 1995, 11.6 mg/l in 2001 and 10.6 mg/l in 2007 on the basis of the average for Finnish soil test samples taken on those respective years (MMM 2003 and Myyrä et al. 2005). fashion as (4) and (5).

4. DATA AND NUTRIENT RUNOFF UNDER FINNISH AGRI-ENVIRONMENTAL PROGRAMME

Following the theory, we develop the empirical counterparts for equations (4) and (5). Data is needed on land allocation, fertilizer application intensities and nutrient runoff. Data on the development of actual land-use and its allocation between the main crops is given by farm statistics; for our counterfactuals the actual land allocation represents revealed profit maximization by farmers. Another given data are fertilizer restrictions of the agri-environmental program during the years 1995-2007. Using annual crop and fertilizer prices we solve privately optimal fertilizer application level in the case where no agri-environmental program would have been implemented. Similarly nutrient runoff (both nitrogen and phosphorus) is estimated by using this modeling approach.

4.1. Data on land use and fertilizer application

Table 1 shows the land use between the main crops in 1994 (just before the beginning of current policy) and during the agri-environmental program. We present the land area of those crops that are external to the analysis under the land use class "Other". It includes crops, such as sugar beet, potatoes, and peas.

Table 1. Land use (ha) in 1994, 1995, 2001, and 2007 (Yearbook of Farm Statistics).

Land use	1994	1995	2001	2007
Wheat	77 600	88 100	115 400	167 900
Barley	505 700	516 200	547 200	550 100
Oats	334 300	329 300	422 700	361 500
Rape	67 200	85 300	73 100	90 200
Hay	257 900	287 100	157 500	103 100
Silage	268 400	300 900	380 900	438 100
Fallow	505 100	223 200	201 900	121 200
Other	285 700	316 300	293 200	423 200
Total	2 301 900	2 146 400	2 191 900	2 255 300

Table 1 shows that during the agri-environmental program periods (1995 – 2007) the total cultivated land increased by 2.1% from 1995 to 2001 and 5.1% from 1995 to 2007 in Finland. Also land allocation between different crops has changed much. Land allocation to wheat cultivation has increased from 4.8% to 8.6% between 1995 and 2007. The share of barley, oats and rape has remained quite stable while land allocated to hay has decreased from 15.7% to 5.3% and land allocated to silage increased from 16.4% to 22.6%. Thus, there has been a clear shift in land use towards more fertilizer intensive crops wheat and silage.

Table 2 presents the actual fertilization constraints and the fertilizer intensity under the hypothetical case, in which farmers optimize on the basis of market prices only and do not participate in the agri-environmental programme (labeled as "N private"). We do not distinguish between manure and chemical fertilizer application. Nitrogen applied can be a combination manure and chemical nitrogen or from chemical nitrogen only.

Table 2. Optimal nitrogen use intensity and nitrogen application constraint in agri-environmental program in 1995, 2001, and 2007.

Crop	1995		2001		2007	
	N private	N constraint	N private	N constraint	N private	N constraint
Wheat	155.3	100	140.4	100	158.8	120
Barley	122.1	90	112.1	90	123.3	100
Oats	97.9	90	92.2	90	94.2	100
Rape	156.3	100	147.3	100	137.2	110
Silage	161.6	180	199.3	180	255.6	240
Hay	128.0	90	133.4	90	176.6	100

Table 2 shows that for most crops the nitrogen application constraint has been binding throughout program years. For wheat, rape and hay the economically optimal application rate has been clearly higher than the constraint, while in the case of oats farmers' economic optimum and nitrogen use constraint are sufficiently close to each other.

4.2. Estimating nutrient runoff under the agri-environmental programme

We estimate the average per hectare nitrogen and phosphorus runoff by employing equations (9) – (10b) for each crop. Total nutrient runoff for each crop is obtained through multiplying the average runoff by the total land area allocated to each crop.

Table 3. Nitrogen, dissolved phosphorus (DRP) and particulate phosphorus (PP) runoff, kg/ha, under constrained fertilizer use intensity in 1995, 2001, and 2007.

Crop	1995			2001			2007		
	N	DRP	PP	N	DRP	PP	N	DRP	PP
Wheat	14.0	0.376	0.479	14.0	0.347	0.462	16.1	0.319	0.444
Barley	13.1	0.376	0.479	13.1	0.347	0.462	14.0	0.319	0.444
Oats	13.1	0.376	0.479	13.1	0.343	0.459	13.5	0.312	0.438
Rape	14.0	0.356	0.467	14.0	0.347	0.462	15.0	0.319	0.444
Silage	7.3	0.585	0.175	8.2	0.534	0.168	12.4	0.506	0.164
Hay	4.5	0.526	0.166	4.5	0.481	0.160	4.8	0.447	0.154
Fallow	5.4	0.461	0.280	5.4	0.407	0.280	5.4	0.373	0.280
Other	9.7	0.671	0.182	10.4	0.636	0.182	12.6	0.602	0.182

The most alarming feature in Table 3 is that due to the relaxation of the nitrogen application constraints, the last phase of the program witnesses the highest per hectare nitrogen runoff. In contrast to this per hectare runoff of particulate and dissolved phosphorus has diminished steadily over time due to the decrease in soil phosphorus. Hence, the total load of phosphorus per hectare has decreased.

Table 4 combines the observed land allocation between the crops and the estimated per hectare average nutrient runoff to produce total nitrogen and phosphorus runoff in tons in 1995,

2001, and 2007 under the Finnish Agri-Environmental Programme. In terms of our theory, Table 4 corresponds to equation (2).

Table 4. Total nitrogen and phosphorus runoff (tons) under constrained fertilizer use in 1995, 2001, and 2007.

Crop	1995		2001		2007	
	N	P	N	P	N	P
Wheat	1894	97	2481	121	4151	166
Barley	10347	568	10969	572	11825	544
Oats	6601	362	8473	438	7461	352
Rape	1834	91	1571	76	2079	89
Silage	3391	257	4883	301	8544	332
Hay	1962	224	1076	114	756	70
Fallow	1129	155	1021	130	613	74
Other	4693	303	4705	271	8184	376
Total	31 851	2056	35 180	2024	43 613	2003

The time path of the nitrogen load shows that despite all efforts, nitrogen runoff has increased during all three phases of the program. Recall that the per hectare nitrogen runoff was the same in 1995 and 2001 for all crops with one exception. Therefore, it is the shift of arable land to more fertilizer intensive crops, such as wheat and silage that increased nitrogen runoff during the second phase of the agri-environmental program. This tendency is re-enforced by the relaxed nitrogen constraint during the third phase. The development of phosphorus runoff follows a different path since the gradual decrease of soil phosphorus content decreases phosphorus runoff in the second and third phase of the program. Thus, nutrient loading evolves to opposite directions; the reason is that phosphorus application is roughly constant across crops, while nitrogen application varies considerably between crops. We can clearly conclude that the Finnish Agri-Environmental Programme has failed to reduce nitrogen loading to inland waters and the Baltic Sea.

5. THE IMPACTS OF THE FINNISH AGRI-ENVIRONMENTAL PROGRAMME ON NUTRIENT RUNOFF: A COUNTERFACTUAL APPROACH

We now turn to the counterfactual analysis. We first assess the effectiveness of the key policy instruments, fertilizer application constraints and mandatory buffer strips, in reducing nutrient runoff. This can be made by isolating the impacts of these intensive margin instruments from changes taking place in the extensive margin that is land allocation, as equation (4) suggest. This helps us to answer the question whether the key instruments have been effective or not. We follow then equation (5) to trace out how much the agri-environmental program has potentially offset nutrient runoff relative to the non-regulated market-based development.

5.1. The effectiveness of instruments targeting nutrient runoff

In Table 5 we examine what would have happened to nutrient loads if agricultural land use would not have changed over the years but would have stayed as it was either in 1994 or 1995. Taking the difference between this and the actual loads under the current policy regime, which is called the Baseline indicates the importance of controlling both entry-exit (total amount of arable land) and extensive (land allocation) margins in addition to intensive margin (fertilizer use intensity and buffer strips).

Table 5. Total nitrogen and phosphorus runoff (tons) under constrained fertilizer use in Baseline and in the case of total cultivated land and its allocation fixed to correspond to that of 1994 or 1995.

Land allocation	1995		2001		2007	
	N	P	N	P	N	P
Baseline	31 851	2056	35 180	2024	43 613	2003
Fixed 1995	31 851	2056	32 398	1938	36 519	1833
Fixed 1994	29 056	1982	29 499	1864	32 915	1764

Table 5 reveals that under either 1995 or 1994 land allocation total nitrogen load would have increased much less and phosphorus load decrease much more than they actually did. Under 1995 land allocation, decrease in phosphorus runoff by 2007 would have been 223 tons and under 1994 land allocation 218 tons, making 11.0% and 10.8% reduction of nutrient loads. Given that the phosphorus fertilization limit has been the same over all three periods, these figures indicate the true impact of phosphorus limits on loads that is purified from the changes in land allocation. Due to changes in land allocation, the actual decrease in phosphorus loads was 53 tons only, representing 2.6% reduction. The difference of these two figures, 165–170 tons, represents simply the increase in total phosphorus load due to the increased land area in cultivation.

During the years 1995 – 2007 nitrogen loads would have increased by 4668 tons (14.7%) under 1995 land allocation and by 3859 tons (13.3%) under 1994 land allocation. Contrast this to the actual increase in nitrogen loads that was 11 762 tons (36.9%). The increase in actual loads is explained by two factors: nitrogen fertilization constraints were relaxed in the third programme period (year 2007); and the amount of cultivated land has increased and more land is allocated to more nitrogen intensive crops. The relaxed nitrogen fertilization constraints increased nitrogen loads by 5753 tons in the baseline, representing 48.9% of the total load increase, relative to a situation where constraints would not have been relaxed. Consequently we can conclude that the impact of the increase in the amount of cultivated land and land allocation change towards more nitrogen intensive crops was 6009 tons, representing 51.1% of the load increase. Table 5 reveals how important role the total amount of land in cultivation (entry-exit margin) and land allocation between the crops (extensive margin) play in determining total nutrient loading from agriculture.

5.2. Preventive impact of the Finnish Agri-Environmental Programme

Let us ask next what would have happened without the program when the farmers are allowed to choose their fertilizer application rates freely on the basis of crop and fertilizer prices. This is our second counterfactual that is defined by equation (5). To answer the question, one must solve nutrient runoff per hectare under market solution and link this to land allocation between the crops.

Table 6 provides information on nutrient loads under both cases. The difference between the two figures indicates how much the Finnish Agri-Environmental Programme has prevented nutrients loads by its presence in each program phase.

Table 6. Total nitrogen and phosphorus runoff (tons) under constrained and free private fertilizer use in 1995, 2001, and 2007.

	1995		2001		2007	
	N	P	N	P	N	P
Constraint	31 851	2056	35 180	2024	43 613	2003
Private	39 083	2164	41 348	2105	51 418	2126
Difference	- 7232	- 108	- 6168	- 81	- 7805	- 123

As Table 6 reveals, the preventive effect of the Finnish Agri-Environmental Programme has been on average 7068 tons of nitrogen and 104 tons of phosphorus. Combining Table 6 with Table 5 allows us to make the following conclusion. The Programme has been somewhat successful in the intensive margin but this success has been outweighed by a failure to control both extensive margin (land allocation) as well as entry-exit margin (total amount of cultivated land). Consequently, nitrogen loads have increased and phosphorus loads have decreased less than expected.

6. COST-BENEFIT ANALYSIS OF WATER PROTECTION POLICY IN THE FINNISH AGRI-ENVIRONMENTAL PROGRAMME

We end the analysis by asking whether the benefits from water quality policy in the Finnish agri-environmental programme exceed the costs, or not. Costs of the policy is the overall amount of annual support payments to farmers targeted to water quality, while the benefits are given by the reduced nutrient runoff damages. Reductions in nutrient loads reduce damages both in inland waters and in the Baltic Sea. As the main goal is to improve the state of the Baltic Sea, we express phosphorus loads as nitrogen equivalent using the Redfield ratio 7.2. The Redfield ratio describes the optimum N/P ratio for the growth of phytoplankton, relevant for algal growth in sea waters. The marginal damage from nitrogen equivalents is assumed constant, so that the damage function is given by

$$d(Z^i) = R_n(N_i + 7.2P_i), \quad (11)$$

where R_n is the constant social marginal damage. Drawing on Gren (2001), the willingness to pay for nutrient load reduction in the Baltic Sea is set to be $R_n = \text{€ } 6.70/\text{kg}$ of N equivalent. Hence, this estimate provides social value of reductions in nutrient runoff. As regards social costs of nutrient runoff reduction we use the budget allocated to water protection measures in the agri-environmental program as a primary measure of social costs of nutrient runoff reduction.

Also, we report a more developed social net benefit estimate by including the policy related transaction costs (PRTC) to the social costs. Our estimate of policy related transaction costs of agri-environmental support is based on Ollikainen et al. (2008) who estimated that the PRTCs of Basic measure support (including fertilizer use constraints and buffer strips) are 1.5% of the total transfer. Finally, the most comprehensive social net-benefit estimate takes also into account the so-called marginal cost of taxation (marginal cost of public funds) as a measure of economic welfare losses due to raising government revenue with distortionary taxes (such as labor taxes). We employ 10% of the total transfer as our estimate of marginal cost of taxation.

Finally, for the purposes of comparison we provide a hypothetical case assuming that only compliance costs were compensated to farmer. A crude estimation of compliance costs of implementing the program is measured as forgone profits when a farmer complies with fertilizer use constraints and buffer strip establishment. Thus, profits forgone show the minimum level required for compensation payment.

Table 7 provides the three estimates of the social net benefits for the Finnish Agri-Environmental Scheme and the net benefits from hypothetical case where only compliance costs are compensated, that is there is no overcompensation.

Table 7. Social net-benefits of agri-environmental program

	1995	2001	2007
Social net benefits of the agri-environmental program			
N-eq reduction, tons	3343	1775	2885
Program outlays, million €	229.6	233.1	276.0
Value of damage reduction, million €	22.4	11.9	19.3
Net benefit, million €	-207.2	-221.2	-256.7
Transaction costs (TC), million €	3.4	3.4	4.0
Net benefit - TCs, million €	-210.5	-224.6	-260.7
Net benefit - TCs - MCT, million €	-233.5	-247.9	-288.3
Social net benefits if only compliance costs were compensated			
N-eq reduction, tons	8011	6754	8686
Profit forgone, million €	-24.1	-12.0	-27.7
Net benefits, million €	29.6	33.2	30.5

Table 7 reveals that the social net benefit of the programme is negative in every program period under all three net benefit measures. This clearly refers to overcompensation of farmers' compliance costs, that is part of the environmental support payments seem to entail farm income support. This can also be verified in the lower part of Table 7, where the direct compliance costs

(i.e. short-run profit forgone) are estimated. The estimated compliance costs vary over time as a function of crop prices and cultivation costs. These direct costs are quite low relative to the environmental payments, because they underestimate the long-run cost burden on farmers. However, even if these costs were doubled or tripled they would still remain relatively small in comparison to program outlays.

The lower part of Table 7 also illustrates what the benefits would have been if the Programme compensated only compliance costs. In this case entry of cultivated land due to Programme would have been zero and the reduction of nitrogen equivalents more than double. This would have made the social net benefits of the program positive with a wide margin of 30 million euros.

7. CONCLUDING REMARKS

We developed a theoretical framework using the interlinkages between the behaviour of agents and the response of environmental systems to the economic decisions. We applied our model to agricultural water protection policy of the Finnish Agri-Environmental Programme, which aims at reducing nutrient runoff from arable lands to the Baltic Sea. Counterfactual analysis allowed us to examine both the unit effectiveness of the measures included in the Programme and its preventive impact.

We find that the Finnish agri-environmental programme has failed to achieve its goals: nitrogen loads have increased and phosphorus loads have decreased only slightly. Our counterfactuals help to trace out the mechanisms leading to this failure. This is not to say that the Finnish Agri-Environmental Programme would have failed entirely. We demonstrated that the loads of both nutrients would have been much higher if no policies would have been implemented. The social cost-benefit analysis of the program showed, nevertheless, strongly negative net benefits. Thus, there is a lot of scope for improving the agri-environmental water protection policies.

ACKNOWLEDGMENT

This paper is a part of the project "Cost-efficient instruments for the Protection of the Baltic Sea" funded by the Ministry of the Environment and Ministry of Agriculture and Forestry. Funding is gratefully acknowledged. Ollikainen also acknowledges funding from Yrjö Jahnsson Foundation.

REFERENCES

- Bäckman, S.T., Vermeulen, S. and Taavitsainen, V.-M. (1997). Long-term fertilizer field trials: comparison of three mathematical response models. *Agricultural and Food Science in Finland* 6: 151-160.
- Gren, I-M., 2001. International versus national action against pollution at the Baltic Sea. *Environmental and Resource Economics* 20, 41-59.
- HELCOM (2010). Atlas of the Baltic Sea. Helsinki Commission, Helsinki, Finland.
- Lankoski, J., Ollikainen M. and Uusitalo, P. (2006). No-till technology: benefits to farmers and the environment? Theoretical analysis and application to Finnish agriculture. *European Review of Agricultural Economics* 33: 193-221.

Ancona - 122nd EAAE Seminar
"Evidence-Based Agricultural and Rural Policy Making"

- Lehtonen, H. (2001). Principles, structure and application of dynamic regional sector model of Finnish agriculture. PhD thesis, Systems Analysis Laboratory, Helsinki University of Technology. Agrifood Research Finland, Economic Research (MTTL). Publ. 98. 264 p.
- MMM (2003). Horisontaalinen maaseudun kehittämisohjelma. EU:n yhteisen maatalouspolitiikan liitännäistoimenpiteet. Manner-Suomi. Helsinki. 229 s.
- Myyrä, S. Ketola, E., Yli-Halla, M. and Pietola, K. (2005). Land improvements under land Tenure insecurity: the case of pH and phosphate in Finland. *Land Economics* 81: 557-569.
- Ollikainen, M., Lankoski, J. and Nuutinen, S. (2008). Policy-related transaction costs of agricultural policies in Finland. *Agricultural and Food Science* 17: 193-209.
- Pietola, L., Tanni, R. & Elonen, P. (1999). Responses of yield and N use of spring sown crops to N fertilization, with special reference to the use of plant growth regulators. *Agricultural and Food Science in Finland* 8:423-440.
- Puustinen M, E. Turtola, M. Kukkonen, J. Koskiahio, J. Linjama, R. Niinioja, S. Tattari (2010). VIHMA – a tool for allocation of measures to control erosion and nutrient loading from Finnish agricultural catchments. *Agriculture, Ecosystems and Environment* 138: 306-317.
- Saarela, I., Järvi, A., Hakkola, H., and Rinne, K. (1995). Fosforilannoituksen porraskokeet 1977-1994. Vuosittain annetun fosforimäärän vaikutus maan viljavuuteen ja peltokasvien satoon monivuotisissa kenttäkokeissa. In English: Phosphorus fertilizer trials, 1977-1994: effects of the rate of annual phosphorus application on soil fertility and yields of field crops in long-term field experiments). Maatalouden tutkimuskeskus, Tiedote 16/95.
- Simmelsgaard, S. (1991). *Estimation of nitrogen leakage functions – Nitrogen leakage as a function of nitrogen applications for different crops on sand and clay soils.* In: Rude, S. (ed.). Nitrogen fertilizers in Danish Agriculture - present and future application and leaching, Report 62, Institute of Agricultural Economics, Copenhagen, 135-150 (in Danish: Kvaelstofgødning i landbruget - behov og udvasking nu og I fremtiden).
- Uusitalo, R. (2004). Potential bioavailability of particular phosphorus in runoff from arable clayey soils. Agrifood Research Reports 53, Doctoral Dissertation, MTT Agrifood Research Finland. Jokioinen: MTT Agrifood Research Finland.
- Uusitalo, R. and Jansson, H. (2002). Dissolved reactive phosphorus in runoff assessed by soil extraction with an acetate buffer. *Agricultural and Food Science in Finland* 11: 343-353.
- Vehkasalo, V. (1999). Ympäristötuen yhteiskunnallinen kannattavuus. In: Maatalouden ympäristöohjelma 1995-1999:n taloudellinen analyysi. Agricultural Economics Research Institute, Publications 90: 42-77.
- Ympäristöministeriö (2000). Vesien suojelun toimenpideohjelma vuoteen 2005. Suomen ympäristö 402. Helsinki. 47 s.